arXiv:1005.0681v2 [math.GR] 10 Mar 2011

CLASSIFICATION OF EQUIVARIANT VECTOR BUNDLES OVER TWO-SPHERE

MIN KYU KIM

Abstract. We classify equivariant topological complex vector bundles over two-sphere under a compact Lie group (not necessarily effective) action. It is shown that nonequivariant Chern classes and isotropy representations at (at most) three points are sufficient to classify equivariant vector bundles except a few cases. To do it, we calculate homotopy of the set of equivariant clutching maps. In other papers, we also give classifications over two-torus, real projective plane, Klein bottle.

1. Introduction

In topology, nonequivariant complex vector bundles can be classified just by calculating their Chern classes under a dimension condition [P]. However, there is no such general result on equivariant vector bundles. Instead, a few results on extreme cases are known. Let us mention four of them. Let a compact Lie group G act on a topological space X.

- If G is trivial, then G-vector bundles over X are just nonequivariant vector bundles, and are classified by their Chern classes in $H^*(X)$.
- In [At, Proposition 1.6.1], [S, p. 132], G-vector bundles over a free G-space X are in one-to-one correspondence with nonequivariant vector bundles over X/G, and are classified by Chern classes in $H^*(X/G)$.
- For a closed subgroup H of G and any point x in $X = G/H$, G-vector bundles over X are classified by their isotropy representations at x which are contained in $\mathrm{Rep}(G_x)$. For this, see [S, p. 130], [B, Proposition II.3.2]. In [CKMS], G-vector bundles over $X = S^1$ are classified by their isotropy representations at (at most) two points.

Since invariants live in $H^*(X)$, $H^*(X/G)$, $\mathrm{Rep}(G_x)$, these results might be considered to have three different types. In this paper, we classify equivariant topological complex vector bundles over two-sphere under a compact Lie group (not necessarily effective) action. Readers will see those different types at once in it. By developing ideas and machineries of it, classification over two-torus is given in [Ki1] which shows that our method is not restrictive. Classifications over real projective plane and Klein bottle are also obtained as corollaries in [Ki2], [Ki3]. After these, classification of equivariant holomorphic vector bundles over Riemann sphere under a holomorphic complex reductive group (not necessarily effective) action will follow which is an equivariant version of Grothendieck's Theorem for holomorphic vector bundles on Riemann sphere [G], and is the motivation for the paper.

To state main results, we need introduce several notations. It is well-known that a topological action on S^2 by a compact Lie group is conjugate to a linear action [Ko, Theorem 1.2], [CK]. Let a compact Lie group G act linearly (not necessarily effectively) on the unit sphere S^2 in $\mathbb{R}^3$ through a representation $\rho : G \to \mathrm{O}(3)$. Let $\mathrm{Vect}_G(S^2)$ be the set of isomorphism classes of topological complex G-vector

---

2000 *Mathematics Subject Classification.* Primary 57S25, 55P91 ; Secondary 20C99.

Key words and phrases. equivariant vector bundle, equivariant homotopy, representation.

bundles over S^2 with the given G-action. For a bundle E in $\mathrm{Vect}_G(S^2)$ and a point x in S^2, denote by E_x the isotropy G_x-representation on the fiber at x. It is needed to decompose $\mathrm{Vect}_G(S^2)$ as the sum of more simple subsemigroups. For this, some terminologies are introduced. Put $H = \ker \rho$, i.e. the kernel of the G-action on S^2. Let $\mathrm{Irr}(H)$ be the set of characters of irreducible complex H-representations which has a G-action defined as

$$(g \cdot \chi)(h) = \chi(g^{-1}hg)$$

for $\chi \in \mathrm{Irr}(H)$, $g \in G$, $h \in H$. Sometimes, we also use the notation $\mathrm{Irr}(H)$ to denote the set of isomorphism classes of irreducible complex H-representations themselves. For $\chi \in \mathrm{Irr}(H)$, an H-representation is called χ-isotypical if its character is a multiple of χ. We slightly generalize this concept. For $\chi \in \mathrm{Irr}(H)$ and a compact Lie group K satisfying $H \lhd K < G$ and $K \cdot \chi = \chi$, a K-representation W is called χ-*isotypical* if $\mathrm{res}_H^K W$ is χ-isotypical, and denote by $\mathrm{Vect}_K(S^2, \chi)$ the set

$$\left\{ \ [E] \in \mathrm{Vect}_K(S^2) \ \middle| \ E_x \text{ is } \chi\text{-isotypical for each } x \in S^2 \ \right\}$$

where S^2 delivers the restricted K-action. In [CKMS], the (*isotypical*) *decomposition* of a G-bundle is defined, and from this a semigroup isomorphism is constructed to satisfy

$$\mathrm{Vect}_G(S^2) \cong \bigoplus_{\chi \in \mathrm{Irr}(H)/G} \mathrm{Vect}_{G_\chi}(S^2, \chi)$$

where G_χ is the isotropy subgroup of G at χ. As a result, our classification is reduced to $\mathrm{Vect}_{G_\chi}(S^2, \chi)$ for each $\chi \in \mathrm{Irr}(H)$. Details are found in [CKMS, Section 2].

The classification of $\mathrm{Vect}_{G_\chi}(S^2, \chi)$ is highly dependent on the G_χ-action on the base space S^2, i.e. on the image $\rho(G_\chi)$, and classification is actually given case by case according to $\rho(G_\chi)$. So, we need to describe $\rho(G_\chi)$ in a moderate way. For this, we would list all possible $\rho(G_\chi)$'s up to conjugacy, and then assign an equivariant simplicial complex structure of S^2 to each finite $\rho(G_\chi)$. Cases of nonzero-dimensional $\rho(G_\chi)$ are relatively simple and separately dealt with as special cases. First, let us define some polyhedra. Let P_m for $m \geq 3$ be the regular m-gon on xy-plane in $\mathbb{R}^3$ whose center is the origin and one of whose vertices is $(1, 0, 0)$. Then,

(1) $|\mathcal{K}_m|$ is defined as the boundary of the convex hull of P_m, $S = (0, 0, -1)$, $N = (0, 0, 1)$,
(2) $|\mathcal{K}_{\mathrm{T}}|$ is defined as the tetrahedron which is the boundary of the convex hull of four points $(\frac{1}{3}, \frac{1}{3}, \frac{1}{3})$, $(-\frac{1}{3}, -\frac{1}{3}, \frac{1}{3})$, $(-\frac{1}{3}, \frac{1}{3}, -\frac{1}{3})$, $(\frac{1}{3}, -\frac{1}{3}, -\frac{1}{3})$, and which is inscribed to $|\mathcal{K}_4|$,
(3) $|\mathcal{K}_{\mathrm{I}}|$ is defined as an icosahedron which has the origin as the center.

With these, denote natural simplicial complex structures on $|\mathcal{K}_m|$, $|\mathcal{K}_{\mathrm{T}}|$, $|\mathcal{K}_{\mathrm{I}}|$ by $\mathcal{K}_m$, $\mathcal{K}_{\mathrm{T}}$, $\mathcal{K}_{\mathrm{I}}$, respectively. Then, it is well-known that each closed subgroup of SO(3) is conjugate to one of the following subgroups [R, Theorem 11]:

(1) $\mathbb{Z}_n$ generated by the rotation a_n through the angle $2\pi/n$ around z-axis,
(2) D_n generated by a_n and the rotation b through the angle π around x-axis,
(3) the tetrahedral group T which is the rotation group of $|\mathcal{K}_{\mathrm{T}}|$,
(4) the octahedral group O which is the rotation group of $|\mathcal{K}_4|$,
(5) the icosahedral group I which is the rotation group of $|\mathcal{K}_{\mathrm{I}}|$,
(6) SO(2) which is the set of rotations around z-axis,
(7) O(2) which is defined as $\langle \mathrm{SO}(2), b \rangle$,
(8) SO(3) itself.

Note that $\mathrm{T} \subset \mathrm{O}$, and pick an element o_0 of $\mathrm{O} \setminus \mathrm{T}$ so that $\mathrm{O} = \langle \mathrm{T}, o_0 \rangle$. And, denote by Z the centralizer $\{\mathrm{id}, -\mathrm{id}\}$ of O(3). In Section 2, it is shown that each closed subgroup of O(3) is conjugate to an R-entry of Table 1.1 (there is no literature on this as far as the author knows). In the table, the notation $\times$ means internal direct product of two subgroups in O(3). Henceforward, it is assumed that $\rho(G_\chi) = R$ for some R. Let O(3) and their subgroups act naturally on $\mathbb{R}^3$. To each finite R, we assign a simplicial complex $\mathcal{K}_R$ of Table 1.1 where $\mathcal{K}_{\mathrm{O}}$ is defined in the below. Each $|\mathcal{K}_R|$ is invariant under the natural R-action on $\mathbb{R}^3$ as shown in Section 3, and is also invariant under the G_χ-action defined through ρ. So, we assume that $|\mathcal{K}_R|$ delivers the G_χ-action, and that $\mathcal{K}_R$ delivers the G_χ-action inherited from it. Henceforward, we consider $|\mathcal{K}_R|$ as the base space instead of the usual two-sphere S^2 when R is finite.

R	$\mathcal{K}_R$	D_R	d^{-1}
D_n, $n > 1$	$\mathcal{K}_n$	$[v^0, b(e^0)]$	S
$\mathbb{Z}_n$	$\mathcal{K}_n$	$\lvert e^0 \rvert$	S
$\mathrm{D}_n \times Z$, odd n	$\mathcal{K}_{2n}$	$[v^0, b(e^0)]$	S
$\langle a_n, -b \rangle$, odd n	$\mathcal{K}_{2n}$	$[b(e^0), v^1] \cup [v^1, b(e^1)]$	S
$\mathbb{Z}_n \times Z$, odd n	$\mathcal{K}_{2n}$	$\lvert e^0 \rvert$	S
$\mathrm{D}_n \times Z$, even n	$\mathcal{K}_n$	$[v^0, b(e^0)]$	S
$\langle a_n, -b \rangle$, even n	$\mathcal{K}_n$	$[v^0, b(e^0)]$	S
$\langle -a_n, b \rangle$, odd $n/2$, $n > 2$	$\mathcal{K}_{n/2}$	$\lvert e^0 \rvert$	S
$\langle -a_n, b \rangle$, even $n/2$	$\mathcal{K}_n$	$[v^0, b(e^0)]$	S
$\langle -a_n, -b \rangle$, odd $n/2$, $n > 2$	$\mathcal{K}_n$	$[b(e^0), v^1] \cup [v^1, b(e^1)]$	S
$\langle -a_n, -b \rangle$, even $n/2$	$\mathcal{K}_n$	$[v^0, b(e^0)]$	S
$\mathbb{Z}_n \times Z$, even n, $n > 2$	$\mathcal{K}_n$	$\lvert e^0 \rvert$	S
$\langle -a_n \rangle$, odd $n/2$, $n > 2$	$\mathcal{K}_{n/2}$	$\lvert e^0 \rvert$	S
$\langle -a_n \rangle$, even $n/2$	$\mathcal{K}_n$	$\lvert e^0 \rvert$	S
T	$\mathcal{K}_{\mathrm{T}}$	$[v^0, b(e^0)]$	$b(f^{-1})$
O	$\mathcal{K}_{\mathrm{O}}$	$[v^0, b(e^0)]$	$b(f^{-1})$
I	$\mathcal{K}_{\mathrm{I}}$	$[v^0, b(e^0)]$	$b(f^{-1})$
$\langle \mathrm{T}, -o_0 \rangle$	$\mathcal{K}_{\mathrm{T}}$	$[v^0, b(e^0)]$	$b(f^{-1})$
$\mathrm{T} \times Z$	$\mathcal{K}_{\mathrm{O}}$	$\lvert e^0 \rvert$	$b(f^{-1})$
$\mathrm{O} \times Z$	$\mathcal{K}_{\mathrm{O}}$	$[v^0, b(e^0)]$	$b(f^{-1})$
$\mathrm{I} \times Z$	$\mathcal{K}_{\mathrm{I}}$	$[v^0, b(e^0)]$	$b(f^{-1})$
O(3)		$\{v^0\}$	v^0
$\mathrm{O}(2) \times Z$		$\{v^0\}$	S
$\langle \mathrm{SO}(2), -b \rangle$		$\{v^0\}$	S
$\langle \mathrm{SO}(2), -a_2 \rangle$		$\{v^0\}$	S
SO(3)		$\{v^0\}$	v^0
O(2)		$\{v^0\}$	S
SO(2)		$\{v^0\}$	S

TABLE 1.1. $\mathcal{K}_R$, D_R, d^{-1} for closed subgroup R

In dealing with equivariant vector bundles over two-sphere, we need to consider isotropy representations at a few points (at most three points) of $|\mathcal{K}_R|$. To specify those points, we introduce some more notations. When $m \geq 3$, denote by v^i the vertex $\exp\left(\frac{2\pi i\sqrt{-1}}{m}\right) \in \mathbb{R}^2$ of $\mathcal{K}_m$, and by e^i the edge of $\mathcal{K}_m$ connecting v^i and v^{i+1} for $i \in \mathbb{Z}_m$. These notations are illustrated in Figure 10.1. When we use the notation $\mathbb{Z}_m$ to denote an index set, it is just the group $\mathbb{Z}/m\mathbb{Z}$ of integers modulo m. In Section 3, $\mathcal{K}_m$, v^i, e^i for $m = 1$, 2 are also defined. We would define similar notations for $\mathcal{K}_{\mathrm{T}}$, $\mathcal{K}_{\mathrm{I}}$. For $\mathcal{K}_{\mathrm{T}}$, $\mathcal{K}_{\mathrm{I}}$, pick two adjacent faces in each case, and call them f^{-1} and f^0. And, label vertices of f^{-1} as v^i for $i \in \mathbb{Z}_3$ to satisfy

(1) v^0, v^1, v^2 are arranged in the clockwise way around f^{-1},
(2) v^0, v^1 are contained in $f^{-1} \cap f^0$.

For $i \in \mathbb{Z}_3$, denote by e^i be the edge connecting v^i and v^{i+1}, and by f^i the face which is adjacent to f^{-1} and contains the edge e^i. We distinguish the superscripts -1 and

2 only for f^i, i.e. $f^{-1} \neq f^2$ in contrast to $v^{-1} = v^2$, $e^{-1} = e^2$. These notations are illustrated in Figure 4.1.(a). Here, we define one more simplicial complex denoted by $\mathcal{K}_{\mathrm{O}}$ which is the same simplicial complex with $\mathcal{K}_4$ but has the same convention of notations v^i, e^i, f^{-1}, f^i with $\mathcal{K}_{\mathrm{T}}$, $\mathcal{K}_{\mathrm{I}}$. Also, put $|\mathcal{K}_{\mathrm{O}}| = |\mathcal{K}_4|$. These notations are illustrated in Figure 4.2.(a). With these notations, we explain for D_R-entry of Table 1.1. To each finite R, we assign a path D_R (called the (closed) *one-dimensional fundamental domain*) in $|\mathcal{K}_R|$ which is listed in the third column of Table 1.1 where $b(\sigma)$ is the barycenter of σ for any simplex σ and $[x, y]$ is the shortest path in $|\mathcal{K}|$ for any simplicial complex $\mathcal{K}$ and two points x, y in $|\mathcal{K}|$. And, let d^0 and d^1 be boundary points of D_R such that d^0 is nearer to v^0 than d^1. Here, we define one more point d^{-1} for each finite R which is listed in the fourth column in Table 1.1. If R is one-dimensional, then denote by D_R the one point set $\{v^0 = (1, 0, 0)\}$, and let d^{-1}, d^0, d^1 be equal to S, v^0, v^0, respectively. Similarly, if R is three-dimensional, then denote by D_R the one point set $\{v^0 = (1, 0, 0)\}$, and let d^{-1}, d^0, d^1 be all equal to v^0. So far, we have defined d^{-1}, d^0, d^1 for each R. Then, $\{S, N\}$ or $\{d^{-1}, d^0, d^1\}$ are wanted points according to R, and we will consider the restriction $E|_{\{S,\ N\}}$ or $E|_{\{d^{-1},\ d^0,\ d^1\}}$ for each E in $\mathrm{Vect}_{G_\chi}(S^2, \chi)$. Define the following semigroup which will be shown to be equal to the set of all the restrictions:

Definition 1.1. For $\chi \in \mathrm{Irr}(H)$, assume that $\rho(G_\chi) = R$ for some R of Table 1.1.

(1) If $R = \mathbb{Z}_n$, $\langle a_n, -b\rangle$, $\mathrm{SO}(2)$, $\langle \mathrm{SO}(2), -b\rangle$, then let $A_{G_\chi}(S^2, \chi)$ be the semigroup of pairs (W_S, W_N) in $\mathrm{Rep}(G_\chi)^2$ satisfying
   i) W_S is χ-isotypical,
   ii) $\mathrm{res}^{G_\chi}_{(G_\chi)_x} W_S \cong \mathrm{res}^{G_\chi}_{(G_\chi)_x} W_N$ for $x = d^0, d^1$.

   And, let $p_{\mathrm{vect}} : \mathrm{Vect}_{G_\chi}(S^2, \chi) \to A_{G_\chi}(S^2, \chi)$ be the semigroup homomorphism defined as $[E] \mapsto (E_S, E_N)$.

(2) Otherwise, let $A_{G_\chi}(S^2, \chi)$ be the semigroup of triples $(W_{d^{-1}}, W_{d^0}, W_{d^1})$ in $\mathrm{Rep}\left((G_\chi)_{d^{-1}}\right) \times \mathrm{Rep}\left((G_\chi)_{d^0}\right) \times \mathrm{Rep}\left((G_\chi)_{d^1}\right)$ satisfying
   i) $W_{d^{-1}}$ is χ-isotypical,
   ii) $W_{d^1} \cong {}^g W_{d^0}$ if there exists $g \in G_\chi$ such that $gd^0 = d^1$,
   iii) for any two points x, x' of three points d^{-1}, d^0, d^1,

$$\mathrm{res}^{(G_\chi)_x}_{(G_\chi)_x \cap (G_\chi)_{x'}} W_x \cong \mathrm{res}^{(G_\chi)_{x'}}_{(G_\chi)_x \cap (G_\chi)_{x'}} W_{x'}.$$

   And, let $p_{\mathrm{vect}} : \mathrm{Vect}_{G_\chi}(S^2, \chi) \to A_{G_\chi}(S^2, \chi)$ be the semigroup homomorphism defined as $[E] \mapsto (E_{d^{-1}}, E_{d^0}, E_{d^1})$.

See Definition 8.1 for the superscript g. Well-definedness of p_{vect} is proved in Lemma 8.2 and Lemma 11.1. Put $I = \{0, 1\}$ and $I^+ = \{-1, 0, 1\}$. And, denote a triple $(W_{d^{-1}}, W_{d^0}, W_{d^1})$ by $(W_{d^i})_{i \in I^+}$. Especially, if $\rho(G_\chi) = \mathrm{SO}(3)$, $\mathrm{O}(3)$, then $A_{G_\chi}(S^2, \chi)$ is equal to the set

$$\left\{ (W_{d^i})_{i \in I^+} \in \mathrm{Rep}\left((G_\chi)_{v_0}\right)^3 \;\middle|\; W_{d^i}\text{'s are the same } \chi\text{-isotypical representation}\right\},$$

and p_{vect} becomes an isomorphism by classification over homogeneous space.

Now, we can state main results. Let $c_1 : \mathrm{Vect}_{G_\chi}(S^2, \chi) \to H^2(S^2)$ be the map defined as $[E] \mapsto c_1(E)$. Denote by l_R the number $|G_\chi|/|G_\chi(D_R)|$ where $G_\chi(D_R)$ is the subgroup of G_χ preserving D_R.

Theorem A. *Assume that* $\rho(G_\chi) = R$ *is equal to one of* $\mathbb{Z}_n$, D_n, T, O, I. *Then,* p_{vect} *is surjective, and*

$$p_{\mathrm{vect}} \times c_1 : \mathrm{Vect}_{G_\chi}(S^2, \chi) \to A_{G_\chi}(S^2, \chi) \times H^2(S^2)$$

is injective. More precisely, bundles in $p_{\mathrm{vect}}^{-1}(\mathbf{W})$ *for each* $\mathbf{W}$ *in* $A_{G_\chi}(S^2,\chi)$ *have all different Chern classes, and* $c_1\big(p_{\mathrm{vect}}^{-1}(\mathbf{W})\big)$ *is equal to* $\{\ \chi(\mathrm{id})\big(l_R k+k_0\big)\ \big|\ k\in\mathbb{Z}\ \}$ *where* k_0 *is dependent on* $\mathbf{W}$.

Theorem B. *Assume that* $\rho(G_\chi)=R$ *is equal to one of* $\mathbb{Z}_n\times Z$ *with odd* n, $\langle -a_n\rangle$ *with even* $n/2$. *For each* $\mathbf{W}$ *in* $A_{G_\chi}(S^2,\chi)$, *its preimage* $p_{\mathrm{vect}}^{-1}(\mathbf{W})$ *has exactly two elements which have the same Chern class. Also,* $[E\oplus E_1]\neq[E\oplus E_2]$ *for any bundles* E, E_1, E_2 *in* $\mathrm{Vect}_{G_\chi}(S^2,\chi)$ *such that* $p_{\mathrm{vect}}([E_1])=p_{\mathrm{vect}}([E_2])$ *and* $[E_1]\neq[E_2]$.

Theorem C. *Assume that* $\rho(G_\chi)=R$ *for some* R *of Table 1.1 which is not equal to any group appearing in the above two theorems. Then,* p_{vect} *is an isomorphism.*

Theorem C says that isotropy representations at some points classify equivariant vector bundles for some R's. Comparing to this, Theorem A, B might be unsatisfactory to some readers because statements on Chern classes, especially k_0, are not so concrete. To handle this, we show the following:

Theorem D. *Assume that* $\rho(G_\chi)=R$ *for some* R *appearing in Theorem A, B. Then,* $\mathrm{Vect}_{G_\chi}(S^2,\chi)$ *is isomorphic to* $\mathrm{Vect}_R(S^2)$ *as semigroups, and* $\mathrm{Vect}_R(S^2)$ *is generated by line bundles. Also,* $A_R(S^2,\mathrm{id})$ *is generated by all the elements with one-dimensional entries. The number of such elements is equal to*

$$\begin{cases} |R_S|\times|R_N| & \text{if } R=\mathbb{Z}_n,\\ |R_{d^{-1}}|\times|R_{d^0}|\times|R_{d^1}| & \text{if } R\neq\mathbb{Z}_n \end{cases}$$

where R_x *is the isotropy subgroup at* $x\in S^2$ *and we denote simply by* id *the trivial character of the trivial group.*

In Section 13, we calculate Chern classes of line bundles in $\mathrm{Vect}_R(S^2)$, and from this we obtain k_0. Also in Section 13, we explain for the reason why we prove the isomorphism of Theorem D only for R's appearing in Theorem A, B.

This paper is organized as follows. In Section 2, we list all closed subgroups of O(3) up to conjugacy. In Section 3, we give an equivariant simplicial complex structure $\mathcal{K}_R$ on S^2 according to finite R, and investigate equivariance of S^2 by calculating isotropy subgroups at vertices and barycenters of $|\mathcal{K}_R|$. Section 4∼12 are divided into three parts. In Section 4∼9, we first deal with cases when $\mathcal{K}_R=\mathcal{K}_{\mathrm{T}}$, $\mathcal{K}_{\mathrm{O}}$, $\mathcal{K}_{\mathrm{I}}$. In Section 4, we introduce a new simplicial complex denoted by $\bar{\mathcal{K}}_R$ which is just a disjoint union of faces of $\mathcal{K}_R$, and consider $|\mathcal{K}_R|$ as a quotient space of the underlying space $|\bar{\mathcal{K}}_R|$. Also, we consider an equivariant vector bundle over S^2 as an equivariant clutching construction of an equivariant vector bundle over $|\bar{\mathcal{K}}_R|$. For this, we define equivariant clutching map. In Section 5, we investigate relations among $\mathrm{Vect}_{G_\chi}(S^2,\chi)$, $A_{G_\chi}(S^2,\chi)$, and homotopy of the set of equivariant clutching maps. From these relations, it is shown that our classification in most cases is obtained by calculation of the homotopy. In Section 6, we develop our machinery called equivariant pointwise gluing which glues an equivariant vector bundle over a finite set through a map called equivariant pointwise clutching map. In calculation of the homotopy, equivariant pointwise clutching map plays a key role because an equivariant clutching map can be considered as a continuous collection of equivariant pointwise clutching maps. Here, the concept of representation extension enters with which equivariant pointwise gluing is described in the language of representation theory. In this way, the homotopy is techniquely related to $A_{G_\chi}(S^2,\chi)$, and calculation of it becomes reduced to calculation of a relative (nonequivariant) homotopy. So, we prove a lemma on relative homotopy in Section 7. In Section 8, we prove technical lemmas needed in dealing with equivariant clutching maps through equivariant pointwise clutching maps. In Section 9, we prove main theorems for cases when $\mathcal{K}_R=\mathcal{K}_{\mathrm{T}}$, $\mathcal{K}_{\mathrm{O}}$, $\mathcal{K}_{\mathrm{I}}$. In Section 10∼11, we second deal with cases when

$\mathcal{K}_R = \mathcal{K}_m$ for some $m \in \mathbb{N}$. In Section 10, we rewrite what we have done in Section 4, 5, 8 to be in accordance with $\mathcal{K}_m$. In Section 11, we prove main theorems for the cases when $\mathcal{K}_R = \mathcal{K}_m$. In Section 12, we third prove Theorem C for cases when R is one-dimensional. In Section 13, we prove Theorem D. Section 14 is the appendix on representation extension.

2. Closed subgroups of O(3)

In this section, we list all closed subgroups of O(3) up to conjugacy. In this section, the notation × is internal direct product of two subgroups in O(3). Since Z is the centralizer of O(3), $\mathrm{O}(3) = \mathrm{SO}(3) \times Z$ and this gives the exact sequence

$$0 \to Z \hookrightarrow \mathrm{O}(3) \xrightarrow{\mathrm{pr}} \mathrm{SO}(3) \to 0$$

where $\mathrm{pr}\,|_{\mathrm{SO}(3)}$ is the identity map. First, we deal with finite subgroups.

Proposition 2.1. *Let R be a finite subgroup of* O(3) *such that $R \not\subset \mathrm{SO}(3)$.*

1. *If $\mathrm{pr}(R)$ is conjugate to $\mathbb{Z}_n$, then*
$$R \text{ is conjugate to } \begin{cases} \mathbb{Z}_n \times Z & \text{if } n \text{ is odd,} \\ \mathbb{Z}_n \times Z \text{ or } \langle -a_n \rangle & \text{if } n \text{ is even.} \end{cases}$$
2. *If $\mathrm{pr}(R)$ is conjugate to D_n, then*
$$R \text{ is conjugate to } \begin{cases} \mathrm{D}_n \times Z \text{ or } \langle a_n, -b \rangle & \text{if } n \text{ is odd,} \\ \mathrm{D}_n \times Z,\ \langle a_n, -b \rangle, & \\ \langle -a_n, b \rangle, \text{ or } \langle -a_n, -b \rangle & \text{if } n \text{ is even.} \end{cases}$$
3. *If $\mathrm{pr}(R)$ is conjugate to* T, *then R is conjugate to* $\mathrm{T} \times Z$.
4. *If $\mathrm{pr}(R)$ is conjugate to* O, *then R is conjugate to* $\mathrm{O} \times Z$ *or* $\langle \mathrm{T}, -o_0 \rangle$.
5. *If $\mathrm{pr}(R)$ is conjugate to* I, *then R is conjugate to* $\mathrm{I} \times Z$.

Proof. We may assume that $\mathrm{pr}(R)$ is equal to one of $\mathbb{Z}_n$, D_n, T, O, I. Denote by R_{rot} the subgroup $R \cap \mathrm{SO}(3)$. Then, R_{rot} is an index two subgroup of R. Denoting $\mathrm{pr}(R)$ by K, $R_{\mathrm{rot}} \subset K$ because $\mathrm{pr}\,|_{\mathrm{SO}(3)}$ is the identity map. Also, since the preimage $\mathrm{pr}^{-1}(g)$ of $g \in \mathrm{SO}(3)$ is equal to $\{g, -g\}$, it is obtained that $R \subset K \times Z$. Here are two possibilities. First, if $Z = \ker \mathrm{pr} \subset R$, then $|R| = 2|K|$ so that

$$R = K \times Z \tag{2.1}$$

from $R \subset K \times Z$. Second, if $Z \not\subset R$, then $\mathrm{pr}\,|_R$ is injective and $|R| = |K|$ so that

$$|K| = 2|R_{\mathrm{rot}}| \tag{2.2}$$

because R_{rot} is an index two subgroup of R. For an element $g_0 \in K - R_{\mathrm{rot}}$ and its preimage $\mathrm{pr}^{-1}(g_0) = \{g_0, -g_0\}$, R is equal to $\langle R_{\mathrm{rot}}, g_0 \rangle$ or $\langle R_{\mathrm{rot}}, -g_0 \rangle$ because $K = \langle R_{\mathrm{rot}}, g_0 \rangle$ is the injective image of R. However, if $R = \langle R_{\mathrm{rot}}, g_0 \rangle$, then $R \subset \mathrm{SO}(3)$ and this contradicts the assumption. So, we obtain

$$R = \langle R_{\mathrm{rot}}, -g_0 \rangle. \tag{2.3}$$

In the remaining proof, we apply (2.1), (2.2), (2.3) to each possible K.

If $K = \mathbb{Z}_n$ with odd n and $Z \subset R$, then $R = \mathbb{Z}_n \times Z$. If $K = \mathbb{Z}_n$ with odd n and $Z \not\subset R$, then K has no index two subgroup so that there exists no possible R_{rot} in K. That is, there exists no such R. Therefore, a proof of (1) for odd n is obtained.

If $K = \mathbb{Z}_n$ with even n and $Z \subset R$, then $R = \mathbb{Z}_n \times Z$. If $K = \mathbb{Z}_n$ with even n and $Z \not\subset R$, then there is the unique index two subgroup $\langle a_n^2 \rangle$ of K which should be equal to R_{rot}. Since $a_n \in K - R_{\mathrm{rot}}$, $R = \langle R_{\mathrm{rot}}, -a_n \rangle = \langle -a_n \rangle$ and this is the proof of (1) for even n.

If $K = \mathrm{D}_n$ with odd n and $Z \subset R$, then $R = \mathrm{D}_n \times Z$. If $K = \mathrm{D}_n$ with odd n and $Z \not\subset R$, then there is the unique index two subgroup $\mathbb{Z}_n$ of D_n which should be equal to R_{rot}. Since $b \in K - R_{\mathrm{rot}}$, $R = \langle R_{\mathrm{rot}}, -b\rangle = \langle a_n, -b\rangle$ and this is a proof of (2) for odd n.

If $K = \mathrm{D}_n$ with even n and $Z \subset R$, then $R = \mathrm{D}_n \times Z$. If $K = \mathrm{D}_n$ with even n and $Z \not\subseteq R$, then index two subgroups of D_n are $\mathbb{Z}_n$, $\langle a_n^2, b\rangle$, $\langle a_n^2, a_n b\rangle$ which are candidates for R_{rot} for some R. If $R_{\mathrm{rot}} = \mathbb{Z}_n$, then $R = \langle R_{\mathrm{rot}}, -b\rangle = \langle a_n, -b\rangle$. If $R_{\mathrm{rot}} = \langle a_n^2, b\rangle$, then $R = \langle R_{\mathrm{rot}}, -a_n\rangle = \langle -a_n, b\rangle$. If $R_{\mathrm{rot}} = \langle a_n^2, a_n b\rangle$, then $R = \langle R_{\mathrm{rot}}, -a_n\rangle = \langle -a_n, a_n b\rangle = \langle -a_n, -b\rangle$. Therefore, a proof of (2) for even n is obtained.

If $K = \mathrm{T}$ or I, then it is well-known that K has no index two subgroup because $\mathrm{T} \cong A_4$ and $\mathrm{I} \cong A_5$. So, $Z \not\subseteq R$ can not happen. Therefore, we obtain a proof for (3) and (5).

If $K = \mathrm{O}$ and $Z \subset R$, then $R = \mathrm{O} \times Z$. If $K = \mathrm{O}$ and $Z \not\subseteq R$, then T is the only index two subgroup of O, so $R_{\mathrm{rot}} = \mathrm{T}$. Since $o_0 \in \mathrm{O} - \mathrm{T}$, we have $R = \langle R_{\mathrm{rot}}, -o_0\rangle = \langle \mathrm{T}, -o_0\rangle$, and this is a proof of (4). □ □

In Proposition 2.1, it can be observed that

(1) D_n with $n = 1$ is conjugate to $\mathbb{Z}_n$ with $n = 2$,
(2) $\langle -a_n, b\rangle$ with $n = 2$ is conjugate to $\langle a_n, -b\rangle$ with $n = 2$,
(3) $\langle -a_n\rangle$ with $n = 2$ is conjugate to $\langle a_n, -b\rangle$ with $n = 1$,
(4) $\langle -a_n, -b\rangle$ with $n = 2$ is conjugate to $\langle a_n, -b\rangle$ with $n = 2$,
(5) $\mathbb{Z}_n \times Z$ with $n = 2$ is conjugate to $\mathrm{D}_n \times Z$ with $n = 1$.

To avoid these repetition, we have put conditions '$n > 1$' or '$n > 2$' in five R-entries of Table 1.1.

Second, we deal with closed nonzero-dimensional subgroups of $\mathrm{O}(3)$.

Proposition 2.2. *Let R be a nonzero-dimensional closed subgroup of* $\mathrm{O}(3)$ *such that* $R \not\subseteq \mathrm{SO}(3)$.

(1) *If* $\mathrm{pr}(R)$ *is conjugate to* $\mathrm{SO}(2)$, *then R is conjugate to the group* $\mathrm{SO}(2) \times Z = \langle \mathrm{SO}(2), -a_2\rangle$.
(2) *If* $\mathrm{pr}(R)$ *is conjugate to* $\mathrm{O}(2)$, *then R is conjugate to* $\langle \mathrm{SO}(2), -b\rangle$ *or* $\mathrm{O}(2) \times Z$.
(3) *If* $\mathrm{pr}(R)$ *is conjugate to* $\mathrm{SO}(3)$, *then R is conjugate to* $\mathrm{O}(3)$.

Proof of this is done in a similar way with Proposition 2.1. We have explained for R-entry of Table 1.1. Here, we calculate some isotropy subgroups for later use.

Lemma 2.3. *For each one-dimensional R of Table 1.1 and its natural action on S^2, isotropy subgroups R_{v^0} and $R_{v^0} \cap R_S$ are calculated as in Table 2.1.*

R	R_{v^0}	$R_{v^0} \cap R_S$
$\mathrm{O}(2) \times Z$	$\langle b, -a_2\rangle$	$\langle -a_2 b\rangle$
$\langle \mathrm{SO}(2), -b\rangle$	$\langle -a_2 b\rangle$	$\langle -a_2 b\rangle$
$\langle \mathrm{SO}(2), -a_2\rangle$	$\langle -a_2\rangle$	$\langle \mathrm{id}\rangle$
$\mathrm{O}(2)$	$\langle b\rangle$	$\langle \mathrm{id}\rangle$
$\mathrm{SO}(2)$	$\langle \mathrm{id}\rangle$	$\langle \mathrm{id}\rangle$

Table 2.1. Isotropy subgroups at $v^0 = (1, 0, 0)$ for closed one-dimensional subgroups of $\mathrm{O}(3)$

3. Equivariant simplicial complex structures for finite subgroups of O(3)

Let a finite R of Table 1.1 act naturally on S^2. In this section, we explain for $\mathcal{K}_R$- and D_R-entries of Table 1.1, and investigate equivariance of $|\mathcal{K}_R|$ by calculating isotropy subgroups at some points. We do these for $\mathrm{pr}(R) = \mathbb{Z}_n$, D_n, and then for $\mathrm{pr}(R) = \mathrm{T}$, O, I.

First, we define $\mathcal{K}_m$ and its v^i, e^i for $m = 1$, 2 as promised in Introduction. Denote by $\mathcal{K}_1$ the simplicial complex which is the same with $\mathcal{K}_4$ but has the following notations:

$$\begin{aligned} v^0 &= (1,0,0), \\ b(e^0) &= (-1,0,0), \\ [v^0, b(e^0)] &= [(1,0,0),(0,1,0)] \cup [(0,1,0),(-1,0,0)], \\ |e^0| &= P_4. \end{aligned}$$

And, denote by $\mathcal{K}_2$ the simplicial complex which is the same with $\mathcal{K}_4$ but has the following notations:

$$\begin{aligned} v^i &= \exp\Big(\ \pi i\sqrt{-1}\ \Big), \\ b(e^i) &= \exp\Big(\ \frac{2\pi(2i+1)\sqrt{-1}}{4}\ \Big), \\ |e^i| &= [v^i, b(e^i)] \cup [b(e^i), v^{i+1}] \end{aligned}$$

for $i \in \mathbb{Z}_2$. Also, let $|\mathcal{K}_1|$ and $|\mathcal{K}_2|$ be equal to $|\mathcal{K}_4|$. Here, we remark that if $\mathcal{K}_R = \mathcal{K}_1$ and $D_R = |e^0|$ in Table 1.1, then d^0 and d^1 are defined as v^0. So, we have finished defining $\mathcal{K}_m$ for all natural number m. Similarly, denote by P_m for $m = 1, 2$ the regular polygon P_4. For reader's convenience, we list all possible R's with $\mathcal{K}_R = \mathcal{K}_1$ or $\mathcal{K}_2$.

R	n	$\mathcal{K}_R$
D_n	2	$\mathcal{K}_2$
$\mathbb{Z}_n$	1	$\mathcal{K}_1$
$\mathbb{Z}_n$	2	$\mathcal{K}_2$
$\mathrm{D}_n \times Z$, odd n	1	$\mathcal{K}_2$
$\langle a_n, -b\rangle$, odd n	1	$\mathcal{K}_2$
$\mathbb{Z}_n \times Z$, odd n	1	$\mathcal{K}_2$
$\mathrm{D}_n \times Z$, even n	2	$\mathcal{K}_2$
$\langle a_n, -b\rangle$, even n	2	$\mathcal{K}_2$

TABLE 3.1. Groups with $\mathcal{K}_R = \mathcal{K}_1$ or $\mathcal{K}_2$

Lemma 3.1. *There are eight R's in Table 1.1 which satisfy* $\mathrm{pr}(R) = \mathbb{Z}_n$, D_n *and* $\mathcal{K}_R = \mathcal{K}_1$, $\mathcal{K}_2$. *They are listed in Table 3.1.*

Now, we explain for $\mathcal{K}_R$-entry of Table 1.1 when $\mathrm{pr}(R) = \mathbb{Z}_n$, D_n . Denoting by m_R the number $|R|/|R_{v^0}|$, we can check that $\mathcal{K}_R$ is equal to $\mathcal{K}_{m_R}$. Since $R \cdot v^0$ is equal to the set of vertices of $\mathcal{K}_{m_R}$ in xy-plane, $|\mathcal{K}_R|$ and $\mathcal{K}_R$ are R-invariant for each R. Here, we calculate R_{v^0}.

Lemma 3.2. *For each finite R of Table 1.1 such that* $\mathrm{pr}(R) = \mathbb{Z}_n$, D_n, *isotropy subgroups R_{v^0}, $R_{v^0} \cap R_S$ are calculated as in Table 3.2 when R acts naturally on S^2.*

Proof. By calculation, $R_{v^0} = \langle b\rangle$ for $R = \mathrm{D}_n \times Z$ with odd n, and $R_{v^0} = \langle -a_n^{n/2}, b\rangle$ for $R = \mathrm{D}_n \times Z$ with even n. Other R's are subgroups of one of these two. So, $R_{v^0} = R \cap (\mathrm{D}_n)_{v^0}$ is easily calculated, and from this the group $R_{v^0} \cap R_S$ is also obtained. □ □

We calculate other isotropy subgroups.

R	R_{v^0}	$R_{v^0} \cap R_S$
D_n, $n > 1$	$\langle b \rangle$	$\langle \mathrm{id} \rangle$
$\mathbb{Z}_n$	$\langle \mathrm{id} \rangle$	$\langle \mathrm{id} \rangle$
$\mathrm{D}_n \times Z$, odd n	$\langle b \rangle$	$\langle \mathrm{id} \rangle$
$\langle a_n, -b \rangle$, odd n	$\langle \mathrm{id} \rangle$	$\langle \mathrm{id} \rangle$
$\mathbb{Z}_n \times Z$, odd n	$\langle \mathrm{id} \rangle$	$\langle \mathrm{id} \rangle$
$\mathrm{D}_n \times Z$, even n	$\langle -a_n^{n/2}, b \rangle$	$\langle -a_n^{n/2} b \rangle$
$\langle a_n, -b \rangle$, even n	$\langle -a_n^{n/2} b \rangle$	$\langle -a_n^{n/2} b \rangle$
$\langle -a_n, b \rangle$, odd $n/2$, $n > 2$	$\langle -a_n^{n/2}, b \rangle$	$\langle -a_n^{n/2} b \rangle$
$\langle -a_n, b \rangle$, even $n/2$	$\langle b \rangle$	$\langle \mathrm{id} \rangle$
$\langle -a_n, -b \rangle$, odd $n/2$, $n > 2$	$\langle -a_n^{n/2} \rangle$	$\langle \mathrm{id} \rangle$
$\langle -a_n, -b \rangle$, even $n/2$	$\langle -a_n^{n/2} b \rangle$	$\langle -a_n^{n/2} b \rangle$
$\mathbb{Z}_n \times Z$, even n, $n > 2$	$\langle -a_n^{n/2} \rangle$	$\langle \mathrm{id} \rangle$
$\langle -a_n \rangle$, odd $n/2$, $n > 2$	$\langle -a_n^{n/2} \rangle$	$\langle \mathrm{id} \rangle$
$\langle -a_n \rangle$, even $n/2$	$\langle \mathrm{id} \rangle$	$\langle \mathrm{id} \rangle$

TABLE 3.2. Isotropy subgroup at v^0 for $\mathrm{pr}(R) = \mathbb{Z}_n$ or D_n .

Lemma 3.3. *Let R be a finite group in Table 1.1 such that* $\mathrm{pr}(R) = \mathbb{Z}_n$, D_n *. Isotropy subgroups $R_{b(e^0)}$, $R_{b(e^0)} \cap R_S$, R_x are calculated as in Table 3.3 where x is a point in $P_{m_R} - \{v^i, b(e^i) | i \in \mathbb{Z}_{m_R}\}$.*

R	$\mathcal{K}_R$	$R_{b(e^0)}$	$R_{b(e^0)} \cap R_S$	R_x
D_n, $n > 1$	$\mathcal{K}_n$	$\langle a_n b \rangle$	$\langle \mathrm{id} \rangle$	$\langle \mathrm{id} \rangle$
$\mathbb{Z}_n$	$\mathcal{K}_n$	$\langle \mathrm{id} \rangle$	$\langle \mathrm{id} \rangle$	$\langle \mathrm{id} \rangle$
$\mathrm{D}_n \times Z$, *odd* n	$\mathcal{K}_{2n}$	$\langle -a_n^{(n+1)/2} b \rangle$	$\langle -a_n^{(n+1)/2} b \rangle$	$\langle \mathrm{id} \rangle$
$\langle a_n, -b \rangle$, *odd* n	$\mathcal{K}_{2n}$	$\langle -a_n^{(n+1)/2} b \rangle$	$\langle -a_n^{(n+1)/2} b \rangle$	$\langle \mathrm{id} \rangle$
$\mathbb{Z}_n \times Z$, *odd* n	$\mathcal{K}_{2n}$	$\langle \mathrm{id} \rangle$	$\langle \mathrm{id} \rangle$	$\langle \mathrm{id} \rangle$
$\mathrm{D}_n \times Z$, *even* n	$\mathcal{K}_n$	$\langle -a_n^{n/2}, a_n b \rangle$	$\langle -a_n^{n/2+1} b \rangle$	$\langle -a_n^{n/2} \rangle$
$\langle a_n, -b \rangle$, *even* n	$\mathcal{K}_n$	$\langle -a_n^{n/2+1} b \rangle$	$\langle -a_n^{n/2+1} b \rangle$	$\langle \mathrm{id} \rangle$
$\langle -a_n, b \rangle$, *odd* $n/2$, $n > 2$	$\mathcal{K}_{n/2}$	$\langle -a_n^{n/2}, a_n^2 b \rangle$	$\langle -a_n^{n/2+2} b \rangle$	$\langle -a_n^{n/2} \rangle$
$\langle -a_n, b \rangle$, *even* $n/2$	$\mathcal{K}_n$	$\langle -a_n^{n/2+1} b \rangle$	$\langle -a_n^{n/2+1} b \rangle$	$\langle \mathrm{id} \rangle$
$\langle -a_n, -b \rangle$, *odd* $n/2$, $n > 2$	$\mathcal{K}_n$	$\langle -a_n^{n/2}, a_n b \rangle$	$\langle -a_n^{n/2+1} b \rangle$	$\langle -a_n^{n/2} \rangle$
$\langle -a_n, -b \rangle$, *even* $n/2$	$\mathcal{K}_n$	$\langle a_n b \rangle$	$\langle \mathrm{id} \rangle$	$\langle \mathrm{id} \rangle$
$\mathbb{Z}_n \times Z$, *even* n, $n > 2$	$\mathcal{K}_n$	$\langle -a_n^{n/2} \rangle$	$\langle \mathrm{id} \rangle$	$\langle -a_n^{n/2} \rangle$
$\langle -a_n \rangle$, *odd* $n/2$, $n > 2$	$\mathcal{K}_{n/2}$	$\langle -a_n^{n/2} \rangle$	$\langle \mathrm{id} \rangle$	$\langle -a_n^{n/2} \rangle$
$\langle -a_n \rangle$, *even* $n/2$	$\mathcal{K}_n$	$\langle \mathrm{id} \rangle$	$\langle \mathrm{id} \rangle$	$\langle \mathrm{id} \rangle$

TABLE 3.3. Isotropy subgroups for $\mathrm{pr}(R) = \mathbb{Z}_n$, D_n .

Proof. Since the R-action on $|\mathcal{K}_R|$ is simplicial, any element g in R_x fixes the whole P_{m_R}. Therefore, $R_x = \langle \mathrm{id} \rangle$ or $\langle -a_n^{n/2} \rangle$ for even n, and it is easy to calculate R_x.

Observing that $R_{b(e^0)}$ is isomorphic to a subgroup of $\mathbb{Z}_2 \times \mathbb{Z}_2$ and that $R_x \subset R_{b(e^0)}$ and $R_{b(e^0)} / R_x \cong \langle \mathrm{id} \rangle$ or $\mathbb{Z}_2$, we can calculate $R_{b(e^0)}$ case by case. □ □

Remark 3.4. In the cases of $\langle a_n, -b \rangle$ with odd n and $\langle -a_n, -b \rangle$ with odd $n/2$, it is observed that $|R_{v^0}| > |R_{b(e^0)}|$, so R does not act transitively on $b(e^i)$'s. And, in the case of $\langle -a_n, -b \rangle$ with odd $n/2$, we additionally calculate $R_{b(e^1)} = \langle -a_n^{n/2}, a_n^3 b \rangle$. □

Here, we explain for D_R.

Lemma 3.5. *For each finite R of Table 1.1 such that* $\mathrm{pr}(R) = \mathbb{Z}_n$, D_n, *the R-orbit of the D_R-entry in Table 1.1 covers P_{m_R}, and D_R is a minimal path satisfying such a property. So, any interior point x of D_R is not moved to other point in D_R by R.*

Proof. First, observe that $|R_{v^0}| = |R_{b(e^0)}|$ if and only if R acts transitively on $b(e^i)$'s because R acts transitively on v^i's and $|\mathcal{K}_R|$ has the same number of v^i's and $b(e^i)$'s. And, R acts transitively on $b(e^i)$'s if and only if the R-orbit of $|e^0|$ cover P_{m_R}. By Table 3.2 and 3.3, $|R_{v^0}| = |R_{b(e^0)}|$ except two cases $\langle a_n, -b\rangle$ with odd n and $\langle -a_n, -b\rangle$ with odd $n/2$. If $|R_{v^0}| = |R_{b(e^0)}|$ and $R_{b(e^0)}/R_x \cong \mathbb{Z}_2$, then $[v^0, b(e^0)]$ can be moved to $[b(e^0), v^1]$ by R so that the R-orbit of $[v^0, b(e^0)]$ covers P_{m_R}. In the other side, if $|R_{v^0}| = |R_{b(e^0)}|$ and $R_{b(e^0)} = R_x$, then the R-orbit of $[v^0, b(e^0)]$ does not cover P_{m_R}. If $|R_{v^0}| < |R_{b(e^0)}|$, then the R-orbit of $|e^0|$ does not cover P_{m_R} because R does not act transitively on $b(e^i)$'s, but the R-orbit of $[b(e^0), v^1] \cup [v^1, b(e^1)]$ covers P_{m_R} because R acts transitively on v^i's. So, the remaining of proof is done by comparing Table 3.2 with Table 3.3. □ □

Now, we repeat these arguments for R's satisfying $\mathrm{pr}(R) = \mathrm{T}, \mathrm{O}, \mathrm{I}$.

Lemma 3.6. *For each finite R of Table 1.1 such that* $\mathrm{pr}(R) = \mathrm{T}, \mathrm{O}, \mathrm{I}$, $|\mathcal{K}_R|$ *is R-invariant. And, R act transitively on vertices, edges, faces of $\mathcal{K}_R$, respectively.*

Remark 3.7. For later use, we need understand the case of $R = \mathrm{T} \times Z$ because it is not equal to the full symmetry of $|\mathcal{K}_R| = |\mathcal{K}_\mathrm{O}|$. In this case, $R_{b(f^{-1})}$ is equal to $\mathrm{T}_{b(f^{-1})} \cong \mathbb{Z}_3$. Also, $-a_4^2, -a_4^2 b$ are in $\mathrm{T} \times Z$ because $b, a_4^2 \in \mathrm{T}$. Here, $-a_4^2$, $-a_4^2 b$ are reflections through the xy-plane, xz-plane, respectively. □

Lemma 3.8. *For each finite R of Table 1.1 such that* $\mathrm{pr}(R) = \mathrm{T}$, O, I, *isotropy subgroups R_{v^0}, $R_{b(e^0)}$, R_x are in Table 3.4 where x is an interior point of $[v^0, b(e^0)]$. In the table, the notation $\cong$ is used when an isotropy subgroup is not equal to $\mathbb{Z}_n = \langle a_n\rangle$ or $\mathrm{D}_n = \langle a_n, b\rangle$ for some n but isomorphic to one of them.*

Lemma 3.9. *For each finite R of Table 1.1 such that* $\mathrm{pr}(R) = \mathrm{T}, \mathrm{O}, \mathrm{I}$, *the R-orbit of the D_R-entry in Table 1.1 covers $|\mathcal{K}_R^{(1)}|$, and D_R is a minimal path satisfying such a property.*

We summarize all results of Section 2, 3 in Table 3.4 so that it will be repeatedly referred. In Table 3.4, $R_{d^{-1}}$ is also calculated. In Section 8, 11, we need the following lemma on isotropy subgroups:

Lemma 3.10. *For each finite R of Table 1.1 such that $R \neq \mathbb{Z}_n$, $\langle a_n, -b\rangle$ for any n, isotropy subgroups R_{d^i}'s satisfy*

1. *$R_{d^{-1}} \cap R_{d^i} = R_{[d^{-1}, d^i]}$ and $[d^{-1}, d^i] \subset |\mathcal{K}_R|^{R_{d^{-1}} \cap R_{d^i}}$ for $i \in I$,*
2. *$R_{d^0} \cap R_{d^1} = R_{D_R}$ and $D_R \subset |\mathcal{K}_R|^{R_{d^0} \cap R_{d^1}}$*

where R_X for a subset X of $|\mathcal{K}_R|$ is the subgroup of R fixing X.

Proof. For the natural R-action on S^2 and two points $x \neq \pm x'$ of S^2, we have

$$(*) \qquad R_x \cap R_{x'} = R_C \quad \text{and} \quad C = (S^2)^{R_x \cap R_{x'}}$$

where C is the great circle containing x and x'. From this, we easily obtain proof for cases when $\mathcal{K}_R = \mathcal{K}_\mathrm{T}$, $\mathcal{K}_\mathrm{O}$, $\mathcal{K}_\mathrm{I}$, or $\mathcal{K}_m$ with $m \geq 3$. Then, there are remaining 8 cases by Lemma 3.1. Four cases of these are $\mathbb{Z}_n$ or $\langle a_n, -b\rangle$ for some n so that four cases are remaining. We can apply (*) to three of remaining four. The remaining case is $\mathbb{Z}_n \times Z$ with $n = 1$. Proof for this is easy. □ □

4. EQUIVARIANT CLUTCHING CONSTRUCTION

Let a compact Lie group G_χ act linearly (not necessarily effectively) on S^2 through a representation $\rho : G_\chi \to \mathrm{O}(3)$. Assume that $\rho(G_\chi) = R$ for some finite R in Table 1.1. In the below, our treatment is different according to R. First, we deal with cases when $\mathrm{pr}(R) = \mathrm{T}, \mathrm{O}, \mathrm{I}$ in Section 4∼9. In a similar way, we deal

R	$\mathcal{K}_R$	D_R	R_{v^0}	$R_{b(e^0)}$	R_x	$R_{d^{-1}}$
D_n, $n > 1$	$\mathcal{K}_n$	$[v^0, b(e^0)]$	$\langle b\rangle$	$\langle a_n b\rangle$	$\langle \mathrm{id}\rangle$	$\mathbb{Z}_n$
$\mathbb{Z}_n$	$\mathcal{K}_n$	$\lvert e^0\rvert$	$\langle \mathrm{id}\rangle$	$\langle \mathrm{id}\rangle$	$\langle \mathrm{id}\rangle$	$\mathbb{Z}_n$
$\mathrm{D}_n \times Z$, odd n	$\mathcal{K}_{2n}$	$[v^0, b(e^0)]$	$\langle b\rangle$	$\langle -a_n^{(n+1)/2} b\rangle$	$\langle \mathrm{id}\rangle$	$\langle a_n, -b\rangle$
$\langle a_n, -b\rangle$, odd n	$\mathcal{K}_{2n}$	$[b(e^0), b(e^1)]$	$\langle \mathrm{id}\rangle$	$\langle -a_n^{(n+1)/2} b\rangle$	$\langle \mathrm{id}\rangle$	$\langle a_n, -b\rangle$
$\mathbb{Z}_n \times Z$, odd n	$\mathcal{K}_{2n}$	$\lvert e^0\rvert$	$\langle \mathrm{id}\rangle$	$\langle \mathrm{id}\rangle$	$\langle \mathrm{id}\rangle$	$\mathbb{Z}_n$
$\mathrm{D}_n \times Z$, even n	$\mathcal{K}_n$	$[v^0, b(e^0)]$	$\langle -a_n^{n/2}, b\rangle$	$\langle -a_n^{n/2}, a_n b\rangle$	$\langle -a_n^{n/2}\rangle$	$\langle a_n, -b\rangle$
$\langle a_n, -b\rangle$, even n	$\mathcal{K}_n$	$[v^0, b(e^0)]$	$\langle -a_n^{n/2} b\rangle$	$\langle -a_n^{n/2+1} b\rangle$	$\langle \mathrm{id}\rangle$	$\langle a_n, -b\rangle$
$\langle -a_n, b\rangle$, odd $n/2$, $n > 2$	$\mathcal{K}_{n/2}$	$\lvert e^0\rvert$	$\langle -a_n^{n/2}, b\rangle$	$\langle -a_n^{n/2}, a_n^2 b\rangle$	$\langle -a_n^{n/2}\rangle$	$\langle a_n^2, -a_n b\rangle$
$\langle -a_n, b\rangle$, even $n/2$	$\mathcal{K}_n$	$[v^0, b(e^0)]$	$\langle b\rangle$	$\langle -a_n^{n/2+1} b\rangle$	$\langle \mathrm{id}\rangle$	$\langle a_n^2, -a_n b\rangle$
$\langle -a_n, -b\rangle$, odd $n/2$, $n > 2$	$\mathcal{K}_n$	$[b(e^0), b(e^1)]$	$\langle -a_n^{n/2}\rangle$	$\langle -a_n^{n/2}, a_n b\rangle$	$\langle -a_n^{n/2}\rangle$	$\langle a_n^2, -b\rangle$
$\langle -a_n, -b\rangle$, even $n/2$	$\mathcal{K}_n$	$[v^0, b(e^0)]$	$\langle -a_n^{n/2} b\rangle$	$\langle a_n b\rangle$	$\langle \mathrm{id}\rangle$	$\langle a_n^2, -b\rangle$
$\mathbb{Z}_n \times Z$, even n, $n > 2$	$\mathcal{K}_n$	$\lvert e^0\rvert$	$\langle -a_n^{n/2}\rangle$	$\langle -a_n^{n/2}\rangle$	$\langle -a_n^{n/2}\rangle$	$\mathbb{Z}_n$
$\langle -a_n\rangle$, odd $n/2$, $n > 2$	$\mathcal{K}_{n/2}$	$\lvert e^0\rvert$	$\langle -a_n^{n/2}\rangle$	$\langle -a_n^{n/2}\rangle$	$\langle -a_n^{n/2}\rangle$	$\langle a_n^2\rangle$
$\langle -a_n\rangle$, even $n/2$	$\mathcal{K}_n$	$\lvert e^0\rvert$	$\langle \mathrm{id}\rangle$	$\langle \mathrm{id}\rangle$	$\langle \mathrm{id}\rangle$	$\langle a_n^2\rangle$
T	$\mathcal{K}_{\mathrm{T}}$	$[v^0, b(e^0)]$	$\cong \mathbb{Z}_3$	$\cong \mathbb{Z}_2$	$\langle \mathrm{id}\rangle$	$\cong \mathbb{Z}_3$
O	$\mathcal{K}_{\mathrm{O}}$	$[v^0, b(e^0)]$	$\cong \mathbb{Z}_4$	$\cong \mathbb{Z}_2$	$\langle \mathrm{id}\rangle$	$\cong \mathbb{Z}_3$
I	$\mathcal{K}_{\mathrm{I}}$	$[v^0, b(e^0)]$	$\cong \mathbb{Z}_5$	$\cong \mathbb{Z}_2$	$\langle \mathrm{id}\rangle$	$\cong \mathbb{Z}_3$
$\langle \mathrm{T}, -o_0\rangle$	$\mathcal{K}_{\mathrm{T}}$	$[v^0, b(e^0)]$	$\cong \mathrm{D}_3$	$\cong \mathbb{Z}_2 \times \mathbb{Z}_2$	$\cong \mathbb{Z}_2$	$\cong \mathrm{D}_3$
$\mathrm{T} \times Z$	$\mathcal{K}_{\mathrm{O}}$	$\lvert e^0\rvert$	$\langle -a_4^2, b\rangle$	$\langle -a_4^2\rangle$	$\langle -a_4^2\rangle$	$\cong \mathbb{Z}_3$
$\mathrm{O} \times Z$	$\mathcal{K}_{\mathrm{O}}$	$[v^0, b(e^0)]$	$\cong \mathrm{D}_4$	$\cong \mathbb{Z}_2 \times \mathbb{Z}_2$	$\cong \mathbb{Z}_2$	$\cong \mathrm{D}_3$
$\mathrm{I} \times Z$	$\mathcal{K}_{\mathrm{I}}$	$[v^0, b(e^0)]$	$\cong \mathrm{D}_5$	$\cong \mathbb{Z}_2 \times \mathbb{Z}_2$	$\cong \mathbb{Z}_2$	$\cong \mathrm{D}_3$
$\mathrm{O}(3)$						
$\mathrm{O}(2) \times Z$		$\{v^0\}$	$\langle b, -a_2\rangle$			$\langle \mathrm{SO}(2), -b\rangle$
$\langle \mathrm{SO}(2), -b\rangle$		$\{v^0\}$	$\langle -a_2 b\rangle$			$\langle \mathrm{SO}(2), -b\rangle$
$\langle \mathrm{SO}(2), -a_2\rangle$		$\{v^0\}$	$\langle -a_2\rangle$			$\mathrm{SO}(2)$
$\mathrm{SO}(3)$						
$\mathrm{O}(2)$		$\{v^0\}$	$\langle b\rangle$			$\mathrm{SO}(2)$
$\mathrm{SO}(2)$		$\{v^0\}$	$\langle \mathrm{id}\rangle$			$\mathrm{SO}(2)$

TABLE 3.4. $\mathcal{K}_R, D_R$, and isotropy subgroups

with cases when $\mathrm{pr}(R) = \mathbb{Z}_n, \mathrm{D}_n$ in Section 10∼11. Then, cases of one-dimensional R's are dealt with.

Assume that $\mathrm{pr}(R) = \mathrm{T}, \mathrm{O}, \mathrm{I}$. Let $\bar{\mathcal{K}}_R$ be the simplicial complex $\amalg_{f \in \mathcal{K}_R} f$, i.e. the disjoint union of faces of $\mathcal{K}_R$. In this section, we would consider $|\mathcal{K}_R|$ as the quotient of the underlying space $|\bar{\mathcal{K}}_R| = \amalg_{f \in \mathcal{K}_R} |f|$. And, we would consider an equivariant vector bundle over $|\mathcal{K}_R|$ as an equivariant clutching construction of an equivariant vector bundle over $|\bar{\mathcal{K}}_R|$. For this, we would define equivariant clutching map and its generalization preclutching map. And, we state equivalent conditions under which a preclutching map is an equivariant clutching map. Before these, we need introduce notations on some relevant simplicial complices. Since we should deal with various cases at the same time, these notations are necessary. Examples of the notations are illustrated in Figure 4.1 and Figure 4.2.

First, we define some notations on $\mathcal{K}_R$ and $\bar{\mathcal{K}}_R$. We denote simply by π, $|\pi|$ natural quotient maps from $\bar{\mathcal{K}}_R$, $|\bar{\mathcal{K}}_R|$ to $\mathcal{K}_R, |\mathcal{K}_R|$, respectively. By definition, $|\pi| \,\Big|_{|f|}$ is bijective for each face $f \in \bar{\mathcal{K}}_R$. From this, the G_χ-actions on $\mathcal{K}_R$, $|\mathcal{K}_R|$ induce G_χ-actions on $\bar{\mathcal{K}}_R$, $|\bar{\mathcal{K}}_R|$ so that π, $|\pi|$ are equivariant, respectively. We use notations $\bar{v}$, $\bar{e}$, $\bar{f}$ to denote a vertex, an edge, a face of $\bar{\mathcal{K}}_R$, respectively. We use the notation $\bar{x}$ to denote an arbitrary point of $|\bar{\mathcal{K}}_R^{(1)}|$. When $\bar{v}$, $\bar{e}$, $\bar{f}$, $\bar{x}$ are understood, we use notations v, e, f, x to denote images $\pi(\bar{v})$, $\pi(\bar{e})$, $\pi(\bar{f})$, $|\pi|(\bar{x})$, respectively. Denote by $\bar{f}^{-1}$, $\bar{f}^i$ faces of $\bar{\mathcal{K}}_R$ such that $\pi(\bar{f}^{-1}) = f^{-1}$ and $\pi(\bar{f}^i) = f^i$, and denote by $\bar{v}^i$, $\bar{e}^i$ simplices of $\bar{f}^{-1}$ such that $\pi(\bar{v}^i) = v^i$ and $\pi(\bar{e}^i) = e^i$. And, denote by $\bar{d}^0$, $\bar{d}^1$ points of $|\bar{f}^{-1}|$ such that $|\pi|(\bar{d}^0) = d^0$ and $|\pi|(\bar{d}^1) = d^1$, and denote by $\bar{D}_R$ the



(a) $\mathcal{K}_{\mathrm{T}}$ near v^0 seen from above (b) $\bar{\mathcal{K}}_{\mathrm{T}}$ near $\{\bar{v}^0_j \mid j \in \mathbb{Z}_3\}$



(c) $\hat{\mathcal{L}}_{\mathrm{T}}$ near $\{\hat{v}^0_{j,\pm} \mid j \in \mathbb{Z}_3\}$

FIGURE 4.1. Relation between $\mathcal{K}_R$, $\bar{\mathcal{K}}_R$, $\hat{\mathcal{L}}_R$ in the case when $\mathcal{K}_R = \mathcal{K}_{\mathrm{T}}$ and $j_R = 3$

path $|\pi|^{-1}(D_R) \cap |\bar{f}^{-1}|$, i.e. $\bar{D}_R = [\bar{d}^0, \bar{d}^1]$. Define the integer j_R as the cardinality of $\pi^{-1}(v^i)$ for $i \in \mathbb{Z}_3$, i.e. $j_R = 3$, 4, 5 according to $\mathcal{K}_R = \mathcal{K}_{\mathrm{T}}$, $\mathcal{K}_{\mathrm{O}}$, $\mathcal{K}_{\mathrm{I}}$, respectively. Let B be the subset $\{\ b(\bar{f}) \mid \bar{f} \in \bar{\mathcal{K}}_R\ \}$ of $|\bar{\mathcal{K}}_R|$ on which R (and G_χ) acts transitively by Lemma 3.6, and B is often confused with $|\pi|(B) = \{\ b(f) \mid f \in \mathcal{K}_R\ \}$. So far, we have defined superscript i for simplices in $\bar{\mathcal{K}}_R$. Next, we define $\bar{x}_j$ with subscript j for any point $\bar{x}$ of $|\bar{\mathcal{K}}^{(1)}_R|$.

Notation 4.1.

(1) For a vertex $\bar{v}$ in $\bar{\mathcal{K}}^{(1)}_R$ and $v = \pi(\bar{v})$, we label vertices in $\pi^{-1}(v)$ with $\bar{v}_j$ to satisfy
  i) $\pi^{-1}(v) = \{\bar{v}_j \mid j \in \mathbb{Z}_{j_R}\}$,
  ii) $\bar{v}_0 = \bar{v}$,
  iii) in each face $|\bar{f}_j|$ containing $\bar{v}_j$ for $j \in \mathbb{Z}_{j_R}$, we can take a small neighborhood U_j of $\bar{v}_j$ so that $|\pi|(U_j)$'s are arranged in the counterclockwise way around v.

(2) For a non-vertex $\bar{x}$ in $|\bar{\mathcal{K}}^{(1)}_R|$ and $x = |\pi|(\bar{x})$, we label two points in $|\pi|^{-1}(x)$ with $\{\bar{x}_j \mid j \in \mathbb{Z}_2\}$ to satisfy $\bar{x}_0 = \bar{x}$.

For simplicity, we denote $(\bar v^i)_j$, $(\bar d^i)_j$ by $\bar v^i_j$, $\bar d^i_j$, respectively.

We need introduce two more simplicial complices. Denote by $\bar{\mathcal{L}}_R$ and $\hat{\mathcal{L}}_R$ the 1-skeleton $\bar{\mathcal{K}}_R^{(1)}$ of $\bar{\mathcal{K}}_R$ and the disjoint union $\amalg_{\bar e \in \bar{\mathcal{L}}_R} \bar e$, respectively. Then, $\bar{\mathcal{L}}_R$ is a subcomplex of $\bar{\mathcal{K}}_R$, and can be regarded as a quotient of $\hat{\mathcal{L}}_R$. These relations are expressed by two natural simplicial maps

$$\imath_{\bar{\mathcal{L}}} : \bar{\mathcal{L}}_R \to \bar{\mathcal{K}}_R, \quad p_{\bar{\mathcal{L}}} : \hat{\mathcal{L}}_R \to \bar{\mathcal{L}}_R$$

where $\imath_{\bar{\mathcal{L}}}$ is the inclusion and $p_{\bar{\mathcal{L}}}$ is the quotient map whose preimage of each vertex and edge of $\bar{\mathcal{L}}_R$ consists of two vertices and one edge of $\hat{\mathcal{L}}_R$, respectively. Two maps on underlying spaces are denoted by

$$\imath_{|\bar{\mathcal{L}}|} : |\bar{\mathcal{L}}_R| \to |\bar{\mathcal{K}}_R|, \quad p_{|\bar{\mathcal{L}}|} : |\hat{\mathcal{L}}_R| \to |\bar{\mathcal{L}}_R|.$$

The G_χ-actions on $\bar{\mathcal{K}}_R$, $|\bar{\mathcal{K}}_R|$ naturally induce G_χ-actions on these relevant simplicial complices $\bar{\mathcal{L}}_R, \hat{\mathcal{L}}_R$ and their underlying spaces. We need introduce notations on simplices of $\hat{\mathcal{L}}_R$ and points of $|\hat{\mathcal{L}}_R|$. We use notations $\hat v$ and $\hat e$ to denote a vertex and an edge of $\hat{\mathcal{L}}_R$, respectively. And, we use the notation $\hat x$ to denote an arbitrary point in $|\hat{\mathcal{L}}_R|$. When $\hat v$, $\hat e$, $\hat x$ are understood, we use notations $\bar v$, $\bar e$, $\bar x$ to denote $p_{\bar{\mathcal{L}}}(\hat v)$, $p_{\bar{\mathcal{L}}}(\hat e)$, $p_{|\bar{\mathcal{L}}|}(\hat x)$, respectively. Two edges $\hat e, \hat e'$ of $\hat{\mathcal{L}}_R$ (and their images $\bar e, \bar e'$ in $\bar{\mathcal{L}}_R$) are called *adjacent* if $\hat e \neq \hat e'$ and $\pi(p_{\bar{\mathcal{L}}}(\hat e)) = \pi(p_{\bar{\mathcal{L}}}(\hat e'))$. And, two faces $\bar f, \bar f'$ of $\bar{\mathcal{K}}_R$ are called *adjacent* if their images f, f' are adjacent.

Next, we introduce superscript i and subscripts $+, -$ for vertices and edges of $\hat{\mathcal{L}}_R$. Before it, we introduce a simplicial map. Let $c : \hat{\mathcal{L}}_R \to \hat{\mathcal{L}}_R$ be the simplicial map whose underlying space map $|c| : |\hat{\mathcal{L}}_R| \to |\hat{\mathcal{L}}_R|$ is defined as

> for any adjacent $\hat e, \hat e' \in \hat{\mathcal{L}}_R$, each point $\hat x$ in $|\hat e|$ is sent to the point $|c|(\hat x)$ in $|\hat e'|$ to satisfy $|\pi|(p_{|\bar{\mathcal{L}}|}(\hat x)) = |\pi|(p_{|\bar{\mathcal{L}}|}(c(\hat x)))$.

For example, $\hat e$ and $c(\hat e)$ are adjacent for any edge $\hat e$ in $\hat{\mathcal{L}}_R$. Easily, c and $|c|$ are G_χ-equivariant. For notational simplicity, we define c also on edges of $\bar{\mathcal{L}}_R$ to satisfy $c(p_{\bar{\mathcal{L}}}(\hat e)) = p_{\bar{\mathcal{L}}}(c(\hat e))$ for each edge $\hat e$.

Notation 4.2.

1. For a vertex $\bar v$ in $\bar{\mathcal{L}}_R$, we label two vertices in $p_{\bar{\mathcal{L}}}^{-1}(\bar v)$ with $\hat v_\pm$ to satisfy
$$p_{\bar{\mathcal{L}}}(c(\hat v_+)) = \bar v_1 \ \text{ and } \ p_{\bar{\mathcal{L}}}(c(\hat v_-)) = \bar v_{-1} = \bar v_{j_R - 1}.$$
2. For a non-vertex $\bar x$ in $|\bar{\mathcal{L}}_R|$, we label the point in $p_{|\bar{\mathcal{L}}|}^{-1}(\bar x)$ with $\hat x_+$ or $\hat x_-$, i.e. $\hat x_+ = \hat x_-$.

For simplicity, denote $\hat x_\pm$ for $\bar x = \bar v^i, \bar v^i_j, \bar d^i, \bar d^i_j$ by $\hat v^i_\pm, \hat v^i_{j,\pm}, \hat d^i_\pm, \hat d^i_{j,\pm}$, respectively. So, if $\bar d^i$ is a barycenter of an edge, then $\hat d^i_+ = \hat d^i_-$. And, denote by $\hat e^i$ the edge in $\hat{\mathcal{L}}_R$ such that $p_{\bar{\mathcal{L}}}(\hat e^i) = \bar e^i$ for $i \in \mathbb{Z}_3$.

Until now, we have finished introducing notations in Figure 4.1. By using these notations, we introduce one-dimensional fundamental domain in $|\hat{\mathcal{L}}_R|$. For $\bar D_R = [\bar d^0, \bar d^1] \subset |\bar{\mathcal{K}}_R|$, we define $\hat D_R$ in $|\hat{\mathcal{L}}_R|$ as $[\hat d^0_+, \hat d^1_-]$ so that $p_{|\bar{\mathcal{L}}|}(\hat D_R) = \bar D_R$. And, denote by $\hat{\mathbf{D}}_R$ the set $(|\pi| \circ p_{|\bar{\mathcal{L}}|})^{-1}(D_R)$ in $|\hat{\mathcal{L}}_R|$ which is equal to

$$[\hat d^0_+, \hat d^1_-] \bigcup |c|\big([\hat d^0_+, \hat d^1_-]\big) \bigcup \Big(\bigcup_{v \in D_R} (\pi \circ p_{\bar{\mathcal{L}}})^{-1}(v) \Big).$$

The union of thick points and edges of Figure 4.2.(b) is $\hat{\mathbf{D}}_R$ in the case of $R = \mathrm{T} \times Z$.

For convenience in calculation, we parameterize each edge of $|\hat{\mathcal{L}}_R|$ linearly by $s \in [0, 1]$ to satisfy

1. $\hat v_+ = 0$, $\hat v_- = 1$, $b(\hat e) = 1/2$ for each vertex $\bar v$ of $\bar{\mathcal{K}}_R$ and each edge $\hat e$ of $\hat{\mathcal{L}}_R$,
2. $|c|(s) = 1 - s$ for each edge $\hat e$ of $\hat{\mathcal{L}}_R$ and $s \in |\hat e|$.



(a) $\mathcal{K}_{\mathrm{O}}$



(b) $\hat{\mathcal{L}}_R$ near $\hat{\mathbf{D}}_R$

FIGURE 4.2. $\hat{\mathcal{L}}_R$ and $\hat{\mathbf{D}}_R$ in the case when $R = \mathrm{T} \times Z$, $\mathcal{K}_R = \mathcal{K}_{\mathrm{O}}$, $D_R = |e^0|$

We repeatedly use this parametrization.

Now, we describe an equivariant vector bundle over $|\mathcal{K}_R|$ as an equivariant clutching construction of an equivariant vector bundle over $|\bar{\mathcal{K}}_R|$. Let V_B be a G_χ-vector bundle over B such that $(\mathrm{res}_H^{G_\chi} V_B)|_{b(\bar{f})}$ is χ-isotypical at each $b(\bar{f})$ in B. If we denote by $V_{\bar{f}}$ the isotropy representation of V_B at each $b(\bar{f})$, then $V_B \cong G_\chi \times_{(G_\chi)_{b(\bar{f})}} V_{\bar{f}}$ because G_χ acts transitively on B. And, $\mathrm{res}_H^{(G_\chi)_{b(\bar{f})}} V_{\bar{f}}$'s are all isomorphic because they are all χ-isotypical. We define $\mathrm{Vect}_{G_\chi}(|\mathcal{K}_R|, \chi)_{V_B}$ as the set

$$\left\{ \quad [E] \in \mathrm{Vect}_{G_\chi}(|\mathcal{K}_R|, \chi) \quad \middle| \quad E|_B \cong V_B \quad \right\}.$$

Similarly, $\mathrm{Vect}_{G_\chi}(|\bar{\mathcal{K}}_R|, \chi)_{V_B}$ is defined. Observe that $\mathrm{Vect}_{G_\chi}(|\bar{\mathcal{K}}_R|, \chi)_{V_B}$ has the unique element $[F_{V_B}]$ for the bundle $F_{V_B} = G_\chi \times_{(G_\chi)_{b(\bar{f}^{-1})}} (|\bar{f}^{-1}| \times V_{\bar{f}^{-1}})$ because

$|\bar{\mathcal{K}}_R| \cong G_\chi \times_{(G_\chi)_{b(\bar{f}^{-1})}} |\bar{f}^{-1}|$ is equivariant homotopically equivalent to B. Henceforward, we use trivializations

$$\begin{array}{lll} (4.1) & |\bar{f}| \times V_{\bar{f}} & \text{for } \big(\operatorname{res}^{G_\chi}_{(G_\chi)_{b(\bar{f})}} F_{V_B}\big)\big|_{|\bar{f}|}, \\ & |\bar{e}| \times \operatorname{res}^{(G_\chi)_{b(\bar{f})}}_{(G_\chi)_{b(\bar{e})}} V_{\bar{f}} & \text{for } \big(\operatorname{res}^{G_\chi}_{(G_\chi)_{b(\bar{e})}} F_{V_B}\big)\big|_{|\bar{e}|} \end{array}$$

for each face $\bar{f}$ and edge $\bar{e} \in \bar{f}$. Observe that each $E \in \mathrm{Vect}_{G_\chi}(|\mathcal{K}_R|, \chi)_{V_B}$ can be constructed by gluing the pull-back bundle $|\pi|^*E$ along edges. Let us describe this more precisely. Let $\widetilde{|\pi|}$ be the bundle morphism covering $|\pi|$

$$\begin{tikzcd} {|\pi|^*E} \arrow[r, "\widetilde{|\pi|}"] \arrow[d] & E \arrow[d] \\ {|\bar{\mathcal{K}}_R|} \arrow[r, "|\pi|"] & {|\mathcal{K}_R|}. \end{tikzcd}$$

Consider the following equivalence relation $\sim$ on vectors of $|\pi|^*E$:

for any two u, u' in $|\pi|^*E$, $u \sim u'$ if and only if $\widetilde{|\pi|}(u) = \widetilde{|\pi|}(u')$.

Since $\widetilde{|\pi|}$ is equivariant, the quotient $|\pi|^*E/\sim$ of $|\pi|^*E$ through the relation delivers the equivariant vector bundle structure inherited from $|\pi|^*E$. Here, note that it suffices to define the relation only on vectors in $|\pi|^*E\big|_{|\bar{\mathcal{L}}_R|}$ to define the quotient which is trivially isomorphic to E. We call the construction of the bundle $|\pi|^*E/\sim$ *equivariant clutching construction.* Since $F_{V_B} \cong |\pi|^*E$ for any E, we may rewrite the construction by using F_{V_B} instead of $|\pi|^*E$. Pick an equivariant isomorphism $\bar{A} : F_{V_B} \to |\pi|^*E$. Then, the relation $\sim$ induces the following equivalence relation $\sim'$ on vectors of F_{V_B} :

for any two u, u' in F_{V_B}, $u \sim' u'$ if and only if $\bar{A}(u) \sim \bar{A}(u')$.

Then, the quotient $F_{V_B}/\sim'$ delivers the equivariant vector bundle structure inherited from F_{V_B}, and trivially $(F_{V_B}/\sim') \cong E$. Let us describe the relation on vectors in $F_{V_B}\big|_{|\bar{\mathcal{L}}_R|}$ more precisely. By using trivialization (4.1) of F_{V_B}, the quotient $F_{V_B}/\sim'$ can be also constructed by gluing F_{V_B} along edges through

$$(4.2) \qquad |\bar{e}| \times V_{\bar{f}} \longrightarrow |c(\bar{e})| \times V_{\bar{f}'}, \quad (\ p_{|\bar{\mathcal{L}}|}(\hat{x}), u\ ) \mapsto \Big(\ p_{|\bar{\mathcal{L}}|}\big(|c|(\hat{x})\big),\ \varphi_{\hat{e}}(\hat{x})\ u\ \Big)$$

via some continuous maps

$$\varphi_{\hat{e}} : |\hat{e}| \to \mathrm{Iso}(V_{\bar{f}}, V_{\bar{f}'})$$

for each edge $\hat{e}$, $\hat{x} \in |\hat{e}|$, $u \in V_{\bar{f}}$ where $\bar{e} = p_{\bar{\mathcal{L}}}(\hat{e})$ and $\bar{e} \in \bar{f}$, $c(\bar{e}) \in \bar{f}'$. Here, the notation Iso with no subscript means the set of nonequivariant isomorphisms. The union $\Phi = \bigcup_{\hat{e} \in \hat{\mathcal{L}}_R} \varphi_{\hat{e}}$ is called an *equivariant clutching map* of E with respect to V_B. The relation $\sim'$ on vectors in $F_{V_B}|_{|\bar{\mathcal{L}}_R|}$ is defined by Φ, and the quotient $F_{V_B}/\sim'$ is denoted by F_{V_B}/Φ. And, the equivariant vector bundle F_{V_B}/Φ is called *determined* by Φ with respect to V_B. When we use the phrase 'with respect to V_B', it is assumed that we use the bundle F_{V_B} and its trivialization (4.1) in gluing. Equivariance of $\widetilde{|\pi|}$ and $\bar{A}$ guarantees equivariance of Φ, i.e.

$$(g \cdot \Phi)(\hat{x}) = g\Phi(g^{-1}\hat{x})g^{-1} = \Phi(\hat{x})$$

for all $g \in G_\chi, \hat{x} \in |\hat{\mathcal{L}}_R|$. We denote by p_Φ the quotient map from F_{V_B} to F_{V_B}/Φ. Here, note that Φ is defined on $|\hat{\mathcal{L}}_R|$. That is why we define $\hat{\mathcal{L}}_R$. Sometimes, we regard Φ as the map

$$p^*_{|\bar{\mathcal{L}}|}F_{V_B} \to p^*_{|\bar{\mathcal{L}}|}F_{V_B}, \quad (\hat{x}, u) \mapsto \big(|c|(\hat{x}), \Phi(\hat{x})u\big)$$

by using trivialization (4.1) for each $(\hat{x}, u) \in |\hat{e}| \times V_{\bar{f}}$ where $\bar{e} \in \bar{f}$. An equivariant clutching map of some bundle in $\mathrm{Vect}_{G_\chi}(|\mathcal{K}_R|, \chi)_{V_B}$ with respect to V_B is called simply an *equivariant clutching map* with respect to V_B, and let Ω_{V_B} be the set of all equivariant clutching maps with respect to V_B. In the next section, we will see that we need calculate the (nonequivariant) homotopy $\pi_0(\Omega_{V_B})$ to classify equivariant vector bundles. To do it, we need to restrict an equivariant clutching map in Ω_{V_B} to $\hat{D}_R$. We explain for this. Let $\Omega_{\hat{D}_R, V_B}$ be the set

$$\{\Phi|_{\hat{D}_R} \mid \Phi \in \Omega_{V_B}\}.$$

If two equivariant clutching maps coincide on $\hat{D}_R$, then they are identical by equivariance and definition of one-dimensional fundamental domain. So, the restriction map $\Omega_{V_B} \to \Omega_{\hat{D}_R, V_B}$, $\Phi \mapsto \Phi|_{\hat{D}_R}$ is bijective, and we obtain a bijection $\pi_0(\Omega_{V_B}) \cong \pi_0(\Omega_{\hat{D}_R, V_B})$ between two homotopies. It is conceivable that it is easier to deal with $\Omega_{\hat{D}_R, V_B}$ than Ω_{V_B} because of smaller domain of definition. This is why we restrict an equivariant clutching map to $\hat{D}_R$. We call a map Φ in Ω_{V_B} the *extension* of $\Phi|_{\hat{D}_R}$ in $\Omega_{\hat{D}_R, V_B}$. And, denote the bundle F_{V_B}/Φ also by $F_{V_B}/\ \Phi|_{\hat{D}_R}$.

Next, we define preclutching map, a generalization of equivariant clutching map. Let $C^0(|\hat{\mathcal{L}}_R|, V_B)$ be the set of continuous functions Φ on $|\hat{\mathcal{L}}_R|$ satisfying $\Phi|_{|\hat{e}|}(\hat{x}) \in \mathrm{Iso}(V_{\bar{f}}, V_{\bar{f}'})$ for each $\hat{e}$ and $\hat{x} \in |\hat{e}|$ where $\bar{e} \in \bar{f}$, $c(\bar{e}) \in \bar{f}'$. Note that we can define the quotient F_{V_B}/Φ also for any $\Phi \in C^0(|\hat{\mathcal{L}}_R|, V_B)$ as we have done in (4.2) though F_{V_B}/Φ need not deliver a suitable equivariant vector bundle structure or even nonequivariant vector bundle structure. Let $C^0(\hat{D}_R, V_B)$ be the set

$$\left\{ \ \Phi|_{\hat{D}_R} \ \middle| \ \Phi \in C^0\big(|\hat{\mathcal{L}}_R|, V_B\big) \ \right\}.$$

A function Φ in $C^0(|\hat{\mathcal{L}}_R|, V_B)$ or a function $\Phi_{\hat{D}_R}$ in $C^0(\hat{D}_R, V_B)$ is called a *preclutching map* with respect to V_B. Then, it is a natural question under which conditions a preclutching map becomes an equivariant clutching map. We can answer this question for a preclutching map in $C^0(|\hat{\mathcal{L}}_R|, V_B)$. A preclutching map Φ in $C^0(|\hat{\mathcal{L}}_R|, V_B)$ is an equivariant clutching map with respect to V_B if and only if it satisfies the following conditions:

N1. $\Phi(|c|(\hat{x})) = \Phi(\hat{x})^{-1}$ for each $\hat{x} \in |\hat{\mathcal{L}}_R|$,

N2. For each vertex $\bar{v} \in \bar{\mathcal{K}}_R$,

$$\Phi(\hat{v}_{j_R-1,+}) \cdots \Phi(\hat{v}_{j,+}) \cdots \Phi(\hat{v}_{0,+}) = \mathrm{id}$$

for $j \in \mathbb{Z}_{j_R}$,

E1. $\Phi(g\hat{x}) = g\Phi(\hat{x})g^{-1}$ for each $\hat{x} \in |\hat{\mathcal{L}}_R|, g \in G_\chi$.

We explain for this more precisely. As a slight generalization of the classical result [At, p. 20, 21], if Φ satisfies Condition N1., N2., then the quotient F_{V_B}/Φ becomes a nonequivariant vector bundle though it need not be an equivariant vector bundle. Moreover, if Φ also satisfies Condition E1., then F_{V_B}/Φ becomes an equivariant vector bundle so that Φ is an equivariant clutching map with respect to V_B. We will answer the same question for a preclutching map in $C^0(\hat{D}_R, V_B)$ in Section 8.

5. Relations among $\mathrm{Vect}_{G_\chi}(S^2, \chi)_{V_B}$, $A_{G_\chi}(S^2, \chi)$, $\pi_0(\Omega_{V_B})$

In this section, we investigate relations among

$$\mathrm{Vect}_{G_\chi}(S^2, \chi)_{V_B}, \quad A_{G_\chi}(S^2, \chi), \quad \pi_0(\Omega_{V_B}).$$

Our classification of the paper is based on these relations. Before it, we state two basic facts on equivariant vector bundles. First, two equivariantly homotopic equivariant clutching maps give isomorphic equivariant vector bundles.

Lemma 5.1. *For two maps* Φ *and* Φ' *in* Ω_{V_B}*, if* Φ *and* Φ' *are homotopic in* Ω_{V_B}*, i.e.* $[\Phi] = [\Phi']$ *in* $\pi_0(\Omega_{V_B})$*, then* $\big[F_{V_B}/\Phi\big] = \big[F_{V_B}/\Phi'\big]$ *in* $\mathrm{Vect}_{G_\chi}(S^2,\chi)_{V_B}$*.*

Proof. This is a slight generalization of the classical result [At, Lemma 1.4.6. and Section 1.6.]. So, we omit the proof. □ □

Lemma 5.1 gives a sufficient condition for isomorphism. Sometimes, we need an equivalent condition. When we consider a map in Ω_{V_B} as defined on $p^*_{|\bar{\mathcal{L}}|}F_{V_B}$, we have the following equivalent condition:

Lemma 5.2. *For any* Φ *and* Φ' *in* Ω_{V_B}*,* $\big[F_{V_B}/\Phi\big] = \big[F_{V_B}/\Phi'\big]$ *in* $\mathrm{Vect}_{G_\chi}(S^2,\chi)_{V_B}$ *if and only if there is a* G_χ*-isomorphism* $\Theta : F_{V_B} \to F_{V_B}$ *such that* $(p^*_{|\bar{\mathcal{L}}|}\Theta)\Phi = \Phi'(p^*_{|\bar{\mathcal{L}}|}\Theta)$ *where* $p^*_{|\bar{\mathcal{L}}|}\Theta : p^*_{|\bar{\mathcal{L}}|}F_{V_B} \to p^*_{|\bar{\mathcal{L}}|}F_{V_B}$ *is the pull-back of* Θ*.*

$$\begin{tikzcd}
p^*_{|\bar{\mathcal{L}}|}F_{V_B} \arrow[r, "p^*_{|\bar{\mathcal{L}}|}\Theta"] \arrow[d, "\Phi"] & p^*_{|\bar{\mathcal{L}}|}F_{V_B} \arrow[d, "\Phi'"] \\
p^*_{|\bar{\mathcal{L}}|}F_{V_B} \arrow[r, "p^*_{|\bar{\mathcal{L}}|}\Theta"] & p^*_{|\bar{\mathcal{L}}|}F_{V_B}
\end{tikzcd}$$

Proof. This is also a slight generalization of the classical result [At, p. 22, (ii) and Section 1.6.]. First, we prove sufficiency. Let $A : F_{V_B}/\Phi \to F_{V_B}/\Phi'$ be a G_χ-isomorphism. Then, we can show that there exists the G_χ-isomorphism $\bar{A} : F_{V_B} \to F_{V_B}$ satisfying the following commutative diagram

$$\begin{tikzcd}
F_{V_B} \arrow[r, "\bar{A}"] \arrow[d, "p_\Phi"] & F_{V_B} \arrow[d, "p_{\Phi'}"] \\
F_{V_B}/\Phi \arrow[r, "A"] & F_{V_B}/\Phi'.
\end{tikzcd} \tag{5.1}$$

From this, $(p^*_{|\mathcal{L}|}\bar{A})\Phi = \Phi'(p^*_{|\mathcal{L}|}\bar{A})$ is obtained.

Next, we prove necessity. If we put $\bar{A} = \Theta$, there exists the unique G_χ-isomorphism A satisfying (5.1) by the assumption. So, we obtain a proof. □ □

Consider the map $\imath_\Omega : \pi_0(\Omega_{V_B}) \to \mathrm{Vect}_{G_\chi}(S^2,\chi)_{V_B}$ mapping $[\Phi]$ to $\big[F_{V_B}/\Phi\big]$. This is well-defined by Lemma 5.1, and also surjective because each bundle in $\mathrm{Vect}_{G_\chi}(S^2,\chi)_{V_B}$ can be considered as an equivariant clutching construction. Then, the map $p_\Omega : \pi_0(\Omega_{V_B}) \to A_{G_\chi}(S^2,\chi)$ defined as $p_\Omega = p_{\mathrm{vect}} \circ \imath_\Omega$ satisfies the following diagram:

$$\begin{tikzcd}
\pi_0(\Omega_{V_B}) \arrow[r, "\imath_\Omega"] \arrow[dr, "p_\Omega"'] & \mathrm{Vect}_{G_\chi}(S^2,\chi)_{V_B} \arrow[d, "p_{\mathrm{vect}}"] \\
 & A_{G_\chi}(S^2,\chi)
\end{tikzcd} . \tag{5.2}$$

Let $p_{\pi_0} : \Omega_{V_B} \to \pi_0(\Omega_{V_B})$ be the natural quotient map. For different elements in $A_{G_\chi}(S^2,\chi)$, their preimages through $(p_\Omega \circ p_{\pi_0})^{-1}$ do not intersect each other so that we obtain a decomposition of Ω_{V_B}. We describe this decomposition more precisely. For each $(W_{d^i})_{i\in I^+} \in A_{G_\chi}(S^2,\chi)$, put

$$V_B = G_\chi \times_{(G_\chi)_{d^{-1}}} W_{d^{-1}}, \quad F_{V_B} = G_\chi \times_{(G_\chi)_{d^{-1}}} (|\bar{f}^{-1}| \times W_{d^{-1}}),$$

and by using these define $\Omega_{(W_{d^i})_{i\in I^+}}$ as the subset $(p_\Omega \circ p_{\pi_0})^{-1}\big((W_{d^i})_{i\in I^+}\big)$ of Ω_{V_B}. Henceforward, we will use these V_B and F_{V_B} whenever we deal with $\Omega_{(W_{d^i})_{i\in I^+}}$.

Then, given a bundle V_B, the set Ω_{V_B} is equal to the disjoint union

$$\bigcup_{(W_{d^i})_{i\in I^+}\in A_{G_\chi}(S^2,\chi) \text{ with } W_{d^{-1}}=V_{\bar{f}^{-1}}} \Omega_{(W_{d^i})_{i\in I^+}}.$$

Since Ω_{V_B} and $\Omega_{\hat{D}_R,V_B}$ are in one-to-one correspondence, we may consider $\imath_\Omega$, p_Ω, p_{π_0} as defined also on $\pi_0(\Omega_{\hat{D}_R,V_B})$, $\pi_0(\Omega_{\hat{D}_R,V_B})$, $\Omega_{\hat{D}_R,V_B}$, respectively. So, if we define $\Omega_{\hat{D}_R,(W_{d^i})_{i\in I^+}}$ as the subset $(p_\Omega \circ p_{\pi_0})^{-1}\big((W_{d^i})_{i\in I^+}\big)$ of $\Omega_{\hat{D}_R,V_B}$, then we obtain the decomposition of $\Omega_{\hat{D}_R,V_B}$

$$\bigcup_{(W_{d^i})_{i\in I^+}\in A_{G_\chi}(S^2,\chi) \text{ with } W_{d^{-1}}=V_{\bar{f}^{-1}}} \Omega_{\hat{D}_R,(W_{d^i})_{i\in I^+}}. \tag{5.3}$$

From this decomposition, it suffices to focus on $\Omega_{\hat{D}_R,(W_{d^i})_{i\in I^+}}$ to understand $\Omega_{\hat{D}_R,V_B}$. Now, we state a classification result.

Proposition 5.3.

1. *Assume that $\pi_0(\Omega_{\hat{D}_R,(W_{d^i})_{i\in I^+}})$ is nonempty and $c_1 : \pi_0(\Omega_{\hat{D}_R,(W_{d^i})_{i\in I^+}}) \to H^2(S^2)$, $[\Phi_{\hat{D}_R}] \mapsto c_1(F_{V_B}/\ \Phi_{\hat{D}_R})$ is injective for each $(W_{d^i})_{i\in I^+}$ in $A_{G_\chi}(S^2)$. Then, p_{vect} is surjective, and*

   $$p_{\mathrm{vect}} \times c_1 : \mathrm{Vect}_{G_\chi}(S^2,\chi) \to A_{G_\chi}(S^2,\chi) \times H^2(S^2)$$

   is injective.
2. *Assume that $\pi_0(\Omega_{\hat{D}_R,(W_{d^i})_{i\in I^+}})$ consists of exactly one element for each $(W_{d^i})_{i\in I^+}$ in $A_{G_\chi}(S^2,\chi)$. Then, p_{vect} is an isomorphism.*

Proof. We prove only (1), and (2) is easier. Surjectivity of p_{vect} is trivial by assumption. For arbitrary $[E] \neq [E']$ in $\mathrm{Vect}_{G_\chi}(S^2,\chi)$, if $p_{\mathrm{vect}}([E]) \neq p_{\mathrm{vect}}([E'])$, then there is nothing to prove. Assume that two elements $[E] \neq [E']$ in the set $\mathrm{Vect}_{G_\chi}(S^2,\chi)$ satisfy $p_{\mathrm{vect}}([E]) = p_{\mathrm{vect}}([E']) = (W_{d^i})_{i\in I^+}$ for some $(W_{d^i})_{i\in I^+}$. Then, it suffices to show $c_1(E) \neq c_1(E')$ to prove injectivity. Put $V_B = G_\chi \times_{(G_\chi)_{d^{-1}}} W_{d^{-1}}$. Then, $E|_B \cong E'|_B \cong V_B$ because their isotropy representations at d^{-1} are all $W_{d^{-1}}$. That is, $[E]$ and $[E']$ are in $\mathrm{Vect}_{G_\chi}(S^2,\chi)_{V_B}$. Since $\imath_\Omega$ is surjective, $E \cong F_{V_B}/\Phi$ and $E' \cong F_{V_B}/\Phi'$ for some Φ and Φ' in Ω_{V_B}, especially in $\Omega_{(W_{d^i})_{i\in I^+}}$. By Lemma 5.1, $[\Phi] \neq [\Phi']$ in $\pi_0(\Omega_{(W_{d^i})_{i\in I^+}})$ because $[E] \neq [E']$. So, $c_1\big(F_{V_B}/\Phi\big) \neq c_1\big(F_{V_B}/\Phi'\big)$ by assumption. Therefore, we obtain a proof for injectivity. □ □

By this lemma, we only have to calculate $\pi_0(\Omega_{\hat{D}_R,(W_{d^i})_{i\in I^+}})$ to classify equivariant vector bundles in many cases. In fact, we can apply this lemma except the case when $R = \rho(G_\chi)$ is equal to $\mathbb{Z}_n \times Z$ with odd n or $\langle -a_n \rangle$ with even $n/2$ as we shall see in Section 11. When we can not apply this lemma, we should apply Lemma 5.2 directly.

6. Equivariant pointwise clutching map

Let Φ be an equivariant clutching map determining E in $\mathrm{Vect}_{G_\chi}(S^2,\chi)_{V_B}$, i.e. the map Φ glues F_{V_B} along $|\bar{\mathcal{L}}_R|$ to give E. Let us investigate this gluing process pointwisely. For each $\bar{x} \in |\bar{\mathcal{L}}_R|$ and $x = |\pi|(\bar{x})$, let $\bar{\mathbf{x}} = \pi^{-1}(x) = \{\bar{x}_j | j \in \mathbb{Z}_m\}$ for some m. Then, the map Φ glues the $(G_\chi)_x$-bundle $\big(\mathrm{res}^{G_\chi}_{(G_\chi)_x} F_{V_B}\big)\big|_{\bar{\mathbf{x}}}$ along $\bar{\mathbf{x}}$ to give the $(G_\chi)_x$-representation E_x, and we call this process *equivariant pointwise gluing.* Here, note that $(G_\chi)_{\bar{x}_j} < (G_\chi)_x$ for each $j \in \mathbb{Z}_m$ and

$$\mathrm{res}^{(G_\chi)_x}_{(G_\chi)_{\bar{x}_j}} E_x \cong (F_{V_B})_{\bar{x}_j} \tag{6.1}$$

by equivariance of Φ. In dealing with equivariant clutching maps, technical difficulties occur in equivariant pointwise gluings because gluing by Φ can be considered as just a continuous collection of equivariant pointwise gluings at points in $|\bar{\mathcal{L}}_R|$. In this section, we prove results on equivariant pointwise gluing. To deal with equivariant pointwise gluing, we need the concept of representation extension. For compact Lie groups $N_1 < N_2$, let W_2 be an N_2-representation and W_1 be an N_1-representation. Then, W_2 is called a *representation extension* or an *N_2-extension* of W_1 if $\operatorname{res}_{N_1}^{N_2} W_2 \cong W_1$. For example, E_x is an $(G_\chi)_x$-extension of $(F_{V_B})_{\bar{x}_j}$ for each $j \in \mathbb{Z}_m$ by (6.1). And, let $\operatorname{ext}_{N_1}^{N_2} W_1$ be the set

$$\{W_2 \in \operatorname{Rep}(N_2) \mid \operatorname{res}_{N_1}^{N_2} W_2 \cong W_1\}.$$

Let us investigate equivariant pointwise gluings more precisely under a little bit general setting. Let a compact Lie group N_2 act on a finite set $\bar{\mathbf{x}} = \{\bar{x}_j | \ j \in \mathbb{Z}_m\}$ for $m \geq 2$, and let N_0 and N_1 be the kernel of the action and the isotropy subgroup $(N_2)_{\bar{x}_0}$, respectively. Let F be an N_2-vector bundle over $\bar{\mathbf{x}}$. Consider an arbitrary map

$$\psi : \bar{\mathbf{x}} \to \amalg_{j \in \mathbb{Z}_m} \operatorname{Iso}(F_{\bar{x}_j}, F_{\bar{x}_{j+1}})$$

such that $\psi(\bar{x}_j) \in \operatorname{Iso}(F_{\bar{x}_j}, F_{\bar{x}_{j+1}})$. Call such a map *pointwise preclutching map* with respect to F. By using ψ, we glue $F_{\bar{x}_j}$'s, i.e. a vector u in $F_{\bar{x}_j}$ is identified with $\psi(\bar{x}_j)u$ in $F_{\bar{x}_{j+1}}$ for each j. Let F/ψ be the quotient of F through this identification, and let $p_\psi : F \to F/\psi$ be the quotient map. Let $\imath_\psi : F_{\bar{x}_0} \to F/\psi$ be the composition of the natural injection $\imath_{\bar{x}_0} : F_{\bar{x}_0} \to F$ and the quotient map p_ψ.

(6.2)
$$\begin{tikzcd}
F_{\bar{x}_0} \arrow[r, "\imath_{\bar{x}_0}"] \arrow[d, "\imath_\psi"'] & F \arrow[dl, "p_\psi"] \\
F/\psi &
\end{tikzcd}$$

We would find conditions on ψ under which the quotient F/ψ inherits an N_2-representation structure from F and the map $\imath_\psi$ becomes an N_1-isomorphism from $F_{\bar{x}_0}$ to $\operatorname{res}_{N_1}^{N_2}(F/\psi)$. For notational simplicity, denote

$$\psi(\bar{x}_{j'}) \cdots \psi(\bar{x}_{j+1})\psi(\bar{x}_j)u$$

by $\psi^{j'-j+1}u$ for $u \in F_{\bar{x}_j}$ and $j \leq j'$ in $\mathbb{Z}$.

Lemma 6.1. *For a pointwise preclutching map ψ with respect to F, the quotient F/ψ carries an N_2-representation structure so that p_ψ is N_2-equivariant and $\imath_\psi$ is an N_1-isomorphism if and only if the following conditions hold :*

1. $\psi^m = \mathrm{id}$ *in* $\operatorname{Iso}(F_{\bar{x}_j})$ *for each* $j \in \mathbb{Z}_m$. *So,* ψ^k *is well-defined for all* $k \in \mathbb{Z}_m$.
2. $\psi^{j_3 - j_1} = g\psi(\bar{x}_{j_2})g^{-1}$ *in* $F_{\bar{x}_{j_1}}$ *for each* $j_1 \in \mathbb{Z}_m$, $g \in N_2$ *when* $g^{-1}\bar{x}_{j_1} = \bar{x}_{j_2}$ *and* $g\bar{x}_{j_2+1} = \bar{x}_{j_3}$ *for some* $j_2, j_3 \in \mathbb{Z}_m$.

Proof. To begin with, it is obvious that $\imath_\psi$ is surjective. And, note that $\imath_\psi$ is injective if and only if Condition (1) holds.

Next, we show that F/ψ carries an N_2-representation structure such that p_ψ is equivariant if and only if Condition (2) holds. The possible group action on F/ψ to guarantee equivariance of p_ψ is as follows:

$$g \cdot u = p_\psi(g\bar{u})$$

for each $g \in N_2, u \in F/\psi, \bar{u} \in p_\psi^{-1}(u)$. This is well-defined if and only if $p_\psi(g\bar{u}) = p_\psi(g\psi^l(\bar{u}))$ for each $l \in \mathbb{Z}$, i.e. $\psi^k g\bar{u} = g\psi^l\bar{u}$ for some k. Putting $\bar{u}' = g\bar{u}$, this can be written as $\psi^k\bar{u}' = g\psi^l g^{-1}\bar{u}'$. Since $g\psi^l g^{-1} = (g\psi g^{-1})^l$, this holds for each l if and only if it holds for $l = 1$, i.e. Condition (2). This gives a proof. □ □

A pointwise preclutching map satisfying (1), (2) of Lemma 6.1 is called an *equivariant pointwise clutching map* with respect to F. Let $\mathcal{A}$ be the set of all equivariant pointwise clutching maps with respect to F, and we topologize $\mathcal{A}$ with the subspace topology of $\prod_{j\in\mathbb{Z}_m}\operatorname{Iso}(F_{\bar{x}_j},F_{\bar{x}_{j+1}})$. An N_2-representation W is called *determined* by $\psi\in\mathcal{A}$ with respect to F if $W\cong F/\psi$. In the next corollary, we can see that representation extension is related to equivariant pointwise clutching map and especially guarantees nonemptiness of $\mathcal{A}$.

Corollary 6.2. *For an N_2-extension W of $F_{\bar{x}_0}$, assume that there exists an N_2-morphism $p:F\to W$ such that $p|_{F_{\bar{x}_j}}$ is a nonequivariant isomorphism for each $j\in\mathbb{Z}_m$. Let ψ be the pointwise preclutching map defined by $\psi(\bar{x}_j)=(p|_{F_{\bar{x}_{j+1}}})^{-1}\circ(p|_{F_{\bar{x}_j}})$ for $j\in\mathbb{Z}_m$. Then, ψ is in $\mathcal{A}$, and there is an N_2-isomorphism $\imath:W\to F/\psi$ such that $p_\psi=\imath\circ p$, i.e. ψ determines W with respect to F. Especially, $\mathcal{A}$ is nonempty.*

Proof. Easy check. □ □

To calculate $\pi_0(\Omega_{V_B})$ later, we need to understand topology of $\mathcal{A}$. For this, we would consider F/ψ as an additional structure over the fixed $F_{\bar{x}_0}$ for each $\psi\in\mathcal{A}$ as follows:

Lemma 6.3. *Define a operation $\star_\psi$ of N_2 on $F_{\bar{x}_0}$ as $g\star_\psi u=\imath_\psi^{-1}p_\psi(gu)$ for $\psi\in\mathcal{A}$, $g\in N_2$, $u\in F_{\bar{x}_0}$. Then,*

1. *$\star_\psi$ is an N_2-action on $F_{\bar{x}_0}$ such that $\imath_\psi:(F_{\bar{x}_0},\star_\psi)\to F/\psi$ is an N_2-isomorphism.*
2. *$g\star_\psi u=\psi^{m-j}gu=g\psi^k u$ when $g\bar{x}_0=\bar{x}_j$ and $g\bar{x}_k=\bar{x}_0$. Especially, $g\star_\psi u=gu$ for $g\in N_1$.*

Henceforth, we consider F/ψ as $(F_{\bar{x}_0},\star_\psi)$.

Proof. To prove (1), we show that $\imath_\psi(g\star_\psi u)=g\cdot\imath_\psi(u)$ for each $g\in N_2$, $u\in F_{\bar{x}_0}$ as follows:

$$\begin{aligned}\imath_\psi(g\star_\psi u)&=\imath_\psi(\imath_\psi^{-1}p_\psi(gu))=p_\psi(gu),\\ g\cdot\imath_\psi(u)&=g\cdot p_\psi(u)=p_\psi(gu).\end{aligned}$$

Since F/ψ already delivers an N_2-action and $\imath_\psi$ is bijective, this shows that $\star_\psi$ is an action. Therefore, we prove (1).

Next, we prove (2). Note that $\imath_\psi^{-1}p_\psi|_{F_{\bar{x}_0}}$ is an identity. Using this,

$$g\star_\psi u=\imath_\psi^{-1}p_\psi(gu)=\imath_\psi^{-1}p_\psi(\psi^{m-j}gu)=\psi^{m-j}gu.$$

Also, $\psi^{m-j}gu=g\psi^k u$ by Lemma 6.1.(2). Therefore, we prove (2). □ □

To obtain more precise results on $\mathcal{A}$, we assume that the N_2-bundle F over $\bar{\mathbf{x}}$ satisfies one of the following two conditions:

F1. N_2 fixes $\bar{\mathbf{x}}=\{\bar{x}_j\mid j\in\mathbb{Z}_m\}$ with $m=2$, and $F_{\bar{x}_0}\cong F_{\bar{x}_1}$,
F2. N_2 acts transitively on $\bar{\mathbf{x}}=\{\bar{x}_j\mid j\in\mathbb{Z}_m\}$ with $m\geq 2$.

These conditions are not too restrictive as the following example shows:

Example 6.1. Given the bundle F_{V_B} over $|\bar{\mathcal{K}}_R|$ of Section 4, let $\bar{\mathbf{x}}=\pi^{-1}(x)=\{\bar{x}_j\mid j\in\mathbb{Z}_m\}$ for each $\bar{x}\in|\bar{\mathcal{L}}_R|$, $x=|\pi|(\bar{x})$, some $m\geq 2$. Put $N_2=(G_\chi)_x$ and $F=\big(\operatorname{res}^{G_\chi}_{(G_\chi)_x}F_{V_B}\big)\big|_{\bar{\mathbf{x}}}$. By Table 3.4 and definition of F_{V_B}, we can check that F satisfies Condition F1. or F2. according to $\bar{x}$. Given a map Φ in Ω_{V_B}, we can define a map ψ in $\mathcal{A}$ as follows:

1. if $\bar{x}$ is a vertex, then $\psi_{\bar{x}}(\bar{x}_j)=\Phi(\hat{x}_{j,+})$ for each $j\in\mathbb{Z}_{j_R}$,
2. if $\bar{x}$ is not a vertex and $\bar{x}=p_{|\bar{\mathcal{L}}|}(\hat{x})$, then

$$\psi(\bar{x}_0)=\Phi(\hat{x})\quad\text{and}\quad\psi(\bar{x}_1)=\Phi\big(|c|(\hat{x})\big).$$

Then, we have $F/\psi \cong \big(F_{V_B}/\Phi\big)_x$ for each x. □

Let $\mathcal{A}^j$ be the set

$$\{\psi(\bar{x}_j) \mid \psi \in \mathcal{A}\},$$

and let $\mathcal{A}^{j,j'}$ be the set

$$\{(\psi(\bar{x}_j), \psi(\bar{x}_{j'})) \mid \psi \in \mathcal{A}\}.$$

In the below, it will be witnessed that $\mathcal{A}$ is homeomorphic to $\mathcal{A}^j$ or $\mathcal{A}^{j,j'}$ in many cases.

Lemma 6.4. *Assume that F satisfies Condition F1. Then, $\mathcal{A}$ is homeomorphic to nonempty $\mathcal{A}^0 = \mathrm{Iso}_{N_2}(F_{\bar{x}_0}, F_{\bar{x}_1})$ through the the evaluation map*

$$\mathcal{A} \to \mathcal{A}^0, \ \psi \mapsto \psi(\bar{x}_0).$$

Proof. By Lemma 6.1.(1), each $\psi \in \mathcal{A}$ satisfies $\psi^2 = \mathrm{id}$ so that ψ is determined by $\psi(\bar{x}_0)$. From this, $\mathcal{A}$ is homeomorphic to $\mathcal{A}^0$ through the evaluation. And, ψ satisfies Lemma 6.1.(2) if and only if $\psi(\bar{x}_j)$ is $\mathrm{Iso}_{N_2}(F_{\bar{x}_j}, F_{\bar{x}_{j+1}})$-valued because N_2 fixes $\bar{x}_0, \bar{x}_1$. Nonemptiness is guaranteed by Condition F1. □ □

In the remaining of this section, we assume that F satisfies Condition F2. so that $F \cong N_2 \times_{N_1} F_{\bar{x}_0}$. Under this assumption, the zeroth homotopy of $\mathcal{A}$ will be related to $\mathrm{ext}_{N_1}^{N_2} F_{\bar{x}_0}$. For this, we need a technical lemma. We would express a map ψ in $\mathcal{A}$ as m endomorphisms of $F_{\bar{x}_0}$. This will be useful in dealing with $\mathcal{A}$. For each $j \in \mathbb{Z}_m$, pick an element $g_j \in N_2$ such that $g_j \bar{x}_j = \bar{x}_0$, and denote by $\mathbf{g}$ the m-tuple $(g_j)_{j \in \mathbb{Z}_m}$. Also, express a pointwise preclutching map ψ by the m-tuple $(\psi(\bar{x}_j))_{j \in \mathbb{Z}_m}$. For each $\psi = (\psi(\bar{x}_j))_{j \in \mathbb{Z}_m}$, define the map

$$\psi_{\mathbf{g}} : \bar{\mathbf{x}} \to \mathrm{Iso}(F_{\bar{x}_0}), \ \bar{x}_j \mapsto g_{j+1}\psi(\bar{x}_j)g_j^{-1},$$

and express it by the m-tuple $(g_{j+1}\psi(\bar{x}_j)g_j^{-1})_{j \in \mathbb{Z}_m}$. And, denote by $\mathcal{A}_{\mathbf{g}}$ the set

$$\{ \ \psi_{\mathbf{g}} \mid \psi \in \mathcal{A} \ \}.$$

Here, we consider an action of $\mathrm{Iso}_{N_1}(F_{\bar{x}_0})$ on these $\psi_{\mathbf{g}}$'s. For $B \in \mathrm{Iso}_{N_1}(F_{\bar{x}_0})$, denote by $B\psi_{\mathbf{g}}B^{-1}$ the m-tuple

$$(Bg_{j+1}\psi(\bar{x}_j)g_j^{-1}B^{-1})_{j \in \mathbb{Z}_m}.$$

Then, this action preserves $\mathcal{A}_{\mathbf{g}}$ as follows:

Lemma 6.5. (1) *For $\psi \in \mathcal{A}$ and $B \in \mathrm{Iso}_{N_1}(F_{\bar{x}_0})$, put*

$$\psi'(\bar{x}_j) = g_{j+1}^{-1}Bg_{j+1}\psi(\bar{x}_j)g_j^{-1}B^{-1}g_j$$

for each $j \in \mathbb{Z}_m$ so that ψ' satisfies $B\psi_{\mathbf{g}}B^{-1} = \psi'_{\mathbf{g}}$. Then, $\psi' \in \mathcal{A}$.

(2) *For two elements ψ, ψ' in $\mathcal{A}$, $F/\psi \cong F/\psi'$ as N_2-representation if and only if $B\psi_{\mathbf{g}}B^{-1} = \psi'_{\mathbf{g}}$ for some $B \in \mathrm{Iso}_{N_1}(F_{\bar{x}_0})$.*

Proof. To prove (1), we would show that ψ' satisfies two conditions of Lemma 6.1. It is easy to show that ψ' satisfies Condition (1) of Lemma 6.1. Condition (2) of Lemma 6.1 is written as $(\psi')^{j_3 - j_1}u = g\psi'(\bar{x}_{j_2})g^{-1}u$ in $F_{\bar{x}_{j_1}}$ for $j_1 \in \mathbb{Z}_m$, $g \in N_2$ when $g\bar{x}_{j_2} = \bar{x}_{j_1}$ and $g\bar{x}_{j_2+1} = \bar{x}_{j_3}$. Here,

$$\begin{aligned}
&g\psi'(\bar{x}_{j_2})g^{-1} \\
=&g\Big(g_{j_2+1}^{-1}Bg_{j_2+1}\psi(\bar{x}_{j_2})g_{j_2}^{-1}B^{-1}g_{j_2}\Big)g^{-1} \\
=&\Big(gg_{j_2+1}^{-1}Bg_{j_2+1}g^{-1}\Big)\Big(g\psi(\bar{x}_{j_2})g^{-1}\Big)\Big(gg_{j_2}^{-1}B^{-1}g_{j_2}g^{-1}\Big).
\end{aligned}$$

Note that $gg_{j_2+1}^{-1} = g_{j_3}^{-1}(g_{j_3}gg_{j_2+1}^{-1})$ and $gg_{j_2}^{-1} = g_{j_1}^{-1}(g_{j_1}gg_{j_2}^{-1})$ where $g_{j_3}gg_{j_2+1}^{-1}$ and $g_{j_1}gg_{j_2}^{-1}$ are in N_1. With these,

$$\begin{aligned} gg_{j_2+1}^{-1}Bg_{j_2+1}g^{-1} &= g_{j_3}^{-1}(g_{j_3}gg_{j_2+1}^{-1})B(g_{j_3}gg_{j_2+1}^{-1})^{-1}g_{j_3} \\ &= g_{j_3}^{-1}Bg_{j_3}, \end{aligned}$$

and

$$\begin{aligned} gg_{j_2}^{-1}B^{-1}g_{j_2}g^{-1} &= g_{j_1}^{-1}(g_{j_1}gg_{j_2}^{-1})B^{-1}(g_{j_1}gg_{j_2}^{-1})^{-1}g_{j_1} \\ &= g_{j_1}^{-1}B^{-1}g_{j_1} \end{aligned}$$

because $B \in \mathrm{Iso}_{N_1}(F_{\bar{x}_0})$. So,

$$\begin{aligned} g\psi'(\bar{x}_{j_2})g^{-1} &= \left(g_{j_3}^{-1}Bg_{j_3}\right)\left(g\psi(\bar{x}_{j_2})g^{-1}\right)\left(g_{j_1}^{-1}B^{-1}g_{j_1}\right) \\ &= \left(g_{j_3}^{-1}Bg_{j_3}\right)\psi^{j_3-j_1}\left(g_{j_1}^{-1}B^{-1}g_{j_1}\right) \\ &= \left(g_{j_3}^{-1}Bg_{j_3}\right)\left(g_{j_3}^{-1}B^{-1}g_{j_3}\psi'(\bar{x}_{j_3-1})g_{j_3-1}^{-1}Bg_{j_3-1}\right)\cdots \\ &\qquad \left(g_{j_1+1}^{-1}B^{-1}g_{j_1+1}\psi'(\bar{x}_{j_1})g_{j_1}^{-1}Bg_{j_1}\right)\left(g_{j_1}^{-1}B^{-1}g_{j_1}\right) \\ &= \psi'(\bar{x}_{j_3-1})\cdots\psi'(\bar{x}_{j_1}) \end{aligned}$$

in $F_{\bar{x}_{j_1}}$, and this proves (1).

Next, we prove (2). For sufficiency, assume that $F/\psi \cong F/\psi'$. Considering F/ψ as $(F_{\bar{x}_0}, \star_\psi)$ by Lemma 6.3, there exists an N_2-isomorphism $B : (F_{\bar{x}_0}, \star_\psi) \to (F_{\bar{x}_0}, \star_{\psi'})$, and especially $B \in \mathrm{Iso}_{N_1}(F_{\bar{x}_0})$ by Lemma 6.3.(2). So, $Bg_j \star_\psi g_0^{-1}u = g_j \star_{\psi'} Bg_0^{-1}u$ for each $j \in \mathbb{Z}_m$, $u \in F_{\bar{x}_0}$, and if we substitute $B^{-1}u$ into u, this is written as

$$(Bg_j\psi^j g_0^{-1}B^{-1})u = (g_j{\psi'}^j g_0^{-1})u$$

by Lemma 6.3.(2). Substituting $j = 1$ in this, we obtain $Bg_1\psi g_0^{-1}B^{-1} = g_1\psi' g_0^{-1}$ on $F_{\bar{x}_0}$. By using this and mathematical induction, we would show $B\psi_{\mathbf{g}}B^{-1} = \psi'_{\mathbf{g}}$. Assume that $Bg_j\psi g_{j-1}^{-1}B^{-1} = g_j\psi' g_{j-1}^{-1}$ on $F_{\bar{x}_0}$ from $j = 1$ to $j = k-1$. Then,

$$\begin{aligned} &(g_k\psi' g_{k-1}^{-1})(g_{k-1}\psi' g_{k-2}^{-1})\cdots(g_1\psi' g_0^{-1}) \\ &= g_k{\psi'}^k g_0^{-1} \\ &= Bg_k\psi^k g_0^{-1}B^{-1} \\ &= (Bg_k\psi g_{k-1}^{-1}B^{-1})(Bg_{k-1}\psi g_{k-2}^{-1}B^{-1})\cdots(Bg_1\psi g_0^{-1}B^{-1}) \\ &= (Bg_k\psi g_{k-1}^{-1}B^{-1})(g_{k-1}\psi' g_{k-2}^{-1})\cdots(g_1\psi' g_0^{-1}). \end{aligned}$$

Comparing the first line with the last, we have $Bg_k\psi g_{k-1}^{-1}B^{-1} = g_k\psi' g_{k-1}^{-1}$. By mathematical induction, we obtain $Bg_j\psi g_{j-1}^{-1}B^{-1} = g_j\psi' g_{j-1}^{-1}$ on $F_{\bar{x}_0}$ for each $j \in \mathbb{Z}_m$, and this shows $B\psi_{\mathbf{g}}B^{-1} = \psi'_{\mathbf{g}}$.

Last, we prove necessity of (2). By assumption, we have $Bg_{j+1}\psi g_j^{-1}B^{-1} = g_{j+1}\psi' g_j^{-1}$ for each $j \in \mathbb{Z}_m$ on $F_{\bar{x}_0}$. Then,

$$\begin{aligned} &Bg_j\psi^j g_0^{-1}B^{-1} \\ =&(Bg_j\psi g_{j-1}^{-1}B^{-1})(Bg_{j-1}\psi g_{j-2}^{-1}B^{-1})\cdots(Bg_1\psi g_0^{-1}B^{-1}) \\ =&(g_j\psi' g_{j-1}^{-1})(g_{j-1}\psi' g_{j-2}^{-1})\cdots(g_1\psi' g_0^{-1}) \\ =&g_j{\psi'}^j g_0^{-1} \end{aligned}$$

for each $j \in \mathbb{Z}_m$ on $F_{\bar{x}_0}$. Acting these on Bg_0u for $u \in F_{\bar{x}_0}$, we obtain

$$Bg_j\psi^j u = g_j{\psi'}^j Bu$$

because $B \in \mathrm{Iso}_{N_1}(F_{\bar{x}_0})$. And, this is written as $Bg_j \star_\psi u = g_j \star_{\psi'} Bu$ by Lemma 6.3.(2). This means that B is equivariant for $\langle N_1, g_j | j \in \mathbb{Z}_m \rangle$ so that B is an N_2-isomorphism between $(F_{\bar{x}_0}, \star_\psi)$ and $(F_{\bar{x}_0}, \star_{\psi'})$. Therefore, we obtain proof. □ □

This lemma means that each orbit of $\mathcal{A}_{\mathbf{g}}$ under the $\mathrm{Iso}_{N_1}(F_{\bar{x}_0})$-action corresponds to a representation. More precisely, we have the following:

Theorem 6.6. *Assume that F satisfies Condition F2. Then, the map*

$$\pi_0(\mathcal{A}) \longrightarrow \mathrm{ext}_{N_1}^{N_2} F_{\bar{x}_0}, \quad [\psi] \mapsto F/\psi$$

is bijective. So, $\mathcal{A}$ is nonempty if and only if $F_{\bar{x}_0}$ has an N_2-extension.

Proof. To begin with, we show that each N_2-extension of $F_{\bar{x}_0}$ is obtained as F/ψ for some $\psi \in \mathcal{A}$. Let W be an N_2-extension of $F_{\bar{x}_0}$ such that $\mathrm{res}_{N_1}^{N_2} W$ is equal to $F_{\bar{x}_0}$. Consider a map $p : N_2 \times_{N_1} F_{\bar{x}_0} \to W$, $[g, u] \mapsto gu$. Since $F \cong N_2 \times_{N_1} F_{\bar{x}_0}$, the map p satisfies conditions of Corollary 6.2. Therefore, there exists ψ such that $F/\psi \cong W$ by Corollary 6.2. That is, surjectivity is proved.

Next, we show that two ψ and ψ' in $\mathcal{A}$ are connected by a path in $\mathcal{A}$ if and only if $F/\psi \cong F/\psi'$. First, we show sufficiency. Let ψ_t for $t \in [0,1]$ be a continuous path in $\mathcal{A}$ such that $\psi_0 = \psi$ and $\psi_1 = \psi'$. Then, all $(F_{\bar{x}_0}, \star_{\psi_t})$'s are isomorphic because the path of their characters are in the discrete space $\mathrm{Rep}(N_2)$. Second, we show necessity. If $F/\psi \cong F/\psi'$, then Lemma 6.5 says that $B\psi_{\mathbf{g}}B^{-1} = \psi'_{\mathbf{g}}$ so that $\psi_{\mathbf{g}}$ and $\psi'_{\mathbf{g}}$ are connected in $\mathcal{A}_{\mathbf{g}}$ by a path because $\mathrm{Iso}_{N_1}(F_{\bar{x}_0})$ is a product of general linear groups by Schur's Lemma. Since the map from $\mathcal{A}$ to $\mathcal{A}_{\mathbf{g}}$ sending ψ to $\psi_{\mathbf{g}}$ is a homeomorphism, ψ and ψ' are connected by a path in $\mathcal{A}$, and we obtain necessity. Therefore, we obtain injectivity.

Nonemptiness is clear from the one-to-one correspondence. □ □

Denote by $\mathcal{A}_\psi$ the path-component of $\mathcal{A}$

$$\{\ \psi' \in \mathcal{A} \mid F/\psi \cong F/\psi'\ \},$$

and by $\mathcal{A}_{\psi,\mathbf{g}}$ the path-component of $\mathcal{A}_{\mathbf{g}}$

$$\{\ \psi'_{\mathbf{g}} \in \mathcal{A}_{\mathbf{g}} \mid F/\psi \cong F/\psi'\ \}.$$

Note that $\mathcal{A}_\psi$ and $\mathcal{A}_{\psi,\mathbf{g}}$ are homeomorphic. To calculate homotopy of equivariant clutching maps in next sections, we need to calculate $\pi_1(\mathcal{A}_\psi)$ for each $\psi \in \mathcal{A}$. To do it, we would investigate the shape of $\mathcal{A}_\psi$.

Lemma 6.7. *For each $\psi \in \mathcal{A}$, $\mathcal{A}_\psi$ is homeomorphic to*

$$\mathrm{Iso}_{N_1}(F/\psi) / \mathrm{Iso}_{N_2}(F/\psi).$$

Proof. In Lemma 6.5.(2), we have shown that $\mathcal{A}_{\psi,\mathbf{g}}$ is equal to the orbit of $\psi_{\mathbf{g}}$ by $\mathrm{Iso}_{N_1}(F_{\bar{x}_0})$. To understand this orbit, we need to know the isotropy subgroup $\mathrm{Iso}_{N_1}(F_{\bar{x}_0})_{\psi_{\mathbf{g}}}$, i.e.

$$\{B \in \mathrm{Iso}_{N_1}(F_{\bar{x}_0}) \mid B\psi_{\mathbf{g}}B^{-1} = \psi_{\mathbf{g}}\}.$$

By definition, $B\psi_{\mathbf{g}}B^{-1} = \psi_{\mathbf{g}}$ is expressed as $Bg_{j+1}\psi(\bar{x}_j)g_j^{-1}B^{-1} = g_{j+1}\psi(\bar{x}_j)g_j^{-1}$ for each $j \in \mathbb{Z}_m$. Since $g_{j+1}\psi g_j^{-1} = g_{j+1}g_j^{-1}g_j\psi g_j^{-1} = (g_{j+1}g_j^{-1})\psi^k$ on $F_{\bar{x}_0}$ for some k by Lemma 6.1.(2), this is written as

$$B(g_{j+1}g_j^{-1} \star_\psi B^{-1}u) = g_{j+1}g_j^{-1} \star_\psi u.$$

Since $g_0 \in N_1$, we have $N_2 = \langle N_1, g_{j+1}g_j^{-1} | j \in \mathbb{Z}_m \rangle$. Therefore, $B \in \mathrm{Iso}_{N_1}(F_{\bar{x}_0})_{\psi_{\mathbf{g}}}$ if and only if $B \in \mathrm{Iso}_{N_2}(F_{\bar{x}_0}, \star_\psi)$. Therefore, $A_{\psi,\mathbf{g}}$ is homeomorphic to the quotient $\mathrm{Iso}_{N_1}(F/\psi) / \mathrm{Iso}_{N_2}(F/\psi)$ because $(F_{\bar{x}_0}, \star_\psi) \cong F/\psi$. □ □

In some special cases, we can understand $\mathcal{A}_\psi$ more precisely. Denote by $\mathcal{A}_\psi^j$ the set

$$\{\ \psi'(\bar{x}_j) \mid \psi' \in \mathcal{A}_\psi\ \},$$

and by $\mathcal{A}_\psi^{j,j'}$ the set

$$\{\ \big(\psi'(\bar{x}_j), \psi'(\bar{x}_{j'})\big) \mid \psi' \in \mathcal{A}_\psi\ \}.$$

Proposition 6.8. *Assume that* $N_2 = \langle N_0, a_0\rangle$ *with some* $a_0 \in N_2$ *such that* $a_0\bar{x}_j = \bar{x}_{j+1}$ *for each* $j \in \mathbb{Z}_m$ *so that* $N_2/N_0 \cong \mathbb{Z}_m$ *and* $N_1 = N_0$. *Then,*

1. *A pointwise preclutching map* ψ *with respect to* F *is in* $\mathcal{A}$ *if and only if* $\psi^m = \mathrm{id}$, $\psi(\bar{x}_0) \in \mathrm{Iso}_{N_0}(F_{\bar{x}_0}, F_{\bar{x}_1})$, *and* $\psi(\bar{x}_j) = a_0^j\psi(\bar{x}_0)a_0^{-j}$ *for each* $j \in \mathbb{Z}_m$.
2. $\mathcal{A}, \mathcal{A}_\psi$ *are homeomorphic to* $\mathcal{A}^0, \mathcal{A}_\psi^0$ *for any* $\psi \in \mathcal{A}$, *respectively.*
3. *If* $F_{\bar{x}_0}$ *is* N_0*-isotypical, then* $\mathcal{A}_\psi$ *is simply connected for each* ψ *in* $\mathcal{A}$.

Proof. For (1), we only have to show that ψ satisfies two conditions of Lemma 6.1 if and only if $\psi^m = \mathrm{id}$, $\psi(\bar{x}_0) \in \mathrm{Iso}_{N_0}(F_{\bar{x}_0}, F_{\bar{x}_1})$, and $\psi(\bar{x}_j) = a_0^j\psi(\bar{x}_0)a_0^{-j}$ for $j \in \mathbb{Z}_m$. First, we prove sufficiency. If we substitute each $h \in N_0$ into g in Condition (2) of Lemma 6.1, $\psi(\bar{x}_j)$ is N_0-equivariant for each $j \in \mathbb{Z}_m$. Also, if we substitute a_0^j into g in the same formula, then we obtain $\psi(\bar{x}_j) = a_0^j\psi(\bar{x}_0)a_0^{-j}$. Next, we prove necessity. Note that $\psi(\bar{x}_j) = a_0^j\psi(\bar{x}_0)a_0^{-j}$ is N_0-equivariant for each $j \in \mathbb{Z}_m$ because N_0 is normal in N_2 and $\psi(\bar{x}_0)$ is N_0-equivariant. An arbitrary g in N_2 is expressed as a_0^jh for some $h \in N_0, j \in \mathbb{Z}_m$. Then, since $g^{-1}\bar{x}_{j_1} = \bar{x}_{j_2}$ and $g\bar{x}_{j_2+1} = \bar{x}_{j_3}$ with $j_2 = j_1 - j, j_3 = j_1 + 1$ for each $j_1 \in \mathbb{Z}_m$, we have

$$\begin{aligned} g\psi(\bar{x}_{j_2})g^{-1}u &= (a_0^jh)\psi(\bar{x}_{j_2})(a_0^jh)^{-1}u \\ &= a_0^j\psi(\bar{x}_{j_2})a_0^{-j}u \\ &= a_0^{j+j_2}\psi(\bar{x}_0)a_0^{-j-j_2}u \\ &= a_0^{j_1}\psi(\bar{x}_0)a_0^{-j_1}u \\ &= \psi(\bar{x}_{j_1})u \end{aligned}$$

for each $u \in F_{\bar{x}_{j_1}}$ so that ψ satisfies Condition (2) of Lemma 6.1. Therefore, we obtain a proof of (1).

(2) is easy because a map ψ in $\mathcal{A}$ is determined by $\psi(\bar{x}_0)$.

Before we prove (3), we prove that the N_0-character of $F_{\bar{x}_0}$ is fixed by N_2. Assume that $F_{\bar{x}_0}$ is χ-isotypical for some $\chi \in \mathrm{Irr}(N_0)$. By Theorem 6.6, existence of ψ guarantees existence of an N_2-extension, say V. For each $g \in N_2$, we have

$$^gF_{\bar{x}_0} \cong \mathrm{res}_{gN_0g^{-1}}^{N_2}\ {}^gV \cong \mathrm{res}_{N_0}^{N_2}\ {}^gV \cong \mathrm{res}_{N_0}^{N_2} V \cong F_{\bar{x}_0} \tag{6.3}$$

by using normality of N_0. By Definition 8.1, ${}^gF_{\bar{x}_0}$ is $(g\cdot\chi)$-isotypical so that $g\cdot\chi = \chi$ for each g.

Now, we prove (3). Put $F_{\bar{x}_0} \cong lU$ for some $l \in \mathbb{N}$ and $U \in \mathrm{Irr}(N_0)$. We already know that the character of U is fixed by N_2. Let $\bar{U}$ be an N_2-extension of U whose existence is guaranteed by Theorem 14.1. By Corollary 14.2, F/ψ is isomorphic to $(l_0\bar{U} \otimes \Omega(0)) \oplus \cdots \oplus (l_{m-1}\bar{U} \otimes \Omega(m-1))$ for some l_k's in $\mathbb{Z}$ so that $l = \sum l_k$. Schur's Lemma says that $\mathrm{Iso}_{N_0}(F/\psi) \cong \mathrm{Iso}_{N_0}(lU) \cong \mathrm{GL}(l,\mathbb{C})$ and $\mathrm{Iso}_{N_2}(F/\psi) \cong \mathrm{GL}(l_0,\mathbb{C}) \times \cdots \times \mathrm{GL}(l_{m-1},\mathbb{C})$. So, $\mathrm{Iso}_{N_0}(F/\psi)/\mathrm{Iso}_{N_2}(F/\psi)$ is homeomorphic to $\mathrm{GL}(l,\mathbb{C})/\mathrm{GL}(l_0,\mathbb{C}) \times \cdots \times \mathrm{GL}(l_{m-1},\mathbb{C})$. By long exact sequence of homotopy of a fibration,

$$\begin{aligned} &\longrightarrow \pi_1\Big(\mathrm{GL}(l_0,\mathbb{C}) \times \cdots \times \mathrm{GL}(l_{m-1},\mathbb{C})\Big) \xrightarrow{i} \pi_1\Big(\mathrm{GL}(l,\mathbb{C})\Big) \\ &\longrightarrow \pi_1\Big(\mathrm{GL}(l,\mathbb{C})/\mathrm{GL}(l_0,\mathbb{C}) \times \cdots \times \mathrm{GL}(l_{m-1},\mathbb{C})\Big) \longrightarrow 0. \end{aligned}$$

Here, i is surjective. So, the quotient space is simply connected, and we obtain a proof of (3). □ □

Proposition 6.9. *Assume that* $N_2 = \langle N_0, a_0, b_0\rangle$ *with some* $a_0, b_0 \in N_2$ *such that* $a_0\bar{x}_j = \bar{x}_{j+1}$ *and* $b_0\bar{x}_j = \bar{x}_{-j+1}$ *for each* $j \in \mathbb{Z}_m$. *So,* $N_2/N_0 \cong \mathrm{D}_m$ *and* $N_1 = \langle N_0, b_0a_0\rangle$. *Then,* $\mathcal{A}$, $\mathcal{A}_\psi$ *are homeomorphic to* $\mathcal{A}^0$, $\mathcal{A}^0_\psi$ *for any* $\psi \in \mathcal{A}$, *respectively.*

Proof. If we restrict the action to $\langle N_0, a_0\rangle$, then the proof is the same with Proposition 6.8.(2). □ □

Proposition 6.10. *Assume that* $N_2 = \langle N_0, \alpha_1, \alpha_2, \alpha_3\rangle$ *with some* α_i*'s in* N_2 *such that* $\alpha_1\bar{x}_j = \bar{x}_{-j+1}$, $\alpha_2\bar{x}_j = \bar{x}_{j+2}$, $\alpha_3\bar{x}_j = \bar{x}_{-j+3}$ *for each* $j \in \mathbb{Z}_m$ *with* $m = 4$. *So,* $N_2/N_0 \cong \mathbb{Z}_2 \times \mathbb{Z}_2$ *and* $N_1 = N_0$. *Then,* $\mathcal{A}$, $\mathcal{A}_\psi$ *are homeomorphic to* $\mathcal{A}^{0,3}$, $\mathcal{A}^{0,3}_\psi$ *for any* $\psi \in \mathcal{A}$, *respectively.*

Proof. It suffices to show that each ψ in $\mathcal{A}$ is determined by $\psi(\bar{x}_0)$, $\psi(\bar{x}_3)$. Substituting $g = \alpha_i$ in Lemma 6.1.(2), we obtain

$$\psi(\bar{x}_2) = \alpha_3\psi(\bar{x}_0)\alpha_3^{-1}, \quad \psi(\bar{x}_1) = \alpha_2\psi(\bar{x}_3)\alpha_2^{-1},$$

so we obtain a proof. □ □

We often need to restrict our arguments on $\mathcal{A}$ to $\{\bar{x}_j, \bar{x}_{j'}\}$ as follows: let $\mathcal{A}_{j,j'}$ with $j \neq j'$ be the set of equivariant pointwise clutching maps with respect to the $N_2(\{\bar{x}_j, \bar{x}_{j'}\})$-bundle $\big(\mathrm{res}^{N_2}_{N_2(\{\bar{x}_j,\bar{x}_{j'}\})} F\big)|_{\{\bar{x}_j,\bar{x}_{j'}\}}$ where $N_2(\{\bar{x}_j, \bar{x}_{j'}\})$ is the subgroup preserving $\{\bar{x}_j, \bar{x}_{j'}\}$. Here, each ψ in $\mathcal{A}_{j,j'}$ is defined on $\{\bar{x}_j, \bar{x}_{j'}\}$, and satisfies

$$\psi(\bar{x}_j) \in \mathrm{Iso}(F_{\bar{x}_j}, F_{\bar{x}_{j'}}) \quad \text{and} \quad \psi(\bar{x}_{j'}) \in \mathrm{Iso}(F_{\bar{x}_{j'}}, F_{\bar{x}_j}).$$

We need to know if the restricted bundle $\big(\mathrm{res}^{N_2}_{N_2(\{\bar{x}_j,\bar{x}_{j'}\})} F\big)|_{\{\bar{x}_j,\bar{x}_{j'}\}}$ satisfies Condition F1. or F2. If F satisfies Condition F1., then

$$\bar{\mathbf{x}} = \{\bar{x}_j, \bar{x}_{j'}\} \quad \text{and} \quad \big(\mathrm{res}^{N_2}_{N_2(\{\bar{x}_j,\bar{x}_{j'}\})} F\big)|_{\{\bar{x}_j,\bar{x}_{j'}\}} = F$$

so that the restricted bundle trivially satisfies Condition F1. And, it is trivial that $\mathcal{A} = \mathcal{A}_{j,j'}$. If F satisfies Condition F2., then $\big(\mathrm{res}^{N_2}_{N_2(\{\bar{x}_j,\bar{x}_{j'}\})} F\big)|_{\{\bar{x}_j,\bar{x}_{j'}\}}$ satisfies Condition F2. because $N_2(\{\bar{x}_j, \bar{x}_{j'}\})$ acts transitively on $\{\bar{x}_j, \bar{x}_{j'}\}$. We obtain a useful lemma on $\mathcal{A}_{j,j'}$.

Lemma 6.11. *The map* $\mathrm{res}_{j,j'} : \mathcal{A} \to \mathcal{A}_{j,j'}$, $\psi \mapsto \mathrm{res}_{j,j'}(\psi)$ *is well-defined where*

$$\mathrm{res}_{j,j'}(\psi)(\bar{x}_j) = \psi^{j'-j}(\bar{x}_j), \quad \mathrm{res}_{j,j'}(\psi)(\bar{x}_{j'}) = \psi^{j-j'}(\bar{x}_{j'}).$$

And, we have the isomorphism

$$\mathrm{res}^{N_2}_{N_2(\{\bar{x}_j,\bar{x}_{j'}\})}(F/\psi) \cong \big(\mathrm{res}^{N_2}_{N_2(\{\bar{x}_j,\bar{x}_{j'}\})} F\big)|_{\{\bar{x}_j,\bar{x}_{j'}\}} \Big/ \mathrm{res}_{j,j'}(\psi)$$

for each $\psi \in \mathcal{A}$.

Proof. By Lemma 6.1, we can check that $\mathrm{res}_{j,j'}(\psi)$ is in $\mathcal{A}_{j,j'}$. And, the injection from $\big(\mathrm{res}^{N_2}_{N_2(\{\bar{x}_j,\bar{x}_{j'}\})} F\big)|_{\{\bar{x}_j,\bar{x}_{j'}\}}$ to F induces the isomorphism through $p_{\mathrm{res}_{j,j'}(\psi)}$ and p_ψ. □ □

Since any ψ in $\mathcal{A}$ glues all fibers of F to obtain a single vector space F/ψ, ψ might be considered to glue each pair of fibers of F. That is, ψ determines the function $\bar{\psi}$ defined on $\bar{\mathbf{x}} \times \bar{\mathbf{x}} - \Delta$ sending a pair $(\bar{x}, \bar{x}')$ to the element $\bar{\psi}(\bar{x}, \bar{x}')$ in $\mathrm{Iso}(F_{\bar{x}}, F_{\bar{x}'})$ such that each u in $F_{\bar{x}}$ is identified with $\bar{\psi}(\bar{x}, \bar{x}')u$ in $F_{\bar{x}'}$ by ψ, i.e. $\bar{\psi}(\bar{x}, \bar{x}')$ satisfies $p_\psi(u) = p_\psi(\bar{\psi}(\bar{x}, \bar{x}')u)$ where Δ is the diagonal. Call $\bar{\psi}$ the *saturation* of ψ. Since the index j is not used in defining $\bar{\psi}$, it is often convenient to use $\bar{\psi}$ instead of ψ. Denote by $\bar{\mathcal{A}}$ the set $\{\bar{\psi} |\ \psi \in \mathcal{A}\}$, and call it the *saturation* of $\mathcal{A}$. And, denote $F/\psi, p_\psi$ also by $F/\bar{\psi}, p_{\bar{\psi}}$, respectively.

7. Some lemmas on fundamental groups

In this section we prove three lemmas needed to calculate homotopy of equivariant clutching maps. Two of them are just rewriting of Schur's Lemma. The other is on relative homotopy.

Lemma 7.1. *For $\chi \in \mathrm{Irr}(H)$, let W be a χ-isotypical H-representation. For the natural inclusion $\imath : \mathrm{Iso}_H(W) \to \mathrm{Iso}(W)$, the map*

$$\imath_* : \pi_1(\mathrm{Iso}_H(W)) \to \pi_1(\mathrm{Iso}(W))$$

is injective and equal to the multiplication by $\chi(id)$ up to sign.

Proof. An example is given instead of a detailed proof. Let U_0 be an irreducible H-representation with the character χ. Put $W = 2U_0$. Assume that $\chi(id) = 3$. By Schur's Lemma, $\mathrm{Iso}_H(W) \cong \mathrm{GL}(2, \mathbb{C})$ and $\imath$ is a map from $\mathrm{GL}(2, \mathbb{C})$ to $\mathrm{GL}(6, \mathbb{C})$ such that

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \mapsto \left(\begin{array}{ccc|ccc} a & 0 & 0 & b & 0 & 0 \\ 0 & a & 0 & 0 & b & 0 \\ 0 & 0 & a & 0 & 0 & b \\ \hline c & 0 & 0 & d & 0 & 0 \\ 0 & c & 0 & 0 & d & 0 \\ 0 & 0 & c & 0 & 0 & d \end{array}\right).$$

Then, proof for this case is easily followed. □ □

Lemma 7.2. *For $\chi \in \mathrm{Irr}(H)$, let N_2 be a compact Lie group such that $H \lhd N_2$ and $N_2/H \cong \mathbb{Z}_m$, and let W be an N_2-representation such that $\mathrm{res}_H^{N_2} W$ is χ-isotypical. For the natural inclusion $\imath : \mathrm{Iso}_{N_2}(W) \to \mathrm{Iso}_H(W)$, the map $\imath_* : \pi_1(\mathrm{Iso}_{N_2}(W)) \to \pi_1(\mathrm{Iso}_H(W))$ is surjective.*

Proof. As in (6.3), the character χ is fixed by N_2. Let U be an irreducible H-representation whose character is equal to χ, and let $\bar{U}$ be an N_2-extension of U whose existence is guaranteed by Theorem 14.1. Then, W can be decomposed as $\oplus_{k \in \mathbb{Z}_m} l_k \bar{U}_k$ for some nonnegative integers l_k's by Corollary 14.2 where we denote by $\bar{U}_k$ the representation $(\bar{U} \otimes \Omega(k))$. Then, Schur's Lemma says that

$$\begin{aligned} \mathrm{Iso}_{N_2}(W) &\cong \mathrm{Iso}_{N_2}(l_0 \bar{U}_0) \times \cdots \times \mathrm{Iso}_{N_2}(l_{m-1} \bar{U}_{m-1}), \\ &\cong \mathrm{GL}(l_0, \mathbb{C}) \times \cdots \times \mathrm{GL}(l_{m-1}, \mathbb{C}), \\ \mathrm{Iso}_H(W) &\cong \mathrm{GL}(l, \mathbb{C}) \end{aligned}$$

where $l = \sum l_k$. Since $\imath$ is the natural inclusion from to $\mathrm{GL}(l_0, \mathbb{C}) \times \cdots \times \mathrm{GL}(l_{m-1}, \mathbb{C})$ to $\mathrm{GL}(l, \mathbb{C})$, we obtain a proof. □ □

Here, recall a notation.

Notation 7.3. Let X be a topological space. For two points y_0 and y_1 in X and a path $\gamma : [0, 1] \to X$ such that $\gamma(0) = y_0$ and $\gamma(1) = y_1$, denote by $\overline{\gamma}$ the function defined as

$$\overline{\gamma} : \pi_1(X, y_0) \longrightarrow \pi_1(X, y_1), \quad [\sigma] \mapsto [\gamma^{-1}.\sigma.\gamma].$$

Lemma 7.4. *Let X be a path connected topological space with an abelian $\pi_1(X)$. Let A and B be path connected subspaces of X. Also, let $\imath_1$ and $\imath_2$ denote inclusions from A and B to X, respectively. Pick two points $y_0 \in A$, $y_1 \in B$, and a path $\gamma : [0, 1] \to X$ such that $\gamma(0) = y_0$ and $\gamma(1) = y_1$. Then, we have a one-to-one correspondence*

$$\begin{aligned} \Pi : \pi_1(X, y_1) \Big/ \Big\{ \overline{\gamma}\big(\imath_{1*} \pi_1(A, y_0)\big) + \imath_{2*} \pi_1(B, y_1) \Big\} &\longrightarrow \Big[[0, 1], 0, 1; X, A, B\Big], \\ [\sigma] &\longmapsto [\gamma.\sigma]. \end{aligned}$$

Proof. The only issue is well-definedness of Π and Π^{-1} which is just a tedious check. □ □

8. Equivariant clutching maps on one-dimensional fundamental domain

Assume that $\rho(G_\chi) = R$ for some finite R in Table 1.1 such that $\mathrm{pr}(R) = \mathrm{T}, \mathrm{O}, \mathrm{I}$. In this section, we find conditions on a preclutching map $\Phi_{\hat{D}_R}$ in $C^0(\hat{D}_R, V_B)$ to guarantee that $\Phi_{\hat{D}_R}$ be the restriction of an equivariant clutching map as promised in Section 4. By using this, we show that $\Omega_{\hat{D}_R,(W_{d^i})_{i\in I}}$ is nonempty for each $(W_{d^i})_{i\in I^+} \in A_{G_\chi}(S^2, \chi)$. For these, we define notations on equivariant pointwise clutching maps with respect to the $(G_\chi)_x$-bundle $\big(\mathrm{res}^{G_\chi}_{(G_\chi)_x} F_{V_B}\big)|_{|\pi|^{-1}(x)}$ for each $\bar{x} \in |\bar{\mathcal{L}}_R|$ and $x = |\pi|(\bar{x})$, and prove some lemmas on them. Then, we apply results of Section 6 in dealing with $\Omega_{\hat{D}_R,(W_{d^i})_{i\in I^+}}$. Concrete calculation of homotopy of equivariant clutching maps for each case is done in the next section.

First, we define a set $\mathcal{A}_{\bar{x}}$ of equivariant pointwise clutching maps for each $\bar{x}$ in $|\bar{\mathcal{L}}_R|$. For each $\bar{x}$ in $|\bar{\mathcal{L}}_R|$ and $x = |\pi|(\bar{x})$, put $\bar{\mathbf{x}} = |\pi|^{-1}(x) = \{\bar{x}_j | j \in \mathbb{Z}_m\}$ for $m = j_R$ or 2 according to whether $\bar{x}$ is a vertex or not. Then, let $\mathcal{A}_{\bar{x}}$ be the set of equivariant pointwise clutching maps with respect to the $(G_\chi)_x$-bundle $\big(\mathrm{res}^{G_\chi}_{(G_\chi)_x} F_{V_B}\big)|_{\bar{\mathbf{x}}}$, i.e. $N_2 = (G_\chi)_x$ and $F = \big(\mathrm{res}^{G_\chi}_{(G_\chi)_x} F_{V_B}\big)|_{\bar{\mathbf{x}}}$ in notations of Section 6 (see Example 6.1). Here, we need to explain for codomain of maps in $\mathcal{A}_{\bar{x}}$. For each $\bar{x} \in |\bar{\mathcal{L}}_R|$ and $\psi_{\bar{x}} \in \mathcal{A}_{\bar{x}}$, $\psi_{\bar{x}}(\bar{x}_j)$ is in

$$\mathrm{Iso}\Big((F_{V_B})_{\bar{x}_j}, (F_{V_B})_{\bar{x}_{j+1}}\Big)$$

for $j \in \mathbb{Z}_{j_R}$ or $\mathbb{Z}_2$. If $\bar{x}_j \in |\bar{f}|$ and $\bar{x}_{j+1} \in |\bar{f}'|$ for some $\bar{f}$, $\bar{f}'$, then $\psi_{\bar{x}}(\bar{x}_j)$ is henceforth regarded as in $\mathrm{Iso}\big(V_{\bar{f}}, V_{\bar{f}'}\big)$. This is justified because we have fixed the trivialization $\big(\mathrm{res}^{G_\chi}_{(G_\chi)_{b(\bar{f})}} F_{V_B}\big)\big|_{|\bar{f}|} = |\bar{f}| \times V_{\bar{f}}$ for each face $\bar{f} \in \bar{\mathcal{K}}_R$. We define one more set $\mathcal{A}_{\bar{e}}$ of equivariant pointwise clutching maps for each edge $\hat{e}$ and its images $\bar{e} = p_{\bar{\mathcal{L}}}(\hat{e})$, $|e| = |\pi|(|\bar{e}|)$. For each $\hat{x}$ in $|\hat{e}|$ and $\bar{x} = p_{|\bar{\mathcal{L}}|}(\hat{x})$, put

$$\bar{x}'_0 = \bar{x}, \quad \bar{x}'_1 = p_{|\bar{\mathcal{L}}|}(|c|(\hat{x})), \quad \text{and} \quad \bar{\mathbf{x}}' = \{\bar{x}'_j | j \in \mathbb{Z}_2\}.$$

Consider the set $\mathcal{A}_{\hat{x}}$ of equivariant pointwise clutching maps with respect to the $(G_\chi)_{|e|}$-bundle $\big(\mathrm{res}^{G_\chi}_{(G_\chi)_{|e|}} F_{V_B}\big)|_{\bar{\mathbf{x}}'}$ where $(G_\chi)_{|e|}$ is the subgroup of G_χ fixing $|e|$, i.e. $N_2 = (G_\chi)_{|e|}$ and $F = \big(\mathrm{res}^{G_\chi}_{(G_\chi)_{|e|}} F_{V_B}\big)|_{\bar{\mathbf{x}}'}$ in notations of Section 6. As for $\mathcal{A}_{\bar{x}}$, each map $\psi_{\hat{x}}$ in $\mathcal{A}_{\hat{x}}$ is considered to satisfy

$$\psi_{\hat{x}}(\bar{x}'_0) \in \mathrm{Iso}(V_{\bar{f}}, V_{\bar{f}'}) \quad \text{and} \quad \psi_{\hat{x}}(\bar{x}'_1) \in \mathrm{Iso}(V_{\bar{f}'}, V_{\bar{f}})$$

when $\bar{x}'_0 \in |\bar{f}|$ and $\bar{x}'_1 \in |\bar{f}'|$. Here, observe that $\mathcal{A}_{\hat{x}}$ is in one-to-one correspondence with $\mathcal{A}_{\hat{y}}$ for any two $\hat{x}, \hat{y}$ in $|\hat{e}|$, i.e. an element $\psi_{\hat{x}}$ in $\mathcal{A}_{\hat{x}}$ and an element $\psi_{\hat{y}}$ in $\mathcal{A}_{\hat{y}}$ are corresponded to each other when $\psi_{\hat{x}}(\bar{x}'_j) = \psi_{\hat{y}}(\bar{y}'_j)$ for $j \in \mathbb{Z}_2$. This is because the $(G_\chi)_{|e|}$-bundle $\big(\mathrm{res}^{G_\chi}_{(G_\chi)_{|e|}} F_{V_B}\big)|_{\bar{\mathbf{x}}'}$ is all isomorphic regardless of $\hat{x} \in |\hat{e}|$. It is very useful to identify all $\mathcal{A}_{\hat{x}}$'s for $\hat{x} \in |\hat{e}|$ in this way, so we denote the identified set by $\mathcal{A}_{\bar{e}}$. That is, each element $\psi_{\bar{e}}$ in $\mathcal{A}_{\bar{e}}$ is considered as contained in $\mathcal{A}_{\hat{x}}$ for any $\hat{x} \in |\hat{e}|$ according to the context.

Next, we would define a G_χ-action on saturations. First, we define notations on saturations. For each $x = |\pi|(\bar{x})$ and $\psi_{\bar{x}} \in \mathcal{A}_{\bar{x}}$, denote saturations of $\mathcal{A}_{\bar{x}}$ and $\psi_{\bar{x}}$ by $\bar{\mathcal{A}}_x$ and $\bar{\psi}_x$, respectively. Since index set is irrelevant in defining $\bar{\mathcal{A}}_x$, the saturation depends not on $\bar{x}$ but on x. This is why we use the subscript x instead of $\bar{x}$. For any $g \in G_\chi$, the function $g \cdot \bar{\psi}_x$ is contained in $\bar{\mathcal{A}}_{gx}$ where $g \cdot \bar{\psi}_x$ is defined as

$$(g \cdot \bar{\psi}_x)(\bar{y}, \bar{y}') = g\bar{\psi}_x(g^{-1}\bar{y}, g^{-1}\bar{y}')g^{-1}$$

for any $\bar{y} \neq \bar{y}'$ in $\pi^{-1}(gx)$. That is, we obtain $g\bar{\mathcal{A}}_x = \bar{\mathcal{A}}_{gx}$. Especially, it is easily shown that $g \cdot \bar{\psi}_x = \bar{\psi}_x$ for each $g \in (G_\chi)_x$ by equivariance of $\psi_{\bar{x}}$. From this, it is noted that if $g'x = gx$ for some $g', g \in G_\chi$, then $g' \cdot \bar{\psi}_x = g \cdot \bar{\psi}_x$ because $g' = g(g^{-1}g')$ with $g^{-1}g' \in (G_\chi)_x$, i.e. $g \cdot \bar{\psi}_x$ is dependent on gx. We have defined a G_χ-action on saturations. Since $\mathcal{A}_{\bar{x}}$ and $\bar{\mathcal{A}}_x$ are in one-to-one correspondence, $\mathcal{A}_{\bar{x}}$'s also deliver the G_χ-action induced from the G_χ-action on $\bar{\mathcal{A}}_x$'s, i.e. $g \cdot \psi_{\bar{x}} \in \mathcal{A}_{g\bar{x}}$ for each $\psi_{\bar{x}} \in \mathcal{A}_{\bar{x}}$ is defined and $\mathcal{A}_{g\bar{x}} = g\mathcal{A}_{\bar{x}}$. Here, we prove a useful lemma on this action. Before it, we explain for the superscript g, and state an elementary fact.

Definition 8.1. Let K be a closed subgroup of a compact Lie group G. For a given element $g \in G$ and $W \in \mathrm{Rep}(K)$, the gKg^{-1}-representation ${}^gW$ is defined to be the vector space W with the new gKg^{-1}-action

$$gKg^{-1} \times W \to W, \quad (k, u) \mapsto g^{-1}kgu$$

for $k \in gKg^{-1}$, $u \in W$.

Lemma 8.2. *Let G be a compact Lie group acting on a topological space X. And, let E be an equivariant vector bundle over X. Then, ${}^gE_x \cong E_{gx}$ for each $g \in G$ and $x \in X$. Also,*

$$\mathrm{res}^{G_x}_{G_x \cap G_{x'}} E_x \cong \mathrm{res}^{G_{x'}}_{G_x \cap G_{x'}} E_{x'}$$

for any two points x, x' in the same component of the fixed set $X^{G_x \cap G_{x'}}$.

By Lemma 8.2 and Lemma 3.10, p_{vect} is well-defined in cases when $\mathrm{pr}(R) = \mathrm{T}, \mathrm{O}, \mathrm{I}$.

Lemma 8.3. *For each $\bar{x} \in |\bar{\mathcal{L}}_R|$, $x = |\pi|(\bar{x})$, $g \in G_\chi$, $\psi_{\bar{x}} \in \mathcal{A}_{\bar{x}}$, we have an $(G_\chi)_{gx}$-isomorphism*

$$\big(\mathrm{res}^{G_\chi}_{(G_\chi)_{gx}} F_{V_B}\big)|_{g\bar{\mathbf{x}}} \big/ g \cdot \bar{\psi}_x \cong {}^g\Big(\big(\mathrm{res}^{G_\chi}_{(G_\chi)_x} F_{V_B}\big)|_{\bar{\mathbf{x}}} \big/ \bar{\psi}_x \Big).$$

Proof. Put $L = \langle g, (G_\chi)_x \rangle$, and let $k_0 \in \mathbb{N}$ be the smallest natural number satisfying $g^{k_0} \in (G_\chi)_x$ where such a number exists because $R = \rho(G_\chi)$ is finite. To prove this lemma by using Lemma 8.2, we would construct an L-bundle $\mathcal{F}$ over the orbit $Lx = \{x, gx, \cdots, g^{k_0-1}x\}$. Put $\bar{\mathcal{F}} = \big(\mathrm{res}^{G_\chi}_L F_{V_B}\big)|_{L\bar{\mathbf{x}}}$ over $L\bar{\mathbf{x}}$, and consider the nonequivariant bundle $\mathcal{F}$ over the orbit Lx whose fiber $\mathcal{F}_{g^kx}$ at g^kx is equal to $\bar{\mathcal{F}}|_{g^k\bar{\mathbf{x}}}/(g^k \cdot \bar{\psi}_x)$ for $k = 0, \cdots, k_0 - 1$. And, let $P : \bar{\mathcal{F}} \to \mathcal{F}$ be the map such that the restriction of P to $\bar{\mathcal{F}}|_{g^k\bar{\mathbf{x}}}$ is equal to $p_{g^k \cdot \bar{\psi}_x}$ where $p_{g^k \cdot \bar{\psi}_x} : \bar{\mathcal{F}}|_{g^k\bar{\mathbf{x}}} \to \bar{\mathcal{F}}|_{g^k\bar{\mathbf{x}}}/(g^k \cdot \bar{\psi}_x)$ is the quotient map of (6.2). Then, we would define an L-action on $\mathcal{F}$ as $lu = P(l\bar{u})$ for each $l \in L, u \in \mathcal{F}$, and any $\bar{u} \in P^{-1}(u)$ so that P becomes L-equivariant. As long as this is well-defined, it is easily shown that it becomes an action because it is defined by the L-action on $\bar{\mathcal{F}}$ through P. So, we would prove well-definedness of the action. For this, it suffices to show $P(l\bar{u}) = P(l\bar{u}')$ for each $l \in L$ and each $\bar{u}, \bar{u}'$ in $\bar{\mathcal{F}}$ satisfying $P(\bar{u}) = P(\bar{u}')$. If $\bar{u} \in (F_{V_B})_{g^k\bar{x}_j}$ and $\bar{u}' \in (F_{V_B})_{g^k\bar{x}_{j'}}$ for some j, j', then $P(\bar{u}) = P(\bar{u}')$ is written as

$$(g^k \cdot \bar{\psi}_x)(g^k\bar{x}_j, g^k\bar{x}_{j'})\bar{u} = \bar{u}'. \tag{*}$$

Note that $l\bar{u} \in (F_{V_B})_{lg^k\bar{x}_j}$ and $l\bar{u}' \in (F_{V_B})_{lg^k\bar{x}_{j'}}$. And, put $l = g^{k'}l'$ with $l' \in L_{g^kx}$ and some integer k' so that $lg^k\bar{\mathbf{x}} = g^{k+k'}\bar{\mathbf{x}}$ because l' fixes $g^k\bar{\mathbf{x}}$. Remembering that the restriction of P to $\bar{\mathcal{F}}|_{lg^k\bar{\mathbf{x}}} = \bar{\mathcal{F}}|_{g^{k+k'}\bar{\mathbf{x}}}$ is equal to $p_{g^{k+k'} \cdot \bar{\psi}_x}$, $P(l\bar{u}) = P(l\bar{u}')$ is shown as

$$\begin{aligned} &(g^{k+k'}\cdot\bar{\psi}_x)(lg^k\bar{x}_j, lg^k\bar{x}_{j'})l\bar{u}\\ =&g^{k'}(g^k\cdot\bar{\psi}_x)(g^{-k'}lg^k\bar{x}_j, g^{-k'}lg^k\bar{x}_{j'})g^{-k'}l\bar{u}\\ =&g^{k'}l'l'^{-1}(g^k\cdot\bar{\psi}_x)(l'g^k\bar{x}_j, l'g^k\bar{x}_{j'})l'\bar{u}\\ =&g^{k'}l'(l'^{-1}g^k\cdot\bar{\psi}_x)(g^k\bar{x}_j, g^k\bar{x}_{j'})\bar{u}\\ =&l(g^k\cdot\bar{\psi}_x)(g^k\bar{x}_j, g^k\bar{x}_{j'})\bar{u}\\ =&l\bar{u}'\end{aligned}$$

where we use (*) in the last line. So, we obtain well-definedness of L-action on $\mathcal{F}$. By definition, isotropy representations $\mathcal{F}_x$ and $\mathcal{F}_{gx}$ are equal to representations

$$\big(\mathrm{res}^{G_\chi}_{(G_\chi)_x} F_{V_B}\big)|_{\bar{\mathbf{x}}}/\bar{\psi}_x \quad \text{and} \quad \big(\mathrm{res}^{G_\chi}_{(G_\chi)_{gx}} F_{V_B}\big)|_{g\bar{\mathbf{x}}}\Big/ g\cdot\bar{\psi}_x,$$

respectively. Then, the lemma follows from Lemma 8.2. □ □

To investigate $\Omega_{\hat{D}_R,(W_{d^i})_{i\in I^+}}$, we need prove a basic lemma on relations between $\mathcal{A}_{\bar{e}^i}$ and $\mathcal{A}_{\bar{x}}$ for $\bar{x}=\bar{v}^i$ or $\bar{v}^{i+1}$. Also, we prove lemmas on evaluation of equivariant pointwise clutching maps.

Lemma 8.4. *Put* $\bar{\mathbf{v}}^k = |\pi|^{-1}(v^k) = \{\bar{x}_j \mid \bar{x}_j = \bar{v}^k_j \text{ for } j\in\mathbb{Z}_{j_R}\}$ *for* $k=i,i+1$. *And, put* $\bar{\mathbf{v}}'^i=\{\bar{v}^i_0,\bar{v}^i_1\}$ *and* $\bar{\mathbf{v}}'^{i+1}=\{\bar{v}^{i+1}_0,\bar{v}^{i+1}_{-1}\}$ *so that* $\bar{v}^i_1$, $\bar{v}^{i+1}_{-1}\in c(\bar{e}^i)$.

1. $\mathcal{A}_{\bar{x}}\subset\mathcal{A}_{\bar{e}^i}$ *for each interior* $\bar{x}$ *in* $|\bar{e}^i|$.
2. $\mathcal{A}_{\bar{e}^i}=\mathcal{A}_{\bar{x}}$ *for each interior* $\bar{x}\neq b(\bar{e}^i)$ *in* $|\bar{e}^i|$. *Moreover,* $\mathcal{A}_{\bar{e}^i}=\mathcal{A}_{b(\bar{e}^i)}$ *if* $(G_\chi)_{b(e^i)}=(G_\chi)_x$ *for* $x=|\pi|(\bar{x})$.
3. $\mathcal{A}^0_{\bar{v}^i}\subset\mathcal{A}^0_{\bar{e}^i}$ *and* $\mathcal{A}^{j_R-1}_{\bar{v}^{i+1}}\subset\mathcal{A}^1_{\bar{e}^i}$.
4. *For each* $\psi_{\bar{v}^i}$ *in* $\mathcal{A}_{\bar{v}^i}$, *we have* $\psi_{\bar{v}^i}(\bar{v}^i_0)=\psi_{\bar{e}^i}(\bar{v}^i_0)$ *for the unique* $\psi_{\bar{e}^i}\in\mathcal{A}_{\bar{e}^i}$, *and*
$$\mathrm{res}^{(G_\chi)_{v^i}}_{(G_\chi)_{|e^i|}}\Big\{\big(\mathrm{res}^{G_\chi}_{(G_\chi)_{v^i}} F_{V_B}\big)|_{\bar{\mathbf{v}}^i}\Big/\psi_{\bar{v}^i}\Big\}\cong\big(\mathrm{res}^{G_\chi}_{(G_\chi)_{|e^i|}} F_{V_B}\big)|_{\bar{\mathbf{v}}'^i}\Big/\psi_{\bar{e}^i}.$$
5. *For each* $\psi_{\bar{v}^{i+1}}$ *in* $\mathcal{A}_{\bar{v}^{i+1}}$, *we have* $\psi_{\bar{v}^{i+1}}(\bar{v}^{i+1}_{-1})=\psi_{\bar{e}^i}(\bar{v}^{i+1}_{-1})$ *for the unique* $\psi_{\bar{e}^i}\in\mathcal{A}_{\bar{e}^i}$, *and*
$$\mathrm{res}^{(G_\chi)_{v^{i+1}}}_{(G_\chi)_{|e^i|}}\Big\{\big(\mathrm{res}^{G_\chi}_{(G_\chi)_{v^{i+1}}} F_{V_B}\big)|_{\bar{\mathbf{v}}^{i+1}}\Big/\psi_{\bar{v}^{i+1}}\Big\}\cong\big(\mathrm{res}^{G_\chi}_{(G_\chi)_{|e^i|}} F_{V_B}\big)|_{\bar{\mathbf{v}}'^{i+1}}\Big/\psi_{\bar{e}^i}.$$

Proof. (1) follows from $(G_\chi)_{|e^i|}\subset(G_\chi)_x$ when $x=|\pi|(\bar{x})$. (2) follows from $(G_\chi)_{|e^i|}=(G_\chi)_x$. Similarly, since $(G_\chi)_{|e^i|}\subset G_\chi(\bar{\mathbf{v}}'^i)$ and $(G_\chi)_{|e^i|}\subset G_\chi(\bar{\mathbf{v}}'^{i+1})$, (3) holds by Lemma 6.11 where $G_\chi(\bar{\mathbf{v}}'^i)$ and $G_\chi(\bar{\mathbf{v}}'^{i+1})$ are subgroups of G_χ preserving $\bar{\mathbf{v}}'^i$ and $\bar{\mathbf{v}}'^{i+1}$, respectively. The first statement of (4) follows by (3) and Lemma 6.4. By Lemma 6.11, we have the second statement of (4). Similarly, (5) is also obtained. □ □

Lemma 8.5. *Assume that* $\mathrm{pr}(R)=\mathrm{T}$, O, I *and* $R\neq\mathrm{T}\times Z$. *For each vertex* $\bar{v}$ *in* $\bar{\mathcal{K}}_R$, *the evaluation map* $\mathcal{A}_{\bar{v}}\to\mathcal{A}^0_{\bar{v}}$, $\psi_{\bar{v}}\mapsto\psi_{\bar{v}}(\bar{v})$ *is homeomorphic.*

Proof. Put $v=|\pi|(\bar{v})$. In these cases, $R_v\cong\mathbb{Z}_m$ or D_m with $m=j_R$ by Table 3.4, and proof is done by Proposition 6.8.(2) and 6.9. □ □

Lemma 8.6. *Assume that* $R=\mathrm{T}\times Z$. *For each vertex* $\bar{v}$ *in* $\bar{\mathcal{K}}_R$, *the evaluation map*

$$\mathcal{A}_{\bar{v}}\to\mathcal{A}^0_{\bar{v}}\times\mathcal{A}^{-1}_{\bar{v}},\ \psi_{\bar{v}}\mapsto\big(\ \psi_{\bar{v}}(\bar{v}_0),\ \psi_{\bar{v}}(\bar{v}_{-1})\ \big)$$

is homeomorphic.

Proof. Put $v=|\pi|(\bar{v})$. In these cases, $R_v\cong\mathbb{Z}_2\times\mathbb{Z}_2$ by Table 3.4, and proof is done by Proposition 6.10. □ □

Now, we state conditions on a preclutching map $\Phi_{\hat{D}_R}$ in $C^0(\hat{D}_R, V_B)$ to guarantee that $\Phi_{\hat{D}_R}$ be the restriction of an equivariant clutching map. When $R = \mathrm{T} \times Z$, pick an element $t_0 \in (G_\chi)_{b(f^{-1})}$ such that $t_0 v^0 = v^1$. So, t_0 satisfies $t_0 \bar{v}^0_j = \bar{v}^1_j$ and $t_0 \bar{v}^0_{j,\pm} = \bar{v}^1_{j,\pm}$ for $j \in \mathbb{Z}_4$ as illustrated in Figure 4.2. Also, we define a terminology.

Definition 8.7. For $\bar{x} \in |\bar{\mathcal{L}}_R|$, $x = |\pi|(\bar{x})$, $\psi_{\bar{x}} \in \mathcal{A}_{\bar{x}}$, $\Phi \in C^0(|\hat{\mathcal{L}}_R|, V_B)$, the map $\psi_{\bar{x}}$ or its saturation $\bar{\psi}_x$ is called *determined* by Φ if Φ satisfies the following condition:

$$\bar{\psi}_x\Big(p_{|\bar{\mathcal{L}}|}(\hat{x}), p_{|\bar{\mathcal{L}}|}(|c|(\hat{x}))\Big) = \Phi(\hat{x})$$

for each $\hat{x} \in (|\pi| \circ p_{|\bar{\mathcal{L}}|})^{-1}(x)$. The condition is concretely written as

(1) if $\bar{x}$ is a vertex, then

$$\psi_{\bar{x}}(\bar{x}_j) = \Phi(\hat{x}_{j,+}) \quad \text{and} \quad \psi_{\bar{x}}^{-1}(\bar{x}_j) = \Phi(\hat{x}_{j,-})$$

for each $j \in \mathbb{Z}_{j_R}$,

(2) if $\bar{x}$ is not a vertex, then

$$\psi_{\bar{x}}(\bar{x}_0) = \Phi(\hat{x}_{0,+}) \quad \text{and} \quad \psi_{\bar{x}}(\bar{x}_1) = \Phi(|c|(\hat{x}_{0,+})).$$

Theorem 8.8. *Assume that* $\mathrm{pr}(R) = \mathrm{T}, \mathrm{O}, \mathrm{I}$. *Then, a preclutching map* $\Phi_{\hat{D}_R}$ *in* $C^0(\hat{D}_R, V_B)$ *is in* $\Omega_{\hat{D}_R, V_B}$ *if and only if there exists the unique* $\psi_{\bar{x}} \in \mathcal{A}_{\bar{x}}$ *for each* $\bar{x} \in \bar{D}_R$ *and* $x = |\pi|(\bar{x})$ *satisfying*

E2. $\bar{\psi}_x\Big(p_{|\bar{\mathcal{L}}|}(\hat{x}), p_{|\bar{\mathcal{L}}|}(|c|(\hat{x}))\Big) = \Phi_{\hat{D}_R}(\hat{x})$ *for each* $\hat{x} \in \hat{D}_R$,

E3. *for each* $\bar{x}, \bar{x}' \in \bar{D}_R$ *and their images* $x = |\pi|(\bar{x}), x' = |\pi|(\bar{x}')$, *if* $x' = gx$ *for some* $g \in G_\chi$, *then* $\bar{\psi}_{x'} = g \cdot \bar{\psi}_x$.

The set $(\psi_{\bar{x}})_{\bar{x} \in \bar{D}_R}$ *is called determined by* $\Phi_{\hat{D}_R}$.

Proof. For necessity, it suffices to construct a map Φ in Ω_{V_B} such that $\Phi|_{\hat{D}_R} = \Phi_{\hat{D}_R}$. For this, we would define Φ on $\hat{\mathbf{D}}_R$, and then extend the domain of definition of Φ to the whole $|\hat{\mathcal{L}}_R|$. First, we define Φ on $\hat{\mathbf{D}}_R$ so that each $\psi_{\bar{x}}$ is determined by Φ, i.e. each $\psi_{\bar{x}}$ and Φ satisfy Definition 8.7. Then, Condition E2. says that $\Phi|_{\hat{D}_R} = \Phi_{\hat{D}_R}$. Next, we define $\Phi(\hat{x}) = g^{-1}\Phi(g\hat{x})g$ for each $\hat{x} \in |\hat{\mathcal{L}}_R|$ and some $g \in G_\chi$ such that $g\hat{x}$ is in $\hat{\mathbf{D}}_R$. We need prove well-definedness of this. Assume that $\hat{y} = g\hat{x}$ and $\hat{y}' = g'\hat{x}$ are in $\hat{\mathbf{D}}_R$ for two elements g, g' in G_χ so that $\hat{y}' = g'g^{-1}\hat{y}$. And, let y and y' be images of $\hat{y}$ and $\hat{y}'$ through $|\pi| \circ p_{|\bar{\mathcal{L}}|}$, respectively. Then, $y' = g'g^{-1}y$. These give us $\bar{\psi}_{y'} = (g'g^{-1}) \cdot \bar{\psi}_y$ by Condition E3. From this, we obtain

$$\begin{aligned} g'^{-1}\Phi(\hat{y}')g' &= g'^{-1}\bar{\psi}_{y'}\Big(p_{|\bar{\mathcal{L}}|}(\hat{y}'), p_{|\bar{\mathcal{L}}|}(|c|(\hat{y}'))\Big)g' \\ &= g^{-1}\bar{\psi}_y\Big(p_{|\bar{\mathcal{L}}|}(\hat{y}), p_{|\bar{\mathcal{L}}|}(|c|(\hat{y}))\Big)g \\ &= g^{-1}\Phi(\hat{y})g \end{aligned}$$

where we use equivariance of $|c|$. So, well-definedness is proved. It is easily checked that Φ satisfies Condition N1., N2., E1. Therefore, Φ is the wanted equivariant clutching map.

For sufficiency, assume that $\Phi_{\hat{D}_R} = \Phi|_{\hat{D}_R}$ for some $\Phi \in \Omega_{V_B}$. Then, we should choose the unique $\psi_{\bar{x}} \in \mathcal{A}_{\bar{x}}$ for each $\bar{x} \in \bar{D}_R$ satisfying Condition E2. and E3. When we show that $\psi_{\bar{x}}$'s satisfy Condition E3., the condition $x' = gx$ holds only if x', x are equal points or x', x are different vertices by definition of one-dimensional fundamental domain. The second situation happens only in the case of $R = \mathrm{T} \times Z$ by Table 1.1, and in other cases D_R contains only one vertex. So, proof for sufficiency is different according to R.

First, assume that $R \neq \mathrm{T} \times Z$. At each $\bar{x} \in \bar{D}_R$, the unique $\psi_{\bar{x}}$ in $\mathcal{A}_{\bar{x}}$ is determined by Φ because Φ satisfies Condition N1., N2., E1. Moreover, it can be checked that

$\psi_{\bar{x}}$ is the unique element satisfying Condition E2. for each $\bar{x}$ by Lemma 8.5. And, as we have seen in the above, we do not need to consider Condition E3. in these cases.

Second, assume that $R = \mathrm{T} \times Z$. At each $\bar{x} \in \bar{D}_R$, the unique $\psi_{\bar{x}}$ in $\mathcal{A}_{\bar{x}}$ is determined by Φ because Φ satisfies Condition N1., N2., E1. Easily, these $\psi_{\bar{x}}$'s satisfy Condition E2., E3. so that it remains to show their uniqueness. For $\bar{x} \in \bar{D}_R - \{\bar{v}^0, \bar{v}^1\}$, $\psi_{\bar{x}}$ is the unique element satisfying Condition E2. by Lemma 6.4. Condition E2. says that

$$(*) \qquad \psi_{\bar{v}^0}(\bar{v}^0) = \Phi_{\hat{D}_R}(\hat{v}^0_+), \quad \psi^{-1}_{\bar{v}^1}(\bar{v}^1) = \Phi_{\hat{D}_R}(\hat{v}^1_-).$$

And, Condition E3. says that $\bar{\psi}_{v^1} = t_0 \cdot \bar{\psi}_{v^0}$, and from this it is obtained that

$$(**) \qquad \psi_{\bar{v}^1}(\bar{v}^1) = t_0 \Phi_{\hat{D}_R}(\hat{v}^0_+) t_0^{-1}.$$

Formulas (*), (**) say that two values $\psi_{\bar{v}^1}(\bar{v}^1)$ and $\psi_{\bar{v}^1}(\bar{v}^1_{-1}) = (\psi^{-1}_{\bar{v}^1}(\bar{v}^1))^{-1}$ are determined by $\Phi_{\hat{D}_R}$. And, this means that $\psi_{\bar{v}^1}$ is unique by Lemma 8.6. By Condition E3., uniqueness of $\psi_{\bar{v}^0}$ is also obtained. □ □

Remark 8.9. This theorem holds even though we might omit the word 'unique' in the statement of the theorem because uniqueness is not used in the proof of necessity. □

By using Theorem 8.8, we would describe $\Omega_{\hat{D}_R, V_B}$ through $\mathcal{A}_{\bar{x}}$'s. Define the following set of equivariant pointwise clutching maps on $\bar{d}^i$'s.

Definition 8.10. Denote by $\bar{A}_{G_\chi}(S^2, V_B)$ the set

$$\{(\psi_{\bar{d}^i})_{i \in I} \mid \psi_{\bar{d}^i} \in \mathcal{A}_{\bar{d}^i} \text{ and } \bar{\psi}_{d^1} = g \cdot \bar{\psi}_{d^0} \text{ if } d^1 = g d^0 \text{ for some } g \in G_\chi\}.$$

An element $(\psi_{\bar{d}^i})_{i \in I}$ in $\bar{A}_{G_\chi}(S^2, V_B)$ is *determined* by $\Phi_{\hat{D}_R} \in \Omega_{\hat{D}_R, V_B}$ if $\psi_{\bar{d}^i}$'s and $\Phi_{\hat{D}_R}$ satisfy Condition E2. and E3. of Theorem 8.8. Also, a triple $(W_{d^i})_{i \in I^+}$ in $A_{G_\chi}(S^2, \chi)$ is *determined* by $(\psi_{\bar{d}^i})_{i \in I}$ in $\bar{A}_{G_\chi}(S^2, V_B)$ if $W_{d^{-1}} \cong V_{\bar{f}^{-1}}$ and W_{d^i} is determined by $\psi_{\bar{d}^i}$ with respect to $\big(\mathrm{res}^{G_\chi}_{(G_\chi)_{d^i}} F_{V_B}\big)|_{|\pi|^{-1}(d^i)}$ for each $i \in I$.

By Theorem 8.8, we can see that for each $\Phi_{\hat{D}_R} \in \Omega_{\hat{D}_R, V_B}$ there exists an element $(\psi_{\bar{d}^i})_{i \in I}$ in $\bar{A}_{G_\chi}(S^2, V_B)$ which is determined by $\Phi_{\hat{D}_R}$. In fact, it can be checked that this element is unique by the proof of Theorem 8.8.

Corollary 8.11. *The set* $\Omega_{\hat{D}_R, V_B}$ *is equal to the set*

$$\Big\{ \Phi_{\hat{D}_R} \in C^0(\hat{D}_R, V_B) \;\Big|\; \Phi_{\hat{D}_R}(\hat{x}) \in \mathcal{A}^0_{\bar{e}^0} \text{ for each } \hat{x} \in [\hat{d}^0_+, \hat{d}^1_-], \text{ and}$$
$$\Phi_{\hat{D}_R}(\hat{d}^0_+) = \psi_{\bar{d}^0}(\bar{d}^0),\ \Phi_{\hat{D}_R}(\hat{d}^1_-) = \psi^{-1}_{\bar{d}^1}(\bar{d}^1) \text{ for some } (\psi_{\bar{d}^i})_{i \in I} \in \bar{A}_{G_\chi}(S^2, V_B) \Big\}.$$

Proof. To prove this corollary, we would rewrite Theorem 8.8 by using $\mathcal{A}_{\bar{e}^0}$ and $\bar{A}_{G_\chi}(S^2, V_B)$. By Theorem 8.8, a preclutching map $\Phi_{\hat{D}_R}$ is in $\Omega_{\hat{D}_R, V_B}$ if and only if a set $\Psi = (\psi_{\bar{x}})_{\bar{x} \in \bar{D}_R}$ is determined by $\Phi_{\hat{D}_R}$. As we have seen in the proof of the theorem, $gx = x'$ with $x \neq x'$ in Condition E3. is possible only when x and x' are d^i's (of course, more precisely when they are vertices). So, Ψ is determined by $\Phi_{\hat{D}_R}$ if and only if $(\psi_{\bar{d}^i})_{i \in I}$ satisfies Condition E2. and E3. and $(\psi_{\bar{x}})_{\bar{x} \in \bar{D}_R - \{\bar{d}^i | i \in I\}}$ satisfies Condition E2. Here, $(\psi_{\bar{d}^i})_{i \in I}$ satisfies Condition E2. if and only if

$$(*) \qquad \Phi_{\hat{D}_R}(\hat{d}^0_+) = \psi_{\bar{d}^0}(\bar{d}^0) \quad \text{and} \quad \Phi_{\hat{D}_R}(\hat{d}^1_-) = \psi^{-1}_{\bar{d}^1}(\bar{d}^1).$$

Lemma 8.4.(3) says that (*) implies $\Phi_{\hat{D}_R}(\hat{x}) \in \mathcal{A}^0_{\bar{e}^i}$ for $\hat{x} = \hat{d}^0_+, \hat{d}^1_-$. So, (*) could be redundantly rewritten as

$$(**) \qquad \Phi_{\hat{D}_R}(\hat{d}^0_+) = \psi_{\bar{d}^0}(\bar{d}^0), \quad \Phi_{\hat{D}_R}(\hat{d}^1_-) = \psi^{-1}_{\bar{d}^1}(\bar{d}^1), \quad \text{and } \Phi_{\hat{D}_R}(\hat{x}) \in \mathcal{A}^0_{\bar{e}^0}$$

for $\hat{x} = \hat{d}^0_+, \hat{d}^1_-$. And, $(\psi_{\bar{d}^i})_{i \in I}$ satisfies Condition E3. if and only if

$$(\psi_{\bar{d}^i})_{i \in I} \in \bar{A}_{G_\chi}(S^2, V_B). \tag{***}$$

Next, we deal with $\psi_{\bar{x}}$'s in $(\psi_{\bar{x}})_{\bar{x} \in \bar{D}_R - \{\bar{d}^i | i \in I\}}$. They satisfy Condition E2. if and only if $\psi_{\bar{x}}(\bar{x}) = \Phi_{\hat{D}_R}(\hat{x}_+)$ for each $\bar{x}$ in $\bar{x} \in \bar{D}_R - \{\bar{d}^i | i \in I\}$. And, this is satisfied if and only if $\Phi_{\hat{D}_R}(\hat{x}_+) \in \mathcal{A}^0_{\bar{x}} = \mathcal{A}^0_{\bar{e}^0}$ and we have chosen $\psi_{\bar{x}}$'s such that $\psi_{\bar{x}}(\bar{x}) = \Phi_{\hat{D}_R}(\hat{x}_+)$ for each $\bar{x}$. In summary, three conditions of this, (**), (***) are equivalent conditions for Ψ to be determined by $\Phi_{\hat{D}_R}$. Therefore, we obtain a proof. □ □

By using this corollary, we would show nonemptiness of $\Omega_{\hat{D}_R, (W_{d^i})_{i \in I^+}}$. For this, we need a lemma.

Lemma 8.12. *For each* $(W_{d^i})_{i \in I^+} \in A_{G_\chi}(S^2, \chi)$, *if we put*

$$V_B = G_\chi \times_{(G_\chi)_{d^{-1}}} W_{d^{-1}}, \quad F_{V_B} = G_\chi \times_{(G_\chi)_{d^{-1}}} (|\bar{f}^{-1}| \times W_{d^{-1}}),$$

then each $\mathcal{A}_{\bar{x}}$ *for* $\bar{x} \in [\bar{d}^0, \bar{d}^1]$ *is nonempty. And, we can pick an element* $(\psi_{\bar{d}^i})_{i \in I}$ *in* $\bar{A}_{G_\chi}(S^2, V_B)$ *which determines* $(W_{d^i})_{i \in I^+}$.

Proof. For each $i \in I$, put $\bar{\mathbf{x}} = |\pi|^{-1}(d^i) = \{\bar{x}_j | \bar{x}_j = \bar{d}^i_j \text{ for } j \in \mathbb{Z}_m\}$ for $m = j_R$ or 2. If we put $F_i = \big(\operatorname{res}^{G_\chi}_{(G_\chi)_{d^i}} F_{V_B}\big)|_{\bar{\mathbf{x}}}$ and $N_2 = (G_\chi)_{d^i}$, then $N_1 = (G_\chi)_{\bar{d}^i}$. Example 6.1 says that F_i satisfies Condition F1. or F2. Definitions of F_i and F_{V_B} say that

$$(F_i)_{\bar{x}_0} \cong \operatorname{res}^{(G_\chi)_{d^{-1}}}_{(G_\chi)_{d^{-1}} \cap (G_\chi)_{d^i}} W_{d^{-1}}$$

because $(G_\chi)_{\bar{x}_0} = (G_\chi)_{[\bar{d}^{-1}, \bar{d}^i]} = (G_\chi)_{[d^{-1}, d^i]}$ is equal to $(G_\chi)_{d^{-1}} \cap (G_\chi)_{d^i}$ by Lemma 3.10. This implies

$$(F_i)_{\bar{x}_0} \cong \operatorname{res}^{(G_\chi)_{d^i}}_{(G_\chi)_{d^{-1}} \cap (G_\chi)_{d^i}} W_{d^i}$$

by Definition 1.1.(4), i.e. W_{d^i} is a $(G_\chi)_{d^i}$-extension of $(F_i)_{\bar{x}_0}$. So, Theorem 6.6 says that $\mathcal{A}_{\bar{d}^i}$ is nonempty. Moreover, we obtain nonemptiness of $\mathcal{A}_{\bar{x}}$ for $\bar{x} \in [\bar{d}^0, \bar{d}^1]$ by Lemma 8.4. To prove the second statement, pick an element $\psi_{\bar{d}^i}$ in $\mathcal{A}_{\bar{d}^i}$ which determines W_{d^i}. If $d^1 = g \cdot d^0$ for some $g \in G_\chi$, then the element $g \cdot \bar{\psi}_{d^0}$ in $\bar{\mathcal{A}}_{\bar{d}^1}$ satisfies

$$F_1 / g \cdot \bar{\psi}_{d^0} \cong {}^g W_{d^0} \cong W_{d^1}$$

by Lemma 8.3 and Definition 1.1.(2). So, we may assume that $\bar{\psi}_{d^1} = g \cdot \bar{\psi}_{d^0}$. And, $W_{d^{-1}} \cong V_{\bar{f}^{-1}}$ by definition of V_B. Then, $(\psi_{\bar{d}^i})_{i \in I}$ is in $\bar{A}_{G_\chi}(S^2, V_B)$ which determines $(W_{d^i})_{i \in I^+}$. Therefore, we obtain a proof. □ □

Proposition 8.13. *For each* $(W_{d^i})_{i \in I^+} \in A_{G_\chi}(S^2, \chi)$, *the set* $\Omega_{\hat{D}_R, (W_{d^i})_{i \in I^+}}$ *is nonempty.*

Proof. Put $V_B = G_\chi \times_{(G_\chi)_{d^{-1}}} W_{d^{-1}}$ so that $\Omega_{\hat{D}_R, (W_{d^i})_{i \in I^+}}$ is contained in $\Omega_{\hat{D}_R, V_B}$. First, we would describe $\Omega_{\hat{D}_R, (W_{d^i})_{i \in I^+}}$ by using Corollary 8.11. By Lemma 8.12, we can pick an element $(\psi_{\bar{d}^i})_{i \in I}$ in $\bar{A}_{G_\chi}(S^2, V_B)$ which determines $(W_{d^i})_{i \in I^+}$. By Theorem 8.8 and Definition 8.10, each element $\Phi_{\hat{D}_R}$ in $\Omega_{\hat{D}_R, (W_{d^i})_{i \in I^+}}$ satisfies

$$\Phi_{\hat{D}_R}(\hat{d}^0_+) = \psi'_{\bar{d}^0}(\bar{d}^0), \ \Phi_{\hat{D}_R}(\hat{d}^1_-) = \psi'^{-1}_{\bar{d}^1}(\bar{d}^1)$$

for some $(\psi'_{\bar{d}^i})_{i \in I} \in \bar{A}_{G_\chi}(S^2, V_B)$, i.e. $(\psi'_{\bar{d}^i})_{i \in I}$ is determined by $\Phi_{\hat{D}_R}$. Easily, $\psi'_{\bar{d}^i} \in (\mathcal{A}_{\bar{d}^i})_{\psi_{\bar{d}^i}}$ for $i \in I$ because $(\psi'_{\bar{d}^i})_{i \in I}$ should determine $(W_{d^i})_{i \in I^+}$. Moreover, Lemma 8.4 says that both $\psi'_{\bar{d}^0}(\bar{d}^0)$ and $\psi'^{-1}_{\bar{d}^1}(\bar{d}^1)$ are in $(\mathcal{A}_{\bar{e}^0})^0$ which determine $\operatorname{res}^{(G_\chi)_{d^0}}_{(G_\chi)_{|e^0|}} W_{d^0}$ and $\operatorname{res}^{(G_\chi)_{d^1}}_{(G_\chi)_{|e^0|}} W_{d^1}$, respectively. Since $(G_\chi)_{|e^0|} = (G_\chi)_{d^0} \cap (G_\chi)_{d^1}$ by Lemma 3.10 and these two representations are isomorphic by definition of $A_{G_\chi}(S^2, \chi)$, Theorem 6.6 says that $\psi'_{\bar{d}^0}(\bar{d}^0)$ and $\psi'^{-1}_{\bar{d}^1}(\bar{d}^1)$ (of course, also $\psi_{\bar{d}^0}(\bar{d}^0)$ and $\psi^{-1}_{\bar{d}^1}(\bar{d}^1)$) are in the same component $(\mathcal{A}_{\bar{e}^0})^0_{\psi_{\bar{e}^0}}$ of $(\mathcal{A}_{\bar{e}^0})^0$ for some $\psi_{\bar{e}^0} \in \mathcal{A}_{\bar{e}^0}$.

Since $\psi_{\bar d^0}(\bar d^0)$ and $\psi_{\bar d^1}(\bar d^1)$ do exist, such a $\psi_{\bar e^0}$ exists and $(\mathcal{A}_{\bar e^0})_{\psi_{\bar e^0}}$ is nonempty. So, $\Omega_{\hat D_R,(W_{d^i})_{i\in I^+}}$ is expressed as

(8.1)

$$\Big\{\ \Phi_{\hat D_R} \in C^0(\hat D_R, V_B)\ \Big|\ \Phi_{\hat D_R}(\hat x) \in (\mathcal{A}_{\bar e^0})^0_{\psi_{\bar e^0}} \text{ for each } \hat x \in [\hat d^0_+, \hat d^1_-], \text{ and}$$
$$\Phi_{\hat D_R}(\hat d^0_+) = \psi'_{\bar d^0}(\bar d^0),\ \Phi_{\hat D_R}(\hat d^1_-) = \psi'^{-1}_{\bar d^1}(\bar d^1) \text{ for some } (\psi'_{\bar d^i})_{i\in I} \in \bar A_{G_\chi}(S^2, V_B)$$
$$\text{such that } \psi'_{\bar d^i} \in (\mathcal{A}_{\bar d^i})_{\psi_{\bar d^i}} \text{ for } i \in I\ \Big\}.$$

However, we can construct a continuous function $\Phi'_{\hat D_R} : [\hat d^0_+, \hat d^1_-] \to (\mathcal{A}_{\bar e^0})^0_{\psi_{\bar e^0}} \subset \mathrm{Iso}(V_{\bar f^{-1}}, V_{\bar f^0})$ such that $\Phi'_{\hat D_R}(\hat d^0_+) = \psi_{\bar d^0}(\bar d^0)$ and $\Phi'_{\hat D_R}(\hat d^1_-) = \psi^{-1}_{\bar d^1}(\bar d^1)$ because $\psi_{\bar d^0}(\bar d^0)$ and $\psi^{-1}_{\bar d^1}(\bar d^1)$ are in the nonempty path connected $(\mathcal{A}_{\bar e^0})^0_{\psi_{\bar e^0}}$. Since $\Phi'_{\hat D_R}$ is contained in the set (8.1), we obtain a proof. □ □

9. Proof for cases when $\mathrm{pr}(\rho(G_\chi)) = \mathrm{T}, \mathrm{O}, \mathrm{I}$

Now, we are ready to calculate homotopy of equivariant clutching maps.

Proposition 9.1. *Assume that* $\mathrm{pr}(R) = \mathrm{T}$, O, I *and* $R \neq \mathrm{T}$, O, I. *Then, Theorem C holds for these R's.*

Proof. We prove the proposition only for the case of $\rho(G_\chi) = \mathrm{T} \times Z$. Proof for other cases are similar. We would show that $\pi_0(\Omega_{\hat D_R,(W_{d^i})_{i\in I^+}})$ is one point set for each $(W_{d^i})_{i\in I^+} \in A_{G_\chi}(S^2, \chi)$ by Proposition 5.3.(2). Put

$$V_B = G_\chi \times_{(G_\chi)_{d^{-1}}} W_{d^{-1}}, \quad F_{V_B} = G_\chi \times_{(G_\chi)_{d^{-1}}} (|\bar f^{-1}| \times W_{d^{-1}})$$

for each $(W_{d^i})_{i\in I^+} \in A_{G_\chi}(S^2, \chi)$.

By Proposition 8.13, we know that $\pi_0(\Omega_{\hat D_R,(W_{d^i})_{i\in I^+}})$ is nontrivial. For two arbitrary $\Phi_{\hat D_R}$ and $\Phi'_{\hat D_R}$ in $\Omega_{\hat D_R,(W_{d^i})_{i\in I^+}}$, let $(\psi_{\bar d^i})_{i\in I}$ and $(\psi'_{\bar d^i})_{i\in I}$ in $\bar A_{G_\chi}(S^2, V_B)$ be two elements determined by $\Phi_{\hat D_R}$ and $\Phi'_{\hat D_R}$, respectively. We would construct a homotopy connecting $\Phi_{\hat D_R}$ and $\Phi'_{\hat D_R}$ in $\Omega_{\hat D_R,(W_{d^i})_{i\in I^+}}$. First, we show that we may assume that $(\psi'_{\bar d^i})_{i\in I} = (\psi_{\bar d^i})_{i\in I}$. Since $\psi_{\bar d^i}$ and $\psi'_{\bar d^i}$ for $i \in I$ determine the same representation W_{d^i}, these two are in the same path component of $\mathcal{A}_{\bar d^i}$ by Theorem 6.6. Take paths $\gamma^i : [0,1] \to \mathcal{A}_{\bar d^i}$ for $i \in I$ such that $\gamma^i(0) = \psi'_{\bar d^i}$ and $\gamma^i(1) = \psi_{\bar d^i}$. In the case of $R = \mathrm{T} \times Z$, we have $d^1 = t_0 \cdot d^0$, and $t_0 \cdot \gamma^0$ satisfies $(t_0 \cdot \gamma^0)(0) = \psi'_{\bar d^1}$ and $(t_0 \cdot \gamma^0)(1) = \psi_{\bar d^1}$ by Definition 8.10. So, we may also assume that $\gamma^1 = t_0 \cdot \gamma^0$. Recall that the parametrization on $|\hat e^0| = [\hat v^0_+, \hat v^1_-]$ by $s \in [0,1]$ satisfies $\hat v^0_+ = 0$, $b(\hat e^0) = 1/2$, $\hat v^1_- = 1$. We construct a homotopy $L(s,t) : [\hat d^i_+, \hat d^{i+1}_-] \times [0,1] \to \mathcal{A}^0_{\bar e^i}$ as

$$\begin{aligned} L(s,t) &= \gamma^i\Big((1-3s)t\Big)(\bar d^i) && \text{for } s \in [0, \tfrac{1}{3}],\\ L(s,t) &= \varphi'^i(3s-1) && \text{for } s \in [\tfrac{1}{3}, \tfrac{2}{3}],\\ L(s,t) &= \gamma^{i+1}\Big((3s-2)t\Big)^{-1}(\bar d^{i+1}) && \text{for } s \in [\tfrac{2}{3}, 1]. \end{aligned}$$

Then, L_t for each $t \in [0,1]$ is in $\Omega_{\hat D_R,(W_{d^i})_{i\in I^+}}$ by Corollary 8.11, and L connects $\Phi'_{\hat D_R}$ with L_1 in $\Omega_{\hat D_R,(W_{d^i})_{i\in I^+}}$ which determines $(\psi_{\bar d^i})_{i\in I}$. So, if we put $\Phi'_{\hat D_R} = L_1$, then we may assume that $\Phi_{\hat D_R}$ and $\Phi'_{\hat D_R}$ determine the same element $(\psi_{\bar d^i})_{i\in I}$ in $\bar A_{G_\chi}(S^2, V_B)$.

Now, we construct a homotopy between $\Phi_{\hat D_R}$ and $\Phi'_{\hat D_R}$. As in the proof of Proposition 8.13, $\psi_{\bar d^0}(\bar d^0)$ and $\psi^{-1}_{\bar d^1}(\bar d^1)$ are in the same component $(\mathcal{A}_{\bar e^0})^0_{\psi_{\bar e^0}}$ for some element $\psi_{\bar e^0}$ in $\mathcal{A}_{\bar e^0}$, and $\Phi_{\hat D_R}$, $\Phi'_{\hat D_R}$ have values in $(\mathcal{A}_{\bar e^0})^0_{\psi_{\bar e^0}}$. Here, note that $(\mathcal{A}_{\bar e^0})_{\psi_{\bar e^0}}$ is simply connected by Lemma 3.8 and Proposition 6.8.(3). By simply connectedness, we can obtain a homotopy $L'(s,t) : [\hat{d}^0_+, \hat{d}^1_-] \times [0,1] \to \mathcal{A}^0_{\bar{e}^0}$ as

$$\begin{array}{lll} L'(s,t) \in (\mathcal{A}_{\bar{e}^0})^0_{\psi_{\bar{e}^0}}, & L'(0,t) = \psi_{\bar{d}^0}(\bar{d}^0), & L'(1,t) = \psi^{-1}_{\bar{d}^1}(\bar{d}^1), \\ L'(s,0) = \Phi_{\hat{D}_R}(s), & L'(s,1) = \Phi'_{\hat{D}_R}(s) & \end{array}$$

for $s,t \in [0,1]$. Then, L' connects $\Phi_{\hat{D}_R}$ and $\Phi'_{\hat{D}_R}$ in $\Omega_{\hat{D}_R,(W_{d^i})_{i\in I^+}}$ by Corollary 8.11. Therefore, we obtain a proof. Here, we remark that simply connectedness is critical in obtaining L'. ☐ ☐

Proposition 9.2. *Assume that R is equal to one of* T, O, I. *Then, Theorem A holds for these cases.*

Proof. When $\rho(G_\chi)$ is given, put

$$V_B = G_\chi \times_{(G_\chi)_{d^{-1}}} W_{d^{-1}}, \quad F_{V_B} = G_\chi \times_{(G_\chi)_{d^{-1}}} (|\bar{f}^{-1}| \times W_{d^{-1}})$$

for each $(W_{d^i})_{i\in I^+} \in A_{G_\chi}(S^2,\chi)$. We can pick an element $(\psi_{\bar{d}^i})_{i\in I}$ in $\bar{A}_{G_\chi}(S^2, V_B)$ which determines $(W_{d^i})_{i\in I^+}$ by Lemma 8.12. In these cases, we have

$$\begin{array}{ll} \bar{D}_R = [\bar{v}^0, b(\bar{e}^0)], & (G_\chi)_{v^0}/H \cong \mathbb{Z}_{j_R}, \\ (G_\chi)_{b(e^0)}/H \cong \mathbb{Z}_2, & (G_\chi)_x/H = \langle \mathrm{id} \rangle \end{array}$$

by Table 3.4. By using these, path components of $\mathcal{A}_{\bar{v}^0}$, $\mathcal{A}_{b(\bar{e}^0)}$ are simply connected by Proposition 6.8.(3), and we have $\mathcal{A}^0_{\bar{e}^0} = \mathrm{Iso}_H(V_{\bar{f}^{-1}}, V_{\bar{f}^0})$ by Lemma 6.4. Here, $\mathcal{A}_{\bar{v}^0}$, $\mathcal{A}_{b(\bar{e}^0)}$, $\mathcal{A}_{\bar{e}^0}$ are all nonempty by Lemma 8.12. Since $d^1 \neq gd^0$ for any $g \in G_\chi$, $\psi'_{\bar{d}^0}$ and $\psi'_{\bar{d}^1}$ in the set (8.1) have no relation. So, the set (8.1) is rewritten as

$$\Big\{ \Phi_{\hat{D}_R} \in C^0(\hat{D}_R, V_B) \;\Big|\; \Phi_{\hat{D}_R}(\hat{x}) \in \mathrm{Iso}_H(V_{\bar{f}^{-1}}, V_{\bar{f}^0}) \text{ for each } \hat{x} \in [\hat{v}^0_+, b(\hat{e}^0)],$$
$$\text{and } \Phi_{\hat{D}_R}(\hat{v}^0_+) \in (\mathcal{A}_{\bar{v}^0})^0_{\psi_{\bar{v}^0}},\ \Phi_{\hat{D}_R}\big(b(\hat{e}^0)\big) \in (\mathcal{A}_{b(\bar{e}^0)})^0_{\psi_{b(\bar{e}^0)}} \Big\}.$$

Since path components $(\mathcal{A}_{\bar{v}^0})^0_{\psi_{\bar{v}^0}}$ and $(\mathcal{A}_{b(\bar{e}^0)})^0_{\psi_{b(\bar{e}^0)}}$ are simply connected, the homotopy $\pi_0(\Omega_{\hat{D}_R,(W_{d^i})_{i\in I^+}})$ is in one-to-one correspondence with the homotopy $\pi_1\Big(\mathrm{Iso}_H(V_{\bar{f}^{-1}}, V_{\bar{f}^0})\Big)$ by Lemma 7.4, and the injection $\imath_0$ from Ω_0 to $\Omega_{\hat{D}_R,(W_{d^i})_{i\in I^+}}$ induces the bijection

$$\pi_0(\imath_0) : \pi_0(\Omega_0) \to \pi_0(\Omega_{\hat{D}_R,(W_{d^i})_{i\in I^+}})$$

where Ω_0 is defined as

$$\Big\{ \Phi_{\hat{D}_R} \in C^0(\hat{D}_R, V_B) \;\Big|\; \Phi_{\hat{D}_R}(\hat{x}) \in \mathrm{Iso}_H(V_{\bar{f}^{-1}}, V_{\bar{f}^0}) \text{ for each } \hat{x} \in [\hat{v}^0_+, b(\hat{e}^0)],$$
$$\text{and } \Phi_{\hat{D}_R}(\hat{v}^0_+) = \psi_{\bar{v}^0}(\bar{v}^0),\ \Phi_{\hat{D}_R}\big(b(\hat{e}^0)\big) = \psi_{b(\bar{e}^0)}(b(\bar{e}^0)) \Big\}.$$

Now, we would show that Chern classes of equivariant vector bundles determined by different classes in $\pi_0\big(\Omega_{\hat{D}_R,(W_{d^i})_{i\in I^+}}\big)$ with respect to V_B are all different by calculating Chern class. For this, we use the parametrization on $\bar{\mathcal{L}}_R$ introduced in Section 4. For notational simplicity, put $* = \psi_{\bar{v}^0}(\bar{v}^0)$ and $*' = \psi_{b(\bar{e}^0)}(b(\bar{e}^0))$. Let $\gamma_0 : [0,1/2] \to \mathrm{Iso}_H(V_{\bar{f}^{-1}}, V_{\bar{f}^0})$ be an element of the set Ω_0 such that $\gamma_0(0) = *$ and $\gamma_0(1/2) = *'$. Also, let $\sigma_0 : [0,1/2] \to \mathrm{Iso}_H(V_{\bar{f}^{-1}}, V_{\bar{f}^0})$ be a loop such that $\sigma_0(0) = \sigma_0(1/2) = *$ and $[\sigma_0]$ is a generator of $\pi_1(\mathrm{Iso}_H(V_{\bar{f}^{-1}}, V_{\bar{f}^0}), *)$. Here, we introduce some notations.

Notation 9.3. For some H-representations W_1 and W_2, let X_1 and X_2 be two spaces $\mathrm{Iso}_H(W_1, W_2)$ and $\mathrm{Iso}_H(W_2, W_1)$, respectively. For paths $\delta_1, \delta_2 : [a,b] \to X_1$ satisfying $\delta_1(b) = \delta_2(a)$, denote by $\delta_1.\delta_2 : [a,b] \to X_1$ the path defined by

$$(\delta_1.\delta_2)(t) = \begin{cases} \delta_1(a + 2(t-a)), & t \in [a, \frac{a+b}{2}], \\ \delta_2(a + 2(t - \frac{a+b}{2})), & t \in [\frac{a+b}{2}, b]. \end{cases}$$

For paths $\delta_3 : [a,b] \to X_1$ and $\delta_4 : [b,c] \to X_1$ satisfying $\delta_3(b) = \delta_4(b)$, denote by $\delta_1 \vee \delta_2 : [a,c] \to X_1$ the path defined by

$$(\delta_3 \vee \delta_4)(t) = \begin{cases} \delta_3(t), & t \in [a,b], \\ \delta_4(t), & t \in [b,c]. \end{cases}$$

Also, for a path $\delta : [a,b] \to X_1$ denote by $\delta^* : [1-b, 1-a] \to X_2$ the path $\delta^*(t) = \delta(1-t)^{-1}$.

Note that $\sigma_0.\gamma_0 \in \Omega_0 \subset \Omega_{\hat{D}_R,(W_{d^i})_{i\in I+}}$. We would show that the difference between $c_1\big(F_{V_B}/\sigma_0.\gamma_0\big)$ and $c_1\big(F_{V_B}/\gamma_0\big)$ is $12\chi(\mathrm{id})$, $24\chi(\mathrm{id})$, $60\chi(\mathrm{id})$ according to $R =$ T, O, I, respectively. Here, $l_R = 12$, 24, 60 because the number of edges of $\mathcal{K}_{\mathrm{T}}$, $\mathcal{K}_{\mathrm{O}}$, $\mathcal{K}_{\mathrm{I}}$ is 6, 12, 30 and D_R is a half of an edge at each case, respectively. For the calculation, we need to describe the equivariant clutching maps precisely. Let $\Phi = \cup\, \varphi_{\hat{e}}$ in $\Omega_{\hat{D}_R,(W_{d^i})_{i\in I+}}$ be the extension of γ_0, i.e. $\varphi_{\hat{e}^0}(t) = \gamma_0(t)$ for $t \in [0, 1/2]$. By Condition N1., $\varphi_{c(\hat{e}^0)}(t) = \gamma_0(1-t)^{-1}$ for $t \in [1/2, 1]$. Pick an element c_0 of $(G_\chi)_{b(e^0)}$ such that $c_0 b(\bar{e}^0) = b\big(c(\bar{e}^0)\big)$. By equivariance of Φ,

$$\begin{aligned} \varphi_{c(\hat{e}^0)}(t) &= \Phi(t) \\ &= c_0\Phi(c_0^{-1}t)c_0^{-1} \\ &= c_0\varphi_{\hat{e}^0}(c_0^{-1}t)c_0^{-1} \\ &= c_0\gamma_0(t)c_0^{-1} \end{aligned}$$

for $t \in [0, 1/2]$. Again by Condition N1.,

$$\begin{aligned} \varphi_{\hat{e}^0}(t) &= \varphi_{c(\hat{e}^0)}(1-t)^{-1} \\ &= c_0\gamma_0(1-t)^{-1}c_0^{-1} \end{aligned}$$

for $t \in [1/2, 1]$. This gives $\varphi_{\hat{e}^0} = \gamma_0 \vee c_0\gamma_0^* c_0^{-1}$ and $\varphi_{c(\hat{e}^0)} = c_0\gamma_0 c_0^{-1} \vee \gamma_0^*$. Let $\Phi' = \cup\varphi'_{\hat{e}}$ be the extension of $\sigma_0.\gamma_0$. Then,

$$\begin{aligned} \varphi'_{\hat{e}^0} &= \sigma_0.\gamma_0 \vee c_0(\gamma_0^*.\sigma_0^*)c_0^{-1} \text{ and} \\ \varphi'_{c(\hat{e}^0)} &= c_0\sigma_0.\gamma_0 c_0^{-1} \vee \gamma_0^*.\sigma_0^* \end{aligned}$$

by using $(\sigma_0.\gamma_0)^* = \gamma_0^*.\sigma_0^*$. Replacing only $\varphi_{\hat{e}}$ and $\varphi_{c(\hat{e}^0)}$ of Φ with $\varphi'_{\hat{e}}$ and $\varphi'_{c(\hat{e}^0)}$, we obtain a new nonequivariant clutching map, say Φ_1. Here, Φ_1 becomes an nonequivariant clutching map because $\Phi_1 = \Phi$ on vertices of $\bar{\mathcal{L}}_R$. The difference $c_1\big(F_{V_B}/\Phi_1\big) - c_1\big(F_{V_B}/\Phi\big)$ is equal to $\pm 2\chi(\mathrm{id})$ (say $+2\chi(\mathrm{id})$) by Lemma 7.1 because $\varphi'_{\hat{e}^0}$ contains two generators σ_0 and $c_0\sigma_0^* c_0^{-1}$. We would repeat this argument for other edges. Pick edges $\hat{e}$ and $c(\hat{e})$ such that $\{\hat{e}, c(\hat{e})\} \neq \{\hat{e}^0, c(\hat{e}^0)\}$. Then, there exists the unique $g_0 \in G_\chi$ up to H such that $g_0\hat{e}^0 = \hat{e}$. By equivariance,

$$\begin{aligned} \varphi_{\hat{e}} &= g_0 \cdot \varphi_{\hat{e}^0} \\ &= g_0 \cdot (\gamma_0 \vee c_0\gamma_0^* c_0^{-1}) \\ &= g_0\gamma_0 g_0^{-1} \vee g_0 c_0\gamma_0^* c_0^{-1} g_0^{-1}, \end{aligned}$$

and

$$\varphi'_{\hat{e}} = g_0\sigma_0.\gamma_0 g_0^{-1} \vee g_0 c_0 \gamma_0^*.\sigma_0^* c_0^{-1} g_0^{-1}.$$

Replacing only $\varphi_{\hat{e}}$ and $\varphi_{c(\hat{e}^0)}$ of Φ_1 with $\varphi'_{\hat{e}}$ and $\varphi'_{c(\hat{e}^0)}$, we obtain a new nonequivariant clutching map, say Φ_2, is obtained, and the difference $c_1\big(F_{V_B}/\Phi_2\big) - c_1\big(F_{V_B}/\Phi_1\big)$ is also equal to $2\chi(\mathrm{id})$ because g_0 preserves the orientation. In this way, if we replace all $\varphi_{\hat{e}}$ of Φ with $\varphi'_{\hat{e}}$ to obtain Φ', then the difference $c_1\big(F_{V_B}/\Phi'\big) - c_1\big(F_{V_B}/\Phi\big)$ is equal to $l_R\chi(\mathrm{id})$. Proposition 5.3 and these calculations prove the proposition. □ □

10. EQUIVARIANT CLUTCHING MAPS WHEN $\mathrm{pr}(\rho(G_\chi)) = \mathbb{Z}_n$, D_n

Assume that $\rho(G_\chi) = R$ for some finite R in Table 1.1 such that $\mathrm{pr}(R) = \mathbb{Z}_n$, D_n. We redefine notations introduced for cases of $\mathrm{pr}(R) = \mathrm{T}$, O, I to be in accordance with these cases, and mimic what we have done in Section 4, 5, 8 with those notations. In our treatment of cases when $R = \mathbb{Z}_n, \langle a_n, -b \rangle$, there are some differences with other cases which are caused by existence of fixed points.



FIGURE 10.1. $\pi : \bar{\mathcal{K}}_R \to \mathcal{K}_R$ when $\mathcal{K}_R = \mathcal{K}_4$

First, we rewrite Section 4. In these cases, $\mathcal{K}_R = \mathcal{K}_{m_R}$ with $m_R = n/2$, n, $2n$ by Table 1.1. Denote by $\mathcal{K}_{R,S}$ the lower simplicial subcomplex of $\mathcal{K}_R$ whose underlying space $|\mathcal{K}_{R,S}|$ is equal to $|\mathcal{K}_R| \cap \{(x,y,z) \in \mathbb{R}^3 \mid z \le 0\}$, and by $\mathcal{K}_{R,N}$ the upper part. Let $\mathcal{L}_R$ be the subcomplex $\mathcal{K}_{R,S} \cap \mathcal{K}_{R,N}$ of $\mathcal{K}_R$ lying on the equator $z = 0$. Put $B = \{S, N\} \subset |\mathcal{K}_R|$ on which R (and G_χ) acts. Since each $|\mathcal{K}_{R,q}|$ for $q \in B$ has a simple equivariant structure for R_q, we consider $\mathcal{K}_R$ as the union of two pieces $\mathcal{K}_{R,q}$ for $q \in B$, not of all faces of $\mathcal{K}_R$. Denote by $\bar{\mathcal{K}}_R$ the disjoint union $\amalg_{q \in B} \mathcal{K}_{R,q}$, and denote by $\bar{\mathcal{K}}_{R,q}$ the subcomplex $\mathcal{K}_{R,q}$ of $\bar{\mathcal{K}}_R$ so that $|\bar{\mathcal{K}}_R| = \amalg_{q \in B} |\bar{\mathcal{K}}_{R,q}|$. We denote simply by π and $|\pi|$ natural quotient maps from $\bar{\mathcal{K}}_R$ and $|\bar{\mathcal{K}}_R|$ to $\mathcal{K}_R$ and $|\mathcal{K}_R|$, respectively. Let $\bar{\mathcal{L}}_R = \pi^{-1}(\mathcal{L}_R)$, and let $\bar{\mathcal{L}}_{R,q}$ be the subcomplex $\bar{\mathcal{L}}_R \cap \bar{\mathcal{K}}_{R,q}$ of $\bar{\mathcal{K}}_R$ for $q \in B$. The subset $|\pi|^{-1}(B)$ in $|\bar{\mathcal{K}}_R|$ is often confused with B in $|\mathcal{K}_R|$. The injection from $\bar{\mathcal{L}}_R$ to $\bar{\mathcal{K}}_R$ and its underlying space map are denoted by $\imath_{\bar{\mathcal{L}}}$ and $\imath_{|\bar{\mathcal{L}}|}$, respectively. The G_χ-actions on $\mathcal{K}_R$, $|\mathcal{K}_R|$ induce G_χ-actions on $\mathcal{L}_R$, $\bar{\mathcal{K}}_R$, $\bar{\mathcal{L}}_R$, and their underlying spaces. We do not need define $\hat{\mathcal{L}}_R$. Here, we introduce the following notations:

$$\begin{aligned}
\bar{v}_q^i &= \bar{\mathcal{L}}_{R,q} \cap \pi^{-1}(v^i), \\
\bar{e}_q^i &= \bar{\mathcal{L}}_{R,q} \cap \pi^{-1}(e^i), \\
\bar{D}_R &= |\bar{\mathcal{L}}_{R,S}| \cap |\pi|^{-1}(D_R), \\
\bar{\mathbf{D}}_R &= |\pi|^{-1}(D_R), \\
\bar{d}^{i'} &= |\bar{\mathcal{L}}_{R,S}| \cap |\pi|^{-1}(d^{i'})
\end{aligned}$$

for $q \in B$, $i \in \mathbb{Z}_{m_R}$, $i' \in I$. Some notations are illustrated in Figure 10.1. Also, let $c : \bar{\mathcal{L}}_R \to \bar{\mathcal{L}}_R$ be the simplicial map whose underlying space map $|c| : |\bar{\mathcal{L}}_R| \to |\bar{\mathcal{L}}_R|$ satisfies

$$\bar{x} \neq |c|(\bar{x}) \quad \text{and} \quad |\pi|(\bar{x}) = |\pi|\big(|c|(\bar{x})\big)$$

for each $\bar{x} \in |\bar{\mathcal{L}}_R|$. And, put $\bar{x}_0 = \bar{x}$ and $\bar{x}_1 = |c|(\bar{x})$ for each $\bar{x} \in |\bar{\mathcal{L}}_R|$. Now, we describe an equivariant vector bundle over $|\mathcal{K}_R|$ as equivariant clutching construction by using an equivariant vector bundle over $|\bar{\mathcal{K}}_R|$. Pick a G_χ-vector bundle V_B over B such that $(\operatorname{res}_H^{G_\chi} V_B)|_q$ is χ-isotypical at each q in B. We denote by V_q the isotropy representations of V_B at q for each $q \in B$. Then, we have

$$V_B \cong G_\chi \times_{(G_\chi)_S} V_S \quad \text{and} \quad \operatorname{res}_H^{(G_\chi)_S} V_S \cong \operatorname{res}_H^{(G_\chi)_N} V_N \tag{10.1}$$

when $R \neq \mathbb{Z}_n$, $\langle a_n, -b \rangle$. Define $\operatorname{Vect}_{G_\chi}(|\mathcal{K}_R|, \chi)_{V_B}$ and $\operatorname{Vect}_{G_\chi}(|\bar{\mathcal{K}}_R|, \chi)_{V_B}$ as before. We observe that $\operatorname{Vect}_{G_\chi}(|\bar{\mathcal{K}}_R|, \chi)_{V_B}$ has the unique element $[F_{V_B}]$ for the bundle

$$F_{V_B} = \begin{cases} \amalg_{q \in B} \, |\bar{\mathcal{K}}_{R,q}| \times V_q & \text{if } R = \mathbb{Z}_n, \ \langle a_n, -b \rangle, \\ G_\chi \times_{(G_\chi)_S} (|\bar{\mathcal{K}}_{R,S}| \times V_S) & \text{if } R \neq \mathbb{Z}_n, \ \langle a_n, -b \rangle. \end{cases}$$

Henceforward, we use trivializations

$$\begin{array}{ll} |\bar{\mathcal{K}}_{R,q}| \times V_q & \text{for } \big(\operatorname{res}_{(G_\chi)_q}^{G_\chi} F_{V_B}\big)\big|_{|\bar{\mathcal{K}}_{R,q}|}, \\ |\bar{e}_q^i| \times \operatorname{res}_{(G_\chi)_{b(\bar{e}_q^i)}}^{(G_\chi)_q} V_q & \text{for } \big(\operatorname{res}_{(G_\chi)_{b(\bar{e}_q^i)}}^{G_\chi} F_{V_B}\big)\big|_{|\bar{e}_q^i|} \end{array} \tag{10.2}$$

for each q and i. Then, each E in $\operatorname{Vect}_{G_\chi}(|\mathcal{K}_R|, \chi)_{V_B}$ can be constructed by gluing $F_{V_B} \cong |\pi|^* E$ along $|\bar{\mathcal{L}}_{R,q}|$'s through

$$|\bar{\mathcal{L}}_{R,q}| \times V_q \to |\bar{\mathcal{L}}_{R,q'}| \times V_{q'}, \ (\bar{x}, u) \mapsto (|c|(\bar{x}), \varphi_q(\bar{x})u)$$

via continuous maps

$$\varphi_q : |\bar{\mathcal{L}}_{R,q}| \to \operatorname{Iso}(V_q, V_{q'})$$

for $\bar{x} \in |\bar{\mathcal{L}}_{R,q}|$, $u \in V_q$, $B = \{q, q'\}$ as we have done in Section 4. The union $\Phi = \cup_{q \in B} \, \varphi_q$ is called an equivariant clutching map of E with respect to V_B. This construction of E is denoted by F_{V_B}/Φ. The map Φ is defined on $|\bar{\mathcal{L}}_R|$, and we also regard Φ as a map

$$\imath_{|\bar{\mathcal{L}}|}^* F_{V_B} \to \imath_{|\bar{\mathcal{L}}|}^* F_{V_B}, \quad (\bar{x}, u) \mapsto \big(|c|(\bar{x}), \Phi(\bar{x})u\big)$$

by using trivialization (10.2) for each $q \in B$, $(\bar{x}, u) \in |\bar{\mathcal{L}}_{R,q}| \times V_q$. Also, Φ should be equivariant. An equivariant clutching map of some bundle in $\operatorname{Vect}_{G_\chi}(|\mathcal{K}_R|, \chi)_{V_B}$ with respect to V_B is called simply an equivariant clutching map with respect to V_B, and let Ω_{V_B} be the set of all equivariant clutching maps with respect to V_B. And, if we define $\Omega_{\bar{D}_R, V_B}$ as the set

$$\{ \ \Phi|_{\bar{D}_R} \mid \Phi \in \Omega_{V_B} \ \},$$

then the restriction map $\Omega_{V_B} \to \Omega_{\bar{D}_R, V_B}$ is bijective, and $\pi_0(\Omega_{V_B}) \cong \pi_0(\Omega_{\bar{D}_R, V_B})$. We call an equivariant clutching map Φ the *extension* of $\Phi|_{\bar{D}_R}$. And, denote also by $F_{V_B}/\ \Phi|_{\bar{D}_R}$ the bundle F_{V_B}/Φ. Let $C^0(|\bar{\mathcal{L}}_R|, V_B)$ be the set of functions Φ defined on $|\bar{\mathcal{L}}_R|$ satisfying $\Phi|_{|\bar{\mathcal{L}}_{R,q}|}(\bar{x}) \in \operatorname{Iso}(V_q, V_{q'})$ for $\bar{x} \in |\bar{\mathcal{L}}_{R,q}|$ and $B = \{q, q'\}$. We can define the quotient F_{V_B}/Φ for any Φ in $C^0(|\bar{\mathcal{L}}_R|, V_B)$. And, let $C^0(\bar{D}_R, V_B)$ be the set

$$\{ \ \Phi|_{\bar{D}_R} \mid \Phi \in C^0(|\bar{\mathcal{L}}_R|, V_B) \ \}.$$

A function Φ in $C^0(|\bar{\mathcal{L}}_R|, V_B)$ or a function $\Phi_{\bar{D}_R}$ in $C^0(\bar{D}_R, V_B)$ is called a *preclutching map* with respect to V_B. A preclutching map Φ in $C^0(|\bar{\mathcal{L}}_R|, V_B)$ is an equivariant clutching map with respect to V_B if and only if it satisfies the following conditions:

N1′. $\Phi(|c|(\bar{x})) = \Phi(\bar{x})^{-1}$ for each $\bar{x} \in |\bar{\mathcal{L}}_R|$, E1$'$. $\Phi(g\bar{x}) = g\Phi(\bar{x})g^{-1}$ for each $\bar{x} \in |\bar{\mathcal{L}}_R|, g \in G_\chi$.

More precisely, if Φ satisfies Condition N1$'$., then the quotient F_{V_B}/Φ becomes a nonequivariant vector bundle. And, if Φ also satisfies Condition E1$'$., then F_{V_B}/Φ becomes an equivariant vector bundle, i.e. Φ is an equivariant clutching map with respect to V_B.

Second, we rewrite Section 5. We obtain Lemma 5.1 and 5.2 by replacing $p_{|\bar{\mathcal{L}}|}$ with $\imath_{|\bar{\mathcal{L}}|}$. Define $\imath_\Omega$, p_Ω, p_{π_0} as before by replacing $\Omega_{\hat{D}_R, V_B}$ with $\Omega_{\bar{D}_R, V_B}$. We decompose $\Omega_{\bar{D}_R, V_B}$. When $R = \mathbb{Z}_n$, $\langle a_n, -b\rangle$, denote a pair (W_S, W_N) in $A_{G_\chi}(S^2, \chi)$ by $(W_q)_{q\in B}$. And, pick the G_χ-vector bundle V_B over B such that $V_B|_q = W_q$ for each $(W_q)_{q\in B} \in A_{G_\chi}(S^2, \chi)$ and $q \in B$. Define $\Omega_{\bar{D}_R, (W_q)_{q\in B}} = (p_\Omega \circ p_{\pi_0})^{-1}\big((W_q)_{q\in B}\big)$. Then, we obtain $\Omega_{\bar{D}_R, V_B} = \Omega_{\bar{D}_R, (W_q)_{q\in B}}$ because V_B and $(W_q)_{q\in B}$ can be regarded as the same G_χ-bundle over B. When $R \neq \mathbb{Z}_n$, $\langle a_n, -b\rangle$, put

$$V_B = G_\chi \times_{(G_\chi)_{d^{-1}}} W_{d^{-1}}, \quad F_{V_B} = G_\chi \times_{(G_\chi)_{d^{-1}}} (|\bar{f}^{-1}| \times W_{d^{-1}})$$

for each $(W_{d^i})_{i\in I^+} \in A_{G_\chi}(S^2, \chi)$. Then, we obtain the decomposition of $\Omega_{\bar{D}_R, V_B}$ by replacing $\Omega_{\hat{D}_R, (W_{d^i})_{i\in I^+}}$ of (5.3) with $\Omega_{\bar{D}_R, (W_{d^i})_{i\in I^+}}$. Proposition 5.3 holds if

1. we replace $(W_{d^i})_{i\in I^+}$, $\Omega_{\hat{D}_R, (W_{d^i})_{i\in I^+}}$ with $(W_q)_{q\in B}$, $\Omega_{\bar{D}_R, (W_q)_{q\in B}}$ when $R = \mathbb{Z}_n$, $\langle a_n, -b\rangle$, respectively.
2. we replace $\Omega_{\hat{D}_R, (W_{d^i})_{i\in I^+}}$ with $\Omega_{\bar{D}_R, (W_{d^i})_{i\in I^+}}$ when $R \neq \mathbb{Z}_n$, $\langle a_n, -b\rangle$.

Third, we rewrite Section 8. To begin with, we define some sets of equivariant pointwise clutching maps. For each $\bar{x}$ in $|\bar{\mathcal{L}}_R|$ and $x = |\pi|(\bar{x})$, put $\bar{\mathbf{x}} = |\pi|^{-1}(x) = \{\bar{x}_j | j \in \mathbb{Z}_2\}$, and let $\mathcal{A}_{\bar{x}}$ be the set of equivariant pointwise clutching maps with respect to the $(G_\chi)_x$-bundle $\big(\operatorname{res}^{G_\chi}_{(G_\chi)_x} F_{V_B}\big)|_{\bar{\mathbf{x}}}$. Also for each edge $\bar{e}$ in $\bar{\mathcal{L}}_R$, $\pi(\bar{e}) = e$, $\bar{x}$ in $|\bar{e}|$, put $\bar{\mathbf{x}} = |\pi|^{-1}(x) = \{\bar{x}_j | j \in \mathbb{Z}_2\}$, and let $\mathcal{A}_{\bar{e}}$ be the set of equivariant pointwise clutching maps with respect to the $(G_\chi)_{|e|}$-bundle $\big(\operatorname{res}^{G_\chi}_{(G_\chi)_{|e|}} F_{V_B}\big)|_{\bar{\mathbf{x}}}$ where $\mathcal{A}_{\bar{e}}$'s for different $\bar{x}$'s are identified as in Section 8. We can define saturations on $\mathcal{A}_{\bar{x}}$ and elements of it, and also group actions on saturations. Evaluation maps $\mathcal{A}_{\bar{x}} \to \mathcal{A}^0_{\bar{x}}$ and $\mathcal{A}_{\bar{e}} \to \mathcal{A}^0_{\bar{e}}$ are bijective for any $\bar{x} \in |\bar{\mathcal{L}}_R|$, $\bar{e} \in \bar{\mathcal{L}}_R$. With these, we can rewrite Theorem 8.8.

Theorem 10.1. *Assume that* $\operatorname{pr}(R) = \mathbb{Z}_n, \mathrm{D}_n$ *. Then, a preclutching map* $\Phi_{\bar{D}_R}$ *in* $C^0(\bar{D}_R, V_B)$ *is in* $\Omega_{\bar{D}_R, V_B}$ *if and only if there exists the unique* $\psi_{\bar{x}} \in \mathcal{A}_{\bar{x}}$ *for each* $\bar{x} \in \bar{D}_R$ *satisfying the following conditions:*

E2$'$. $\psi_{\bar{x}}(\bar{x}) = \Phi_{\bar{D}_R}(\bar{x})$ *for each* $\bar{x} \in \bar{D}_R$,

E3$'$. *for each* $\bar{x}, \bar{x}' \in \bar{D}_R$ *and* $x = |\pi|(\bar{x}), x' = |\pi|(\bar{x}')$, *if* $x' = gx$ *for some* $g \in G_\chi$, *then* $\bar{\psi}_{x'} = g \cdot \bar{\psi}_x$.

The set $(\psi_{\bar{x}})_{\bar{x}\in\bar{D}_R}$ *is called determined by* $\Phi_{\bar{D}_R}$.

In cases when $R = \mathbb{Z}_n$, $\langle a_n, -b\rangle$, Theorem 10.1 is sufficient to calculate the homotopy $\pi_0(\Omega_{\bar{D}_R, V_B})$, but in other cases we need more. In the remaining of this section, assume that $R \neq \mathbb{Z}_n$, $\langle a_n, -b\rangle$. With exactly the same definition of $\bar{A}_{G_\chi}(S^2, V_B)$, we can rewrite Corollary 8.11.

Corollary 10.2. *The set* $\Omega_{\bar{D}_R, V_B}$ *is equal to the set*

$$\Big\{ \Phi_{\bar{D}_R} \in C^0(\bar{D}_R, V_B) \Big| \; \Phi_{\bar{D}_R}(\bar{x}) \in \mathcal{A}^0_{\bar{e}} \text{ for each } \bar{x} \text{ and any } \bar{e} \text{ such that } \bar{x} \in |\bar{e}|, \text{ and}$$
$$\Phi_{\bar{D}_R}(\bar{d}^0) = \psi_{\bar{d}^0}(\bar{d}^0),\ \Phi_{\bar{D}_R}(\bar{d}^1) = \psi_{\bar{d}^1}(\bar{d}^1) \text{ for some } (\psi_{\bar{d}^i})_{i\in I} \in \bar{A}_{G_\chi}(S^2, V_B) \Big\}.$$

By using this, we would show nonemptiness of $\Omega_{\bar{D}_R, (W_{d^i})_{i\in I^+}}$. For this, we need a lemma.

Lemma 10.3. *For each* $(W_{d^i})_{i\in I^+} \in A_{G_\chi}(S^2,\chi)$*, if we put*

$$V_B = G_\chi \times_{(G_\chi)_{d^{-1}}} W_{d^{-1}}, \quad F_{V_B} = G_\chi \times_{(G_\chi)_{d^{-1}}} (|\bar{\mathcal{K}}_{R,S}| \times W_{d^{-1}}),$$

then each $\mathcal{A}_{\bar{x}}$ *for* $\bar{x} \in [\bar{d}^0, \bar{d}^1]$ *is nonempty. And, we can pick an element* $(\psi_{\bar{d}^i})_{i\in I}$ *in* $\bar{A}_{G_\chi}(S^2, V_B)$ *which determines* $(W_{d^i})_{i\in I^+}$*.*

Proof. For each $i \in I$, put $\bar{\mathbf{x}} = |\pi|^{-1}(d^i) = \{\bar{x}_j | \bar{x}_j = \bar{d}^i_j \text{ for } j \in \mathbb{Z}_2\}$. If we put $F_i = \big(\operatorname{res}^{G_\chi}_{(G_\chi)_{d^i}} F_{V_B}\big)|_{\bar{\mathbf{x}}}$ and $N_2 = (G_\chi)_{d^i}$, then $N_1 = (G_\chi)_{\bar{d}^i}$. By Table 3.4 and (10.1), at least one of F_i's satisfies Condition F2. except two cases $R = \mathbb{Z}_n \times Z$ with odd n, $\langle -a_n \rangle$ with even $n/2$. Then, we obtain nonemptiness as in Lemma 8.12 in these cases. In the remaining two cases, $(G_\chi)_{\bar{x}} = H$ for each $\bar{x} \in [\bar{d}^0, \bar{d}^1]$ by Table 3.4, and F_i's satisfy Condition F1. because $\operatorname{res}^{(G_\chi)_S}_H V_S \cong \operatorname{res}^{(G_\chi)_N}_H V_N$ by (10.1). So, we obtain nonemptiness by Lemma 6.4. The second statement is proved as in Lemma 8.12. □ □

By using these, we obtain nonemptiness of $\Omega_{\bar{D}_R,(W_{d^i})_{i\in I^+}}$.

Proposition 10.4. *For each* $(W_{d^i})_{i\in I^+} \in A_{G_\chi}(S^2,\chi)$*, the set* $\Omega_{\bar{D}_R,(W_{d^i})_{i\in I^+}}$ *is nonempty.*

11. Proof for cases when $\operatorname{pr}(\rho(G_\chi)) = \mathbb{Z}_n$, D_n

In this section, we calculate homotopy of equivariant clutching maps in cases when $\operatorname{pr}(R) = \mathbb{Z}_n$, D_n . We parameterize each edge $|\bar{e}^i_S|$ in $|\bar{\mathcal{K}}_{R,S}|$ for $0 \le i \le m_R - 1$ by the interval $[i, i+1]$ linearly to satisfy $\bar{v}^i_S \mapsto i$ when $m_R \ge 3$. The vertex $\bar{v}^0_S$ is parameterized by 0 or m_R according to context. First, we deal with cases when $R = \mathbb{Z}_n$, $\langle a_n, -b \rangle$. Similarly to Lemma 8.2, we easily obtain the following lemma:

Lemma 11.1. *Assume that* $R = \mathbb{Z}_n$, $\langle a_n, -b \rangle$*. Then,*

$$\operatorname{res}^{G_\chi}_{(G_\chi)_x} E_S \cong \operatorname{res}^{G_\chi}_{(G_\chi)_x} E_N$$

for each E *in* $\operatorname{Vect}_{G_\chi}(|\mathcal{K}_R|,\chi)$ *and* $x \in |\mathcal{L}_R|$*.*

Proof. In these cases, the great circle passing through N, S, x is fixed by $(G_\chi)_x$ because G_χ fixes B. From this, we obtain a proof by Lemma 8.2. □ □

This lemma says that p_{vect} is well-defined in cases when $R = \mathbb{Z}_n$, $\langle a_n, -b \rangle$. In other cases, p_{vect} is well-defined by Lemma 8.2 and Lemma 3.10. For a path γ defined on $[0,1]$, define a path γ_{-i} on $[i, i+1]$ as $\gamma_{-i}(t) = \gamma(t-i)$ for $i \in \mathbb{Z}$.

Proposition 11.2. *Assume that* $R = \mathbb{Z}_n$*. Then, Theorem A holds for the case.*

Proof. For simplicity, we prove the proposition only when $m_R = n \ge 3$. Other cases are similarly proved. For each $(W_q)_{q\in B}$ in $A_{G_\chi}(S^2,\chi)$, pick the G_χ-bundle V_B over B such that $V_B|_q = W_q$ for $q \in B$. Proof is similar to Proposition 9.2.

In this case, $(G_\chi)_x = H$ for each point x in $|\mathcal{L}_R|$ by Table 3.4 so that $\mathcal{A}_{\bar{x}} = \mathcal{A}_{\bar{e}}$ and $\mathcal{A}^0_{\bar{x}} = \operatorname{Iso}_H(V_S, V_N)$ by Lemma 6.4 for each $\bar{x} \in |\bar{\mathcal{L}}_{R,S}|$ and $\bar{e} \in \bar{\mathcal{L}}_{R,S}$ which are nonempty by Lemma 10.3. Let g_0 be an element in G_χ such that $\rho(g_0) = a_n$, i.e. $g_0 d^0 = d^1$. Similarly to Corollary 10.2, $\Omega_{\bar{D}_R,(W_q)_{q\in B}} = \Omega_{\bar{D}_R,V_B}$ is equal to

$$\Big\{\Phi_{\bar{D}_R} \in C^0(\bar{D}_R, V_B) \mid \Phi_{\bar{D}_R}(\bar{x}) \in \operatorname{Iso}_H(V_S,V_N) \text{ for each } \bar{x} \in \bar{D}_R, \\ \text{and } \Phi_{\bar{D}_R}(\bar{d}^1) = g_0 \Phi_{\bar{D}_R}(\bar{d}^0) g_0^{-1}\Big\}$$

by Theorem 10.1. Pick a point $*$ in $\operatorname{Iso}_H(V_S,V_N)$, and put $*' = g_0 * g_0^{-1}$. Since $\operatorname{Iso}_H(V_S,V_N)$ is path-connected, we may assume that each element $\Phi_{\bar{D}_R}$ in $\Omega_{\bar{D}_R,V_B}$

has $*, *'$ at $\bar{d}^0, \bar{d}^1$, respectively. Then, it is easy that $\pi_0(\Omega_{\bar{D}_R, V_B}) \cong \mathbb{Z}$ by Lemma 7.4.

Let $\gamma_0 : \bar{D}_R = [0,1] \to \mathrm{Iso}_H(V_S, V_N)$ be an arbitrary element of $\Omega_{\bar{D}_R, V_B}$ such that $\gamma_0(0) = *$, $\gamma_0(1) = *'$, and let $\sigma_0 : [0,1] \to \mathrm{Iso}_H(V_S, V_N)$ be a loop such that $\sigma_0(0) = \sigma_0(1) = *$ and $[\sigma_0]$ is a generator of $\pi_1\big(\mathrm{Iso}_H(V_S, V_N), *\big)$. So, $\sigma_0.\gamma_0$ is contained in $\Omega_{\bar{D}_R, V_B}$. We would show that the difference $c_1\big(F_{V_B}/\sigma_0.\gamma_0\big) - c_1\big(F_{V_B}/\gamma_0\big)$ is equal to $n\chi(id)$ up to sign where $l_R = n$. For this, we need to describe the equivariant clutching maps precisely. Let $\Phi = \cup_{q \in B}\, \varphi_q$ be the extension of γ_0, i.e. $\varphi_S(t) = \gamma_0(t)$ for $t \in [0,1]$. By equivariance of Φ,

$$\begin{aligned}\varphi_S(t) &= g_0^i \varphi_S(t-i) g_0^{-i} \\ &= g_0^i \gamma_0(t-i) g_0^{-i} \\ &= g_0^i (\gamma_0)_{-i}(t) g_0^{-i}\end{aligned}$$

for $0 \le i \le n-1$ and $t \in [i, i+1]$. That is,

$$\varphi_S = \gamma_0 \vee \big(g_0(\gamma_0)_{-1} g_0^{-1}\big) \vee \cdots \vee \big(g_0^{n-1}(\gamma_0)_{-n+1} g_0^{-n+1}\big).$$

And, let $\Phi' = \cup_{q \in B}\, \varphi'_q$ be the extension of $\sigma_0.\gamma_0$. Then,

$$\varphi'_S = (\sigma_0.\gamma_0) \vee \big(g_0(\sigma_0.\gamma_0)_{-1} g_0^{-1}\big) \vee \cdots \vee \big(g_0^{n-1}(\sigma_0.\gamma_0)_{-n+1} g_0^{-n+1}\big). \tag{11.1}$$

Here, $g_0^i(\sigma_0.\gamma_0)_{-i} g_0^{-i}$ for $0 \le i \le n-1$ is equal to

$$g_0^i(\sigma_0)_{-i}\ g_0^{-i} \cdot g_0^i(\gamma_0)_{-i}\ g_0^{-i},$$

and we can move $g_0^i(\sigma_0)_{-i}\ g_0^{-i}$'s in (11.1) (nonequivariantly) homotopically to $*$. For example,

$$\begin{aligned}&[(\sigma_0.\gamma_0) \vee \big(g_0(\sigma_0)_{-1}\ g_0^{-1} \cdot g_0(\gamma_0)_{-1}\ g_0^{-1}\big)] \\ =&[(\sigma_0.\gamma_0.g_0\sigma_0 g_0^{-1}) \vee \big(g_0(\gamma_0)_{-1}\ g_0^{-1}\big)] \\ =&[\big(\sigma_0.\bar{\gamma}_0^{-1}(g_0\sigma_0 g_0^{-1}).\gamma_0\big) \vee \big(g_0(\gamma_0)_{-1}\ g_0^{-1}\big)] \\ =&[(\sigma_0.\sigma_0.\gamma_0) \vee \big(g_0(\gamma_0)_{-1}\ g_0^{-1}\big)].\end{aligned}$$

Since φ'_S contains n σ_0's, difference between Chern classes is $n\chi(id)$ up to sign by Lemma 7.1. And, we can conclude that c_1 on $\Omega_{\bar{D}_R, V_B}$ is injective and its image is equal to the set $\{\chi(\mathrm{id})(nk + k_0) \mid k \in \mathbb{Z}\}$ where k_0 is dependent on the pair. Therefore, we obtain a proof by Proposition 5.3.(1). □ □

Proposition 11.3. *Assume that $R = \langle a_n, -b\rangle$. Then, Theorem C holds for the case.*

Proof. For simplicity, we prove the proposition only when $m_R \ge 3$. Other cases are similarly proved. For each $(W_q)_{q \in B}$ in $A_{G_\chi}(S^2, \chi)$, pick the G_χ-bundle V_B over B such that $V_B|_q = W_q$ for $q \in B$.

In this case, $(G_\chi)_x = H$ and $\mathcal{A}^0_{\bar{x}} = \mathrm{Iso}_H(V_S, V_N)$ for each interior $\bar{x} \in \bar{D}_R$ and its image $x = |\pi|(\bar{x})$ by Table 3.4 and Lemma 6.4. Here, $\mathrm{Iso}_H(V_S, V_N)$ is nonempty by Lemma 10.3. Similarly, $\mathcal{A}^0_{\bar{x}} = \mathrm{Iso}_{(G_\chi)_x}(V_S, V_N)$ is nonempty for $\bar{x} = \bar{d}^0, \bar{d}^1$ and its image $x = |\pi|(\bar{x})$. Similarly to Corollary 10.2, $\Omega_{\bar{D}_R, (W_q)_{q \in B}} = \Omega_{\bar{D}_R, V_B}$ is equal to

$$\Big\{\Phi_{\bar{D}_R} \in C^0(\bar{D}_R, V_B) \mid \Phi_{\bar{D}_R}(\bar{x}) \in \mathrm{Iso}_H(V_S, V_N) \text{ for each } \bar{x} \in \bar{D}_R, \\ \text{and } \Phi_{\bar{D}_R}(\bar{x}) \in \mathcal{A}^0_{\bar{x}} \text{ for } \bar{x} = \bar{d}^0, \bar{d}^1\Big\}$$

by Theorem 10.1. Then, we obtain a proof by Lemma 7.4 because the inclusion from $\mathrm{Iso}_{(G_\chi)_x}(V_S, V_N)$ to $\mathrm{Iso}_H(V_S, V_N)$ for $x = d^0, d^1$ induces surjection in the level of fundamental groups by Lemma 7.2. □ □

Proposition 11.4. *Assume that R is equal to one of the following:*
$\mathrm{D}_n \times Z$, *even* n, $\langle -a_n, -b\rangle$, *odd* $n/2$, $\langle -a_n, b\rangle$, *odd* $n/2$,
$\mathbb{Z}_n \times Z$, *even* n, $\langle -a_n\rangle$, *odd* $n/2$.
Then, Theorem C holds for these cases.

Proof. Proof is similar to Proposition 9.1. □ □

Proposition 11.5. *Assume that $R = \mathrm{D}_n$. Then, Theorem A holds for the case where $l_R = 2n$.*

Proof. Proof is similar to Proposition 9.2. □ □

Proposition 11.6. *Assume that R is equal to one of the following:*
$\mathrm{D}_n \times Z$, *odd* n, $\langle -a_n, b\rangle$, *even* $n/2$, $\langle -a_n, -b\rangle$, *even* $n/2$.
Then, Theorem C holds for these cases.

Proof. For each $(W_{d^i})_{i\in I^+}$ in $A_{G_\chi}(S^2,\chi)$, put $V_B = G_\chi \times_{(G_\chi)_{d^{-1}}} W_{d^{-1}}$. In these cases, $\mathcal{A}_{\bar{x}}$ is nonempty for each $\bar{x} \in \bar{D}_R$ by Lemma 10.3. Let us investigate $\mathcal{A}_{\bar{x}}$ more precisely. First, $(G_\chi)_x = H$ and $\mathcal{A}^0_{\bar{x}} = \mathrm{Iso}_H(V_S, V_N)$ for each interior $\bar{x} \in \bar{D}_R$ and its image $x = |\pi|(\bar{x})$ by Table 3.4 and Lemma 6.4. Also, $(G_\chi)_x/H \cong \mathbb{Z}_2$ for $x = d^0$ or d^1 (say it d^0), and $\rho\big((G_\chi)_x\big)$ has an element of the form $-a^i_n b$ fixing S for some i by Table 3.4. From this, $(G_\chi)_{d^0}$ fixes the great circle containing S, N, d^0 so that V_S and V_N are $(G_\chi)_{d^0}$-isomorphic as in Lemma 11.1. And, $\mathcal{A}^0_{\bar{d}^0}$ is homeomorphic to $\mathrm{Iso}_{(G_\chi)_{d^0}}(V_S, V_N)$ by Lemma 6.4. Pick an element $\psi_{\bar{d}^1}$ in $\mathcal{A}_{\bar{d}^1}$ which determines W_{d^1}. Note that d^0 and d^1 are not in a G_χ-orbit because one is a vertex and the other is not a vertex in D_R by Table 1.1. Similarly to (8.1) of Proposition 8.13, $\Omega_{\bar{D}_R,(W_{d^i})_{i\in I^+}}$ is equal to

$$\Big\{\Phi_{\bar{D}_R} \in C^0(\bar{D}_R, V_B) \mid \Phi_{\bar{D}_R}(\bar{x}) \in \mathrm{Iso}_H(V_S, V_N) \text{ for each } \bar{x} \in \bar{D}_R,$$
$$\Phi_{\bar{D}_R}(\bar{d}^0) \in \mathrm{Iso}_{(G_\chi)_{d^0}}(V_S, V_N), \text{ and } \Phi_{\bar{D}_R}(\bar{d}^1) \in (\mathcal{A}_{\bar{d}^1})^0_{\psi_{\bar{d}^1}}\Big\}$$

by Theorem 10.1 which is nonempty by Proposition 10.4. Here, $(\mathcal{A}_{\bar{d}^1})_{\psi_{\bar{d}^1}}$ is simply connected by Proposition 6.8 because $(G_\chi)_{d^1}/H \cong \mathbb{Z}_2$ and $\bar{d}^1$, $|c|(\bar{d}^1)$ are in a G_χ-orbit by Table 3.4. And, the inclusion from $\mathrm{Iso}_{(G_\chi)_{d^0}}(V_S, V_N)$ to $\mathrm{Iso}_H(V_S, V_N)$ induces surjection in the level of fundamental groups by Lemma 7.2. Therefore, we obtain a proof by Lemma 7.4. □ □

Now, we deal with remaining two cases $R = \mathbb{Z}_n \times Z$ with odd n or $\langle -a_n\rangle$ with even $n/2$. For a path γ defined on $[0,1]$, define a path γ^{tr} on $[0,1]$ as $\gamma^{tr}(t) = \gamma(1-t)$ for $t \in [0,1]$.

Proposition 11.7. *Assume that $R = \mathbb{Z}_n \times Z$ with odd n or $\langle -a_n\rangle$ with even $n/2$. For each triple $(W_{d^i})_{i\in I^+} \in A_{G_\chi}(S^2,\chi)$, we have*

$$\pi_0(\Omega_{\bar{D}_R,(W_{d^i})_{i\in I^+}}) \cong \mathbb{Z}.$$

And, $c_1\big(F_{V_B}/\Phi\big)$ is constant for Φ in $\Omega_{\bar{D}_R,(W_{d^i})_{i\in I^+}}$ when $V_B = G_\chi \times_{(G_\chi)_{d^{-1}}} W_{d^{-1}}$.

Proof. In these cases, $\mathcal{A}_{\bar{x}}$ is nonempty for each $\bar{x} \in \bar{D}_R$ by Lemma 10.3. Since $(G_\chi)_x = H$ for each $\bar{x} \in \bar{D}_R$ and its image $x = |\pi|(\bar{x})$ by Table 3.4, the triple satisfies

$$W_{d^{-1}} = V_S \text{ and } W_{d^i} \cong \mathrm{res}^{(G_\chi)_S}_H V_S$$

for $i \in I$ by Definition 1.1, i.e. each triple is determined by the third entry $W_{d^{-1}}$. So, $\Omega_{\bar{D}_R,(W_{d^i})_{i\in I^+}} = \Omega_{\bar{D}_R,V_B}$ because V_B is also determined by $W_{d^{-1}}$. And, $\mathcal{A}^0_{\bar{x}} = \mathcal{A}^0_{\bar{e}} = \mathrm{Iso}_H(V_S, V_N)$ for $\bar{x} \in |\bar{\mathcal{L}}_{R,S}|$, $\bar{e} \in \bar{\mathcal{L}}_{R,S}$ by Lemma 6.4 so that $\pi_0(\mathcal{A}^0_{\bar{x}}) \cong \mathbb{Z}$ by Schur's Lemma because V_S is H-isotypical.

Let $g_0 \in G_\chi$ be an element such that

(1) $\rho(g_0) = -a_n^{n+1/2}$ if $\rho(G_\chi) = \mathbb{Z}_n \times Z$ with odd n,
(2) $\rho(g_0) = -a_n^{n/2+1}$ if $\rho(G_\chi) = \langle -a_n \rangle$ with even $n/2$.

In both cases, $g_0 \bar v_S^i = \bar v_N^{i+1}$ for each $i \in \mathbb{Z}_{m_R}$ in $\bar{\mathcal{L}}_R$. Similarly to Corollary 10.2, $\Omega_{\bar D_R,(W_{d^i})_{i\in I^+}} = \Omega_{\bar D_R,V_B}$ is equal to

$$\Big\{ \Phi_{\bar D_R} \in C^0(\bar D_R, V_B) \mid \Phi_{\bar D_R}(\bar x) \in \mathrm{Iso}_H(V_S,V_N) \text{ for each } \bar x \in \bar D_R, \\ \text{and } \Phi_{\bar D_R}(\bar d^1) = g_0 \Phi^{-1}_{\bar D_R}(\bar d^0) g_0^{-1} \Big\}$$

by Theorem 10.1. Pick an element $*$ in $\mathcal{A}^0_{\bar d^0} = \mathrm{Iso}_H(V_S,V_N)$, and put $*' = g_0 *^{-1} g_0^{-1}$ in $\mathcal{A}^0_{\bar d^1} = \mathrm{Iso}_H(V_S,V_N)$. Since $\mathrm{Iso}_H(V_S,V_N)$ is path-connected, we may assume that each element $\Phi_{\bar D_R}$ in $\Omega_{\bar D_R,V_B}$ satisfies

$$\Phi_{\bar D_R}(\bar d^0) = * \text{ and } \Phi_{\bar D_R}(\bar d^1) = *'.$$

So, $\pi_0(\Omega_{\bar D_R,V_B}) \cong \mathbb{Z}$ by Lemma 7.4 because $\pi_1\big(\mathrm{Iso}_H(V_S,V_N)\big) \cong \mathbb{Z}$.

Next, we calculate the first Chern class. Our calculation is done in a similar way with Proposition 11.2. Pick an arbitrary element $\Phi_{\bar D_R}$ in $\Omega_{\bar D_R,V_B}$ such that $\Phi_{\bar D_R}(\bar d^0) = *$ and $\Phi_{\bar D_R}(\bar d^1) = *'$. Put $\gamma(t) = \Phi_{\bar D_R}(t)$ on $t \in [0,1]$. Take a loop $\sigma_0 : [0,1] \to \mathrm{Iso}_H(V_S,V_N)$ such that $\sigma_0(0) = \sigma_0(1) = *$ and $[\sigma_0]$ is a generator of $\pi_1(\mathrm{Iso}_H(V_S,V_N),*)$. To prove our result, we only have to show that $c_1\big(F_{V_B}/\sigma_0.\gamma\big) = c_1\big(F_{V_B}/\gamma\big)$ where $\sigma_0.\gamma$ is in $\Omega_{\bar D_R,V_B}$ by Corollary 10.2. Let $\Phi = \cup_{q\in B}\, \varphi_q$ be the extension of γ, i.e. $\varphi_S(t) = \gamma(t)$ on $t \in [0,1]$. Then, equivariance of Φ shows that

$$\begin{aligned} \varphi_S(t) &= g_0 \varphi_N(g_0^{-1}t) g_0^{-1} \\ &= g_0 \varphi_S(g_0^{-1}t)^{-1} g_0^{-1} \\ &= g_0 \gamma(t-1)^{-1} g_0^{-1} \end{aligned}$$

for $t \in [1,2]$. That is, $\varphi_S = \gamma \wedge g_0 \gamma_{-1}^{-1} g_0^{-1}$ on $[0,2]$. By using this, if $\Phi' = \cup_{q\in B}\, \varphi'_q$ is the extension of $\sigma_0.\gamma$, then we have

$$\begin{aligned} \varphi'_S &= \sigma_0.\gamma \wedge g_0\big((\sigma_0)_{-1}^{-1} \cdot \gamma_{-1}^{-1}\big) g_0^{-1} \\ &= \sigma_0.\gamma \wedge \Big(g_0 (\sigma_0)_{-1}^{-1} g_0^{-1} \cdot g_0 \gamma_{-1}^{-1} g_0^{-1} \Big) \end{aligned}$$

on $[0,2]$. Here, two generators $[\sigma_0]$ in $\pi_1\big(\mathrm{Iso}_H(V_S,V_N),*\big)$ and $\big[g_0(\sigma_0)_{-1}^{-1}g_0^{-1}\big]$ in $\pi_1\big(\mathrm{Iso}_H(V_S,V_N),*'\big)$ cancel each other in $\pi_1(\mathrm{Iso}_H(V_S,V_N))$. This is in contrast with the calculation of Proposition 11.2. Since the whole φ_S and φ'_S are equivariantly determined by γ and $\sigma_0.\gamma$ as in the proof of Proposition 9.2, we obtain $c_1\big(F_{V_B}/\sigma_0.\gamma\big) = c_1\big(F_{V_B}/\gamma\big)$. □ □

The proposition says that Proposition 5.3 does not applies to these two cases. So, we need to apply Lemma 5.2 to these cases. For this, we prove two technical lemmas.

Lemma 11.8. *Assume that $R = \mathbb{Z}_n \times Z$ with odd n or $\langle -a_n \rangle$ with even $n/2$. For each $(W_{d^i})_{i\in I^+} \in A_{G_\chi}(S^2,\chi)$, put $V_B = G_\chi \times_{(G_\chi)_{d^{-1}}} W_{d^{-1}}$. Let $\eta_S : |\bar{\mathcal{K}}_{R,S}| \times V_S \to |\bar{\mathcal{K}}_{R,S}| \times V_S$ be a $(G_\chi)_S$-isomorphism. For an element g in G_χ such that $gS = N$, let $\eta_N : |\bar{\mathcal{K}}_{R,N}| \times V_N \to |\bar{\mathcal{K}}_{R,N}| \times V_N$ be defined by $\eta_N(x) = g\eta_S(g^{-1}x)g^{-1}$. Then,*

(1) *$\eta = \cup_{q\in B}\, \eta_q$ is the unique G_χ-isomorphism of F_{V_B} extending η_S.*
(2) *For a map $\Phi = \bigcup_{q\in B}\, \varphi_q$ in Ω_{V_B}, the map $\Phi' = \bigcup_{q\in B}\, \varphi'_q$ with $\varphi'_q = \eta_{q'} \varphi_q \eta_q^{-1}$ is also an equivariant clutching map satisfying the following commutative diagram*

$$\begin{tikzcd}
{|\bar{\mathcal{L}}_{R,q}| \times V_q} \arrow[r, "\eta_q|_{|\bar{\mathcal{L}}_{R,q}|}"] \arrow[d, "\varphi_q"] & {|\bar{\mathcal{L}}_{R,q}| \times V_q} \arrow[d, "\varphi'_q"] \\
{|\bar{\mathcal{L}}_{R,q'}| \times V_{q'}} \arrow[r, "\eta_{q'}|_{|\bar{\mathcal{L}}_{R,q'}|}"] & {|\bar{\mathcal{L}}_{R,q'}| \times V_{q'}}
\end{tikzcd}$$

when $B = \{q, q'\}$.

Proof. Well-definedness is an only issue here, and normality of $(G_\chi)_S = (G_\chi)_N$ in G_χ is used for this. □ □

Remark 11.9.

(1) In this lemma, η gives a G_χ-isomorphism between F_{V_B}/Φ and F_{V_B}/Φ'.

(2) Let $\eta_{S,t}$ for $t \in [0,1]$ be a homotopy of $(G_\chi)_S$-isomorphisms of $|\bar{\mathcal{K}}_{R,S}| \times V_S$. Let $\Phi'_t = \bigcup_{q \in B} \varphi'_{q,t}$ for each $t \in [0,1]$ be the equivariant clutching map determined by $\eta_{S,t}$ and Φ in this lemma. Then, F_{V_B}/Φ'_0 and F_{V_B}/Φ'_1 are G_χ-isomorphic by Lemma 5.1. □

Lemma 11.10. *Assume that* $R = \mathbb{Z}_n \times Z$ *with odd* n *or* $\langle -a_n \rangle$ *with even* $n/2$. *For each* $(W_{d^i})_{i \in I^+} \in A_{G_\chi}(S^2, \chi)$, *put* $V_B = G_\chi \times_{(G_\chi)_{d^{-1}}} W_{d^{-1}}$. *Then, each* G_χ-*isomorphism* $\eta = \cup_{q \in B} \eta_q$ *of* F_{V_B} *is equivariantly homotopic to a* G_χ-*isomorphism* $\eta' = \cup_{q \in B} \eta'_q$ *of* F_{V_B} *such that*

$$\eta'_S|_{|\bar{e}^0_S| \cup |\bar{e}^1_S|} \ = \ \gamma \wedge (\gamma^{tr})_{-1}$$

for some loop $\gamma : \bar{D}_R = [0,1] \to \mathrm{Iso}_H(V_S)$ *satisfying* $\gamma(\bar{v}^0_S) = \gamma(\bar{v}^1_S) = \mathrm{id}$ *where* $(\gamma^{tr})_{-1}$ *is defined on* $[1,2]$.

Proof. Note that $R_S = \mathbb{Z}_{m_R/2}$ for both cases, and pick an element $g_1 \in (G_\chi)_S$ such that $g_1 \bar{v}^0_S = \bar{v}^2_S$. Consider $|\bar{\mathcal{K}}_{R,S}|$ as the quotient $|\bar{\mathcal{L}}_{R,S}| \times [0,1]/|\bar{\mathcal{L}}_{R,S}| \times 0$, and parameterize points of it by $(\bar{x}, t)$ with $\bar{x} \in |\bar{\mathcal{L}}_{R,S}|$ and $t \in [0,1]$. Let $\gamma_i : [1/2, 1] \to \mathrm{Iso}_H(V_S)$ be paths such that

$$\gamma_i(1/2) = \eta_S(\bar{v}^i_S), \quad \gamma_i(1) = \mathrm{id}, \quad \gamma_2(t) = g_1 \gamma_0(t) g_1^{-1}$$

for each t, $i \in \{0, 2\}$. Then, define

$$\eta'_S : |\bar{\mathcal{L}}_{R,S}| \times [0,1/2] \bigcup \Big(\cup_{i=0,2} \ (\bar{v}^i_S \times [1/2,1]) \Big) \longrightarrow \mathrm{Iso}_H(V_S)$$

to satisfy

$$\begin{aligned} \eta'_S(\bar{x}, t) &= \eta_S(\bar{x}, 2t) && \text{for } t \in [0, 1/2], \\ \eta'_S(\bar{v}^i_S, t) &= \gamma_i(t) && \text{for } t \in [1/2, 1], i \in \{0, 2\}. \end{aligned}$$

We can extend η'_S to $|\bar{\mathcal{L}}_{R,S}| \times [0,1/2] \bigcup \big(|\bar{e}^0_S| \ \cup \ |\bar{e}^1_S| \big) \times [1/2, 1]$. Then, η'_S is equivariantly extended to all $|\bar{\mathcal{K}}_{R,S}|$ so that η'_S is a $(G_\chi)_S$-isomorphism of $|\bar{\mathcal{K}}_{R,S}| \times V_S$. Since η'_S is defined on $|\bar{\mathcal{K}}_{R,S}|$, $\eta'_S|_{|\bar{\mathcal{L}}_{R,S}|}$ is a (nonequivariantly) homotopically trivial map when we consider $\eta'_S|_{|\bar{\mathcal{L}}_{R,S}|}$ as a map from S^1 to $\mathrm{Iso}_H(V_S)$. Let $\alpha : |\bar{e}^0_S| \cup |\bar{e}^1_S| \to \mathrm{Iso}_H(V_S)$ be the restriction $\eta'_S|_{|\bar{e}^0_S| \cup |\bar{e}^1_S|}$. Note that $\alpha(\bar{v}^0_S) = \alpha(\bar{v}^2_S) = \mathrm{id}$. Since $\eta'_S|_{|\bar{\mathcal{L}}_{R,S}|}$ is the G_χ-orbit of α, α should be also (nonequivariantly) homotopically trivial. Therefore, we may assume that $\alpha = \gamma \wedge (\gamma^{tr})_{-1}$ for some loop $\gamma : \bar{D}_R \to \mathrm{Iso}_H(V_S)$ satisfying $\gamma(\bar{v}^0_S) = \gamma(\bar{v}^1_S) = \mathrm{id}$. □ □

Now, we can prove Theorem B.

Proof of Theorem B. For each $(W_{d^i})_{i \in I^+}$, put $V_B = G_\chi \times_{(G_\chi)_{d^{-1}}} W_{d^{-1}}$. Since $\Omega_{\bar{D}_R, V_B} = \Omega_{\bar{D}_R, (W_{d^i})_{i \in I^+}}$ as we have seen in the proof of Proposition 11.7, we have $p^{-1}_{\mathrm{vect}}\big((W_{d^i})_{i \in I^+}\big) = \mathrm{Vect}_{G_\chi}(S^2, \chi)_{V_B}$. So, it suffices to show that the set $\mathrm{Vect}_{G_\chi}(S^2, \chi)_{V_B}$ has exactly two elements for each V_B to prove the first statement.

Let $\Phi = \cup_{q \in B}\ \varphi_q$ be a map in Ω_{V_B}. Put $\Phi(\bar{d}^0) = *$ and $\Phi(\bar{d}^1) = *'$. Let $g_0 \in G_\chi$ be the element in the proof of Proposition 11.7 such that $g_0 \bar{v}_S^i = \bar{v}_N^{i+1}$ for each i. We would determine which classes of $\Omega_{\bar{D}_R, V_B}$ give isomorphic equivariant vector bundles. Take a loop $\sigma_0 : \bar{D}_R = [0,1] \to \mathrm{Iso}_H(V_S)$ such that $\sigma_0(0) = \sigma_0(1) = \mathrm{id}$ and $[\sigma_0]$ is a generator of $\pi_1\big(\mathrm{Iso}_H(V_S), \mathrm{id}\big)$. For each G_χ-isomorphism $\eta = \cup_{q \in B}\ \eta_q$ of F_{V_B}, we may assume that $\eta_S|_{|\bar{e}_S^0| \cup |\bar{e}_S^1|} = \sigma_0^j \wedge (\sigma_0^{tr})_{-1}^j$ for some $j \in \mathbb{Z}$ by Lemma 11.10. Put

$$\varphi'_S(\bar{x}) = \eta_N(|c|(\bar{x}))\varphi_S(\bar{x})\eta_S(\bar{x})^{-1}$$

for all $\bar{x} \in \bar{D}_R$. By equivariance,

$$\eta_N(|c|(\bar{x})) = g_0^{-1}\eta_S\big(g_0|c|(\bar{x})\big)g_0 = g_0^{-1}\eta_S(\bar{x}+1)g_0.$$

Since $\eta_S|_{|\bar{e}_S^0| \cup |\bar{e}_S^1|} = \sigma_0^j \wedge (\sigma_0^{tr})_{-1}^j$,

$$\varphi'_S(\bar{x}) = g_0^{-1}\ \sigma_0^{tr}(\bar{x})^j\ g_0\ \varphi_S(\bar{x})\ \sigma_0(\bar{x})^{-j}$$

for $\bar{x} \in \bar{D}_R$. And, this path is nonequivariantly homotopic to $\varphi_S(\bar{x})\sigma_0(\bar{x})^{-2j}$ in $\big[[0,1], 0, 1; \mathrm{Iso}_H(V_S, V_N), *, *'\big]$. Since each class in $\Omega_{\bar{D}_R, V_B}$ is nonequivariantly homotopic to one of $\varphi_S(\bar{x})\sigma_0(\bar{x})^i$ for $i \in \mathbb{Z}$ by Proposition 11.7, this says that the equivariant vector bundle determined by an equivariant clutching map on $\bar{D}_R$ with respect to V_B depends only on the parity of i by Lemma 5.2. Therefore, $\mathrm{Vect}_{G_\chi}(S^2, \chi)_{V_B}$ have two different G_χ-bundles. By the arguments on parity, we similarly obtain the second statement. ☐ ☐

12. Proof for cases when $\mathrm{pr}(\rho(G_\chi)) = \mathrm{SO}(2)$, $\mathrm{O}(2)$

Assume that $\rho(G_\chi) = R$ for some one-dimensional R in Table 1.1. In these cases, S^2 can not have equivariant simplicial complex structure. So, we introduce new notations. Consider the disjoint union $\bar{S}^2 = S_S^2 \amalg S_N^2$ of the lower and upper hemispheres S_S^2 and S_N^2 of S^2, and denote by $\bar{S}_S^2$ and $\bar{S}_N^2$ hemispheres S_S^2 and S_N^2 in $\bar{S}^2$, respectively. Denote by π the natural quotient maps from $\bar{S}^2$ to S^2. Denote by $\bar{S}_S^1$ and $\bar{S}_N^1$ boundaries of $\bar{S}_S^2$ and $\bar{S}_N^2$ in $\bar{S}^2$, respectively. And, denote by $\bar{S}^1$ the disjoint union $\bar{S}_S^1 \amalg \bar{S}_N^1$ which is the preimage of the equator through π. Put $B = \{S, N\} \subset S^2$ on which R (and G_χ) acts. The subset $\pi^{-1}(B)$ in $\bar{S}^2$ is often confused with B in S^2. Denote by $\bar{v}_q^0$ be the point $\pi^{-1}(v^0) \cap \bar{S}_q^1$ for $q \in B$ where $v^0 = (1,0,0)$. Note that if $R = \mathrm{SO}(2)$, $\langle \mathrm{SO}(2), -b \rangle$, then R fixes B, and otherwise R acts transitively on B. We can redefine notations V_B, F_{V_B}, Φ, F_{V_B}/Φ, $\imath_\Omega$, p_Ω, p_{π_0}, Ω_{V_B}, $\cdots$ of Section 10 by replacing $|\mathcal{K}_R|$, $|\bar{\mathcal{K}}_R|$, $|\bar{\mathcal{K}}_{R,q}|$, $|\bar{\mathcal{L}}_R|$, $|\bar{\mathcal{L}}_{R,q}|$ with S^2, $\bar{S}^2$, $\bar{S}_q^2$, $\bar{S}^1$, $\bar{S}_q^1$, respectively. Here, notations for cases when $R = \mathrm{SO}(2)$, $\langle \mathrm{SO}(2), -b \rangle$ and $R \neq \mathrm{SO}(2)$, $\langle \mathrm{SO}(2), -b \rangle$ are redefined in the same way with cases when $R = \mathbb{Z}_n$, $\langle a_n, -b \rangle$ and $R \neq \mathbb{Z}_n$, $\langle a_n, -b \rangle$ of Section 10, respectively.

Put $\bar{\mathbf{x}} = \{\bar{x}_j \mid j \in \mathbb{Z}_2\}$ with $\bar{x}_0 = \bar{v}_S^0$ and $\bar{x}_1 = \bar{v}_N^0$. Let $\mathcal{A}_{\bar{v}_S^0}$ be the set of equivariant pointwise clutching maps with respect to the $(G_\chi)_{v^0}$-bundle $\big(\mathrm{res}_{(G_\chi)_{v^0}}^{G_\chi} F_{V_B}\big)|_{\bar{\mathbf{x}}}$.

Proposition 12.1. *Assume that R is equal to one of the following:*

$$\mathrm{O}(2) \times Z, \qquad \mathrm{O}(2), \qquad \langle \mathrm{SO}(2),\ -a_2 \rangle.$$

Then, Theorem C holds for these cases.

Proof. Put $V_B = G_\chi \times_{(G_\chi)_{d^{-1}}} W_{d^{-1}}$ for each $(W_{d^i})_{i \in I^+}$ in $A_{G_\chi}(S^2, \chi)$. By Proposition 5.3.(2), we only have to show that $\pi_0(\Omega_{(W_{d^i})_{i \in I^+}})$ consists of exactly one element for each $(W_{d^i})_{i \in I^+}$. First, we show nonemptiness of $\mathcal{A}_{\bar{v}_S^0}$. For those R's, the $(G_\chi)_{v^0}$-bundle $F = \big(\mathrm{res}_{(G_\chi)_{v^0}}^{G_\chi} F_{V_B}\big)|_{\bar{\mathbf{x}}}$ satisfies Condition F2. by Table 3.4. Since $d^0 = d^1 = v^0$, we have $W_{d^0} \cong W_{d^1}$ which is an $(G_\chi)_{v^0}$ extension of $F_{\bar{x}_0} =$

$\mathrm{res}^{(G_\chi)_S}_{(G_\chi)_S \cap (G_\chi)_{v^0}} W_{d^{-1}}$ by Definition 1.1. This means that $\mathcal{A}_{\bar{v}^0_S}$ is nonempty by Theorem 6.6 as in the proof of Lemma 8.12. Pick an element $\psi_{\bar{v}^0_S}$ in $\mathcal{A}_{\bar{v}^0_S}$ which determines W_{d^0}

Consider the evaluation map at $\bar{v}^0_S$

$$\Omega_{(W_{d^i})_{i \in I^+}} \to (\mathcal{A}_{\bar{v}^0_S})^0_{\psi_{\bar{v}^0_S}}, \quad \Phi \mapsto \Phi(\bar{v}^0_S).$$

By definition of $\Omega_{(W_{d^i})_{i \in I^+}}$, the isotropy representation $(F_{V_B}/\Phi)_{d^0}$ for each Φ is isomorphic to W_{d^0} so that $\Phi(\bar{v}^0_S)$ is contained in $(\mathcal{A}_{\bar{v}^0_S})^0_{\psi_{\bar{v}^0_S}}$, i.e. the evaluation map is well-defined. We show that it is a one-to-one correspondence. For those R's, G_χ acts transitively on $\bar{S}^1$. So, each map Φ in Ω_{V_B} is determined by $\Phi(\bar{v}^0_S)$ through equivariance, and the evaluation is injective. For each ψ in $(\mathcal{A}_{\bar{v}^0_S})_{\psi_{\bar{v}^0_S}}$, we construct a map Φ to satisfy

$$\Phi(g\bar{v}^0_S) = g\psi(\bar{v}^0_S)g^{-1}.$$

for each $g \in G_\chi$, especially $\Phi|_{\bar{\mathbf{x}}} = \psi$. We can show that this is a well-defined equivariant clutching map with respect to V_B so that the evaluation is surjective. And, it induces the bijection from $\pi_0(\Omega_{(W_{d^i})_{i \in I^+}})$ to a one point set $\pi_0\big((\mathcal{A}_{\bar{v}^0_S})_{\psi_{\bar{v}^0_S}}\big)$. Therefore, we obtain a proof. □ □

Proposition 12.2. *Assume that R is equal to one of* $\mathrm{SO}(2)$, $\langle \mathrm{SO}(2), -b \rangle$. *Then, Theorem C holds for these cases.*

Proof. Pick a G_χ-vector bundle V_B over B such that $V_B|_q = W_q$ for each pair $(W_q)_{q \in B} \in A_{G_\chi}(S^2, \chi)$ and $q \in B$. By Proposition 5.3.(2), we only have to show that $\pi_0(\Omega_{(W_q)_{q \in B}})$ consists of exactly one element for each $(W_q)_{q \in B}$. For those R's, the $(G_\chi)_{v^0}$-bundle $F = \big(\mathrm{res}^{G_\chi}_{(G_\chi)_{v^0}} F_{V_B}\big)|_{\bar{\mathbf{x}}}$ satisfies Condition F1. by Table 3.4 and Definition 1.1. This means that $\mathcal{A}^0_{\bar{v}^0_S}$ is isomorphic to nonempty $\mathrm{Iso}_H(V_S, V_N)$ by Proposition 6.4. Consider the evaluation map at $\bar{v}^0_S$

$$\Omega_{(W_q)_{q \in B}} \to \mathcal{A}^0_{\bar{v}^0_S}, \ \Phi \mapsto \Phi(\bar{v}^0_S).$$

In these cases, G_χ acts transitively only on $\bar{S}^1_S$, but each map Φ in Ω_{V_B} is determined by $\Phi(\bar{v}^0_S)$ through equivariance and inverse. It can be shown that the evaluation is bijective as in Proposition 12.1. And, it induces the bijective map from $\pi_0(\Omega_{(W_q)_{q \in B}})$ to $\pi_0\big(\mathcal{A}^0_{\bar{v}^0_S}\big)$ which is one-point set. Therefore, we obtain a proof. □ □

13. Equivariant line bundles over effective G_χ/H-actions

In this section, we prove Theorem D and calculate Chern classes. Since H is the kernel of the G_χ-action on S^2, S^2 delivers the G_χ/H-action. Since $\rho(G_\chi) = R$ for some R of Table 1.1 by assumption, we may assume that G_χ/H is equal to R and the G_χ/H-action is equal to the R-action on S^2.

Proof of Theorem D. We prove this only for the case when $R = \mathbb{Z}_n$ because other cases are proved similarly. Let U be the H-representation with the character χ. Let $\bar{U}$ be a $(G_\chi)_q$-extension of U for $q \in B$ whose existence is guaranteed by Theorem 14.1. Pick a bundle E in $\mathrm{Vect}_{G_\chi}(S^2, \chi)$, and put $(W_q)_{q \in B} = p_{\mathrm{vect}}(E)$. Then, W_q's are direct sums of $\bar{U} \otimes \Omega(l)$'s by Corollary 14.2 because $(G_\chi)_q/H \cong \mathbb{Z}_n$ for $q \in B$. Pick arbitrary direct summands $\bar{U} \otimes \Omega(l_0)$ and $\bar{U} \otimes \Omega(l_1)$ of W_S and W_N, respectively. Define $(W'_q)_{q \in B}$ by

$$W'_S = \bar{U} \otimes \Omega(l_0) \quad \text{and} \quad W'_N = \bar{U} \otimes \Omega(l_1).$$

By definition of $A_{G_\chi}(S^2,\chi)$, the pair $(W'_q)_{q\in B}$ is contained in $A_{G_\chi}(S^2,\chi)$. Since $p_{\text{vect}}^{-1}\Big((W'_q)_{q\in B}\Big)$ is nonempty by Theorem A, there exists a bundle L with rank $\chi(\text{id})$ in $\text{Vect}_{G_\chi}(S^2,\chi)$. Existence of L proves the isomorphism by [CKMS, Lemma 2.2]. By similar arguments, we can show that $A_R(S^2,\text{id})$ is generated by all the elements with one-dimensional entries. By using this and Theorem A, B, we can show that $\text{Vect}_R(S^2)$ is generated by line bundles.

Now, we calculate the number of elements in $A_R(S^2,\text{id})$ with one-dimensional entries. By Definition 1.1, each pair $(W_q)_{q\in B}$ in $\text{Rep}(R)^2$ is in $A_R(S^2,\text{id})$, i.e. there is no relation between W_S and W_N. Since the number of one-dimensional representations in $\text{Rep}(R)$ is equal to $n=|R_S|=|R_N|$, we obtain a proof. □ □

Remark 13.1. We explain for the reason why we prove the isomorphism of Theorem D only for R's appearing in Theorem A, B. In the proof of Theorem D, existence of $(G_\chi)_q$-extensions of U for all q or $(G_\chi)_{d^i}$-extensions of U for all i is critical according to R. But, such existence is not guaranteed if R_q or R_{d^i} is isomorphic to D_{2m} for some m by [CMS, Corollary 3.5.(2)], and almost all R's appearing in Theorem C satisfy that R_q or R_{d^i} is isomorphic to D_{2m} for some q or i according to R. So, we can not obtain the isomorphism for such R's. The inextensibility does not happen in dealing with equivariant vector bundles over circle. □

In the below, we use the notation $\mathbf{W}$ to denote an element in $A_R(S^2,\text{id})$ with one-dimensional entries. We would calculate Chern classes of line bundles in $\text{Vect}_R(S^2)$. Especially, we would calculate k_0 of Theorem A which is dependent on $\mathbf{W}$. Denote it by $k_0(\mathbf{W})$ to stress its dependency. By Theorem A, $k_0(\mathbf{W})$ is determined up to $l_R\cdot\mathbb{Z}$, i.e. $k_0(\mathbf{W})$ lives in $\mathbb{Z}_{l_R}$. More precisely, Theorem A says that $k_0(\mathbf{W})$ is congruent modulo l_R to $c_1(L)$ for any line bundle $L\in p_{\text{vect}}^{-1}(\mathbf{W})$. So, we will calculate $c_1(L) \pmod{l_R}$ for one bundle L in $p_{\text{vect}}^{-1}(\mathbf{W})$. In doing so, $c_1(L)$ is expressed by using $\mathbf{W}=(L_q)_{q\in B}$ or $\mathbf{W}=(L_{d^i})_{i\in I^+}$ according to R. When $n\in\mathbb{N}$ is understood, put $\xi_0=\exp(2\pi\sqrt{-1}/n)$, and let $\Omega(l)$ for $l\in\mathbb{Z}_n$ be the one-dimensional $\mathbb{Z}_n$-representation satisfying $a_n\cdot v=\xi_0^l v$ for each $v\in\mathbb{C}$. Then, we have the following well-known result:

Lemma 13.2. *Assume that $R=\mathbb{Z}_n$. For any line bundle L in $\text{Vect}_R(S^2)$, if $L_q\cong\Omega(l_q)$ for $q\in B$, $l_q\in\mathbb{Z}_n$, then $c_1(L)\equiv l_N-l_S \pmod{n}$.*

We obtain similar results for cases when $R=\text{D}_n$, $\mathbb{Z}_n\times Z$ with odd n, or $\langle -a_n\rangle$ with even $n/2$.

Lemma 13.3. *Assume that $R=\text{D}_n$. For any line bundle L in $\text{Vect}_R(S^2)$, if $L_q\cong\Omega(l_q)$ for $q\in B$, $l_q\in\mathbb{Z}_n$, then $l_N\equiv -l_S \pmod{n}$, and $c_1(L)$ is congruent modulo $2n$ to*

$$\begin{cases} -2l_S & \text{if } L_{d^0}\cong L_{d^1}, \\ -2l_S+n & \text{if } L_{d^0}\ncong L_{d^1}. \end{cases}$$

Proof. The first statement easily follows from the relation $ba_nb^{-1}=a_n^{-1}$. To prove the second statement, we would construct line bundles L' in $\text{Vect}_R(S^2)$ such that $L'_q\cong\Omega(l_q)$. Pick the R-bundle V_B over B such that

$$a_n\cdot v=\xi_0^{l_q}v \quad\text{and}\quad b\cdot v=v$$

for $q\in B$ and $v\in V_q=\mathbb{C}$. And, define F_{V_B} as $\amalg_{q\in B}\,|\bar{\mathcal{K}}_{R,q}|\times V_q$ such that

$$g\cdot(\bar{x},v)=(g\cdot\bar{x},g\cdot v)$$

for $g\in R$, $q\in B$, $\bar{x}\in|\bar{\mathcal{K}}_{R,q}|$, $v\in V_q$. We calculate $\mathcal{A}_{\bar{x}}$. Let φ_0 be the element id in $\text{Iso}(V_S,V_N)=\text{Iso}(\mathbb{C})$. Then, we can show the following:

1. $\mathcal{A}^0_{\bar{v}^0_S} = \{\varphi_0,\ -\varphi_0\}$. And, $\varphi_0, -\varphi_0$ determine $\langle b\rangle$-representations $\Omega(0)$, $\Omega(1)$, respectively.
2. $\mathcal{A}^0_{b(\bar{e}^0_S)} = \{\xi_0^{-l_S}\varphi_0,\ -\xi_0^{-l_S}\varphi_0\}$. And, $\xi_0^{-l_S}\varphi_0$, $-\xi_0^{-l_S}\varphi_0$ determine $\langle a_n b\rangle$-representations $\Omega(0)$, $\Omega(1)$, respectively.
3. $\mathcal{A}^0_{\bar{x}} = \mathrm{Iso}(V_S, V_N) = \mathrm{Iso}(\mathbb{C})$ for each interior $\bar{x}$ of $[\bar{v}^0_S, b(\bar{e}^0_S)]$.

By using the parametrization on $|\bar{\mathcal{L}}_{R,S}|$, we can define two equivariant clutching maps Φ and Φ' with respect to V_B which satisfy

$$\Phi(t) = \exp\Big(- \frac{4\pi t\ l_S\sqrt{-1}}{n} \Big)\varphi_0 \quad \text{and} \quad \Phi'(t) = \exp\Big(- \frac{4\pi t\ \big(l_S + \frac{n}{2}\big)\sqrt{-1}}{n} \Big)\varphi_0$$

for $t \in [0, n]$. And, we can show the following:

1. $L'_{d^{-1}} \cong \Omega(l_S)$, $L'_{d^0} \cong \Omega(0)$, $L'_{d^1} \cong \Omega(0)$, and $c_1(L') \equiv -2l_S \pmod{2n}$ for $L' = F_{V_B}/\Phi$,
2. $L'_{d^{-1}} \cong \Omega(l_S)$, $L'_{d^0} \cong \Omega(1)$, $L'_{d^1} \cong \Omega(1)$, and $c_1(L') \equiv -2l_S \pmod{2n}$ for $L' = F_{V_B}/-\Phi$,
3. $L'_{d^{-1}} \cong \Omega(l_S)$, $L'_{d^0} \cong \Omega(0)$, $L'_{d^1} \cong \Omega(1)$, and $c_1(L') \equiv -2l_S + n \pmod{2n}$ for $L' = F_{V_B}/\Phi'$,
4. $L'_{d^{-1}} \cong \Omega(l_S)$, $L'_{d^0} \cong \Omega(1)$, $L'_{d^1} \cong \Omega(0)$, and $c_1(L') \equiv -2l_S + n \pmod{2n}$ for $L' = F_{V_B}/-\Phi'$.

Images of these four line bundles through p_{vect} are four $\mathbf{W}$'s in $A_R(S^2, \mathrm{id})$ whose $W_{d^{-1}}$-entry is isomorphic to $\Omega(l_S)$. Therefore, we obtain a proof. □ □

Remark 13.4. We explain for how to calculate $k_0(\mathbf{W})$ for cases when $R = \mathrm{T}$, O, I. Let $\mathbf{W} = p_{\mathrm{vect}}(L)$ for some line bundle L in $\mathrm{Vect}_R(S^2)$. Then, it suffices to calculate $c_1(L) \pmod{l_R}$. Note $l_R = |R|$ in these cases. For a 2-Sylow subgroup P of R, observe that the restricted P-action on S^2 is conjugate to D_m for some m. Then, we can calculate $c_1(L) \pmod{2m}$ by applying Lemma 13.3 to $\mathrm{res}^R_P L$ where $|\,\mathrm{D}_m\,| = 2m$. For other prime number p dividing $|R|$ and a p-Sylow subgroup P of R, observe that the restricted P-action on S^2 is conjugate to $\mathbb{Z}_p$, and we can calculate $c_1(L) \pmod{p}$ by applying Lemma 13.2 to $\mathrm{res}^R_P L$ where $|\mathbb{Z}_p| = p$. So, we can calculate $c_1(L) \pmod{l_R}$ by Chinese Remainder Theorem because $l_R = |R|$. □

Lemma 13.5. *Assume that $R = \mathbb{Z}_n \times Z$ with odd n. For any line bundle L in $\mathrm{Vect}_R(S^2)$, if $L_q \cong \Omega(l_q)$ for $q \in B$, $l_q \in \mathbb{Z}_n$, then $l_N \equiv l_S \pmod{n}$, and $c_1(L)$ is trivial.*

Proof. The first statement easily follows because a_n and $-\mathrm{id}$ commute in R. To prove the second statement, we would construct a line bundle L' in $\mathrm{Vect}_R(S^2)$ such that $L'_q \cong \Omega(l_q)$. Pick the R-bundle V_B over B such that

$$a_n \cdot v = \xi_0^{l_q} v \quad \text{and} \quad -\mathrm{id} \cdot v = v$$

for $q \in B$ and $v \in V_q = \mathbb{C}$. And, define F_{V_B} as $\amalg_{q\in B}\ |\bar{\mathcal{K}}_{R,q}| \times V_q$ such that

$$g \cdot (\bar{x}, v) = (g \cdot \bar{x}, g \cdot v)$$

for $g \in R$, $q \in B$, $\bar{x} \in |\bar{\mathcal{K}}_{R,q}|$, $v \in V_q$. Then, we can define the equivariant clutching map Φ with respect to V_B which satisfies $\Phi(\bar{x}) = \mathrm{id} \in \mathrm{Iso}(V_S, V_N) = \mathrm{Iso}(\mathbb{C})$ for each $\bar{x} \in \bar{\mathcal{L}}_{R,S}$. For $L' = F_{V_B}/\Phi$, the Chern class $c_1(L')$ is trivial. Since two equivariant vector bundles in $\mathrm{Vect}_R(S^2)$ with the same isotropy representation at each d^i have the same Chern class by Theorem B, we obtain a proof. □ □

Lemma 13.6. *Assume that $R = \langle -a_n\rangle$ with even $n/2$. For any line bundle L in $\mathrm{Vect}_R(S^2)$, if $L_q \cong \Omega(l_q)$ for $q \in B$, $l_q \in \mathbb{Z}_{n/2}$, then $l_N \equiv l_S \pmod{n/2}$, and $c_1(L)$ is trivial.*

Proof. Similarly to Lemma 13.3 and Lemma 13.5, the first statement can be proved by using the fact that R is a cyclic group. To prove the second statement, we would construct a line bundle L' in $\mathrm{Vect}_R(S^2)$ such that $L'_q \cong \Omega(l_q)$. Put $c_0 = \exp\left(\frac{2\pi l_S\sqrt{-1}}{n}\right)$ so that $c_0^2 = \xi_0^{l_S}$ where $\xi_0 = \exp\left(\frac{2\pi\sqrt{-1}}{n/2}\right)$. Pick the R-bundle V_B over B such that

$$-a_n \cdot v = c_0 v$$

for $q \in B$ and $v \in V_q = \mathbb{C}$. And, define F_{V_B} as $\amalg_{q\in B} |\bar{\mathcal{K}}_{R,q}| \times V_q$ such that

$$g \cdot (\bar{x}, v) = (g \cdot \bar{x}, g \cdot v)$$

for $g \in R$, $q \in B$, $\bar{x} \in |\bar{\mathcal{K}}_{R,q}|$, $v \in V_q$. Then, the remaining is the same with Lemma 13.5. □ □

14. Appendix: representation extension

Let N_0 and N_2 be compact Lie groups such that $N_0 \lhd N_2$ and $N_2/N_0 \cong \mathbb{Z}_m$. Let a_0 be a fixed generator of N_2/N_0, and let $\Omega(l)$ be the representation defined by

$$N_2/N_0 \times \mathbb{C} \to \mathbb{C}, \quad (a_0, z) \mapsto \exp(2\pi l\sqrt{-1}/m)z$$

for $l \in \mathbb{Z}_m$. We also consider $\Omega(l)$ to be an N_2-representation via the projection $N_2 \to N_2/N_0$. Then, we obtain the following result from [CMS]:

Theorem 14.1. *For $U \in \mathrm{Irr}(N_0)$, if the character of U is fixed by N_2, then there exists an N_2-extension of U. If $\bar{U}$ is an N_2-extension of U, then the number of mutually nonisomorphic N_2-extensions of U is m and they are $\bar{U}\otimes\Omega(l)$ for $l \in \mathbb{Z}_m$.*

Proof. By [CMS, Theorem 3.2.], U has m mutually nonisomorphic N_2-extensions. Call one of them $\bar{U}$. By [CMS, Proposition 3.1.] and its proof, each extension of U is expressed as $\bar{U} \otimes \Omega(l)$ for some $l \in \mathbb{Z}_m$. □ □

Corollary 14.2. *Let U be an irreducible N_0-representation whose character is fixed by N_2, and $\bar{U}$ be an N_2-extension of U. If W be an N_2-representation such that $\mathrm{res}^{N_2}_{N_0} W$ is U-isotypical, W is a direct sum of $\bar{U} \otimes \Omega(l)$'s.*

Proof. First, we prove that the induced representation $\mathrm{ind}^{N_2}_{N_0} U$ is isomorphic to the direct sum $\oplus_{l\in\mathbb{Z}_m}(\bar{U}\otimes\Omega(l))$. By Frobenius reciprocity, $\mathrm{Hom}_{N_2}(\bar{U}\otimes\Omega(l), \mathrm{ind}^{N_2}_{N_0} U) \cong \mathrm{Hom}_{N_0}(\mathrm{res}^{N_2}_{N_0} \bar{U} \otimes \Omega(l), U)$ is one-dimensional, and this means that each $\bar{U} \otimes \Omega(l)$ for $l \in \mathbb{Z}_m$ is a subrepresentation of $\mathrm{ind}^{N_2}_{N_0} U$ by Schur's Lemma. So, $\mathrm{ind}^{N_2}_{N_0} U$ is isomorphic to the direct sum $\oplus_{l\in\mathbb{Z}_m}(\bar{U} \otimes \Omega(l))$.

We may assume that W is irreducible. We only have to show that W is one of $\bar{U} \otimes \Omega(l)$'s. Since $\mathrm{res}^{N_2}_{N_0} W$ is U-isotypical, $\mathrm{res}^{N_2}_{N_0} W \cong lU$ for some integer l. By Frobenius reciprocity, $\mathrm{Hom}_{N_2}(W, \mathrm{ind}^{N_2}_{N_0} U) \cong \mathrm{Hom}_{N_0}(\mathrm{res}^{N_2}_{N_0} W, U)$. Since $\mathrm{res}^{N_2}_{N_0} W$ is isomorphic to lU, we obtain that $\mathrm{Hom}_{N_0}(\mathrm{res}^{N_2}_{N_0} W, U)$ is l-dimensional by Schur's Lemma. But, since $\bar{U}\otimes\Omega(l)$'s are all different and W is irreducible, Schur's Lemma says that $\mathrm{Hom}_{N_2}(W, \mathrm{ind}^{N_2}_{N_0} U)$ is at most one-dimensional, i.e. $l \leq 1$. Therefore, l is equal to 1, and this gives a proof. □ □

References

[At] M. F. Atiyah, *K-Theory*, Addison-Wesley, 1989.

[B] G. E. Bredon, *Introduction to compact transformation groups*, Academic Press, New York and London, 1972.

[BD] T. Bröcker, T. tom Dieck, *Representations of compact Lie groups*, Graduate Texts in Mathematics, 98, Springer-Verlag, New York, 1985.

[CKMS] J.-H. Cho, S. S. Kim, M. Masuda, D. Y. Suh, *Classification of Equivariant Complex Vector Bundles over a Circle*, J. Math. Kyoto Univ. **41** (2001), 517-534.

[CK] A. Constantin, B. Kolev, *The theorem of Kerekjarto on periodic homeomorphisms of the disk and the sphere*, Enseign. Math. (2) **40** (1994), 193–204.

[CMS] J.-H. Cho, M. Masuda, D. Y. Suh, *Extending representations of H to G with discrete G/H*, J. Korean Math. Soc. **43** (2006), 29–43.

[G] A. Grothendieck, *Sur la classification des fibres holomorphes sur la sphere de Riemann*, Am. J. Math. **(**79) (1957), 121–138 .

[Ki1] M. K. Kim, *Classification of equivariant vector bundles over two-torus*, arXiv:1005.0682v2.

[Ki2] M. K. Kim, *Classification of equivariant vector bundles over real projective plane*, preprint.

[Ki3] M. K. Kim, *Classification of equivariant vector bundles over Klein bottle*, in preparation.

[Ko] B. Kolev, *Sous-groupes compacts d'homeomorphismes de la sphere*, Enseign. Math. (2) 52 (2006), no. 3-4, 193–214.

[P] F. P. Peterson, *Some remarks on Chern classes*, Ann. Math. (2) **69** (1959), 414–420.

[R] E. G. Rees, *Notes on geometry*, Universitext, Springer-Verlag, Berlin-New York, 1983.

[S] G. Segal, *Equivariant K-theory*, Inst. Hautes Études Sci. Publ. Math. **34** (1968), 129–151.

Department of Mathematics Education, Gyeongin National University of Education, San 59-12, Gyesan-dong, Gyeyang-gu, Incheon, 407-753, Korea

E-mail address: `mkkim@kias.re.kr`